
References

- 1 Behrman RE, Vaughan VC, Nelson WF (eds): Nelson Textbook of Pediatrics, ed 13. Philadelphia, Saunders, 1987, appendix.
- 2 Brohee D, Vanhaeverbeek M, Kennes B, Neve P: Leukocyte and lymphocyte subsets after a short pharmacological stress by intravenous epinephrine and hydrocortisone in healthy humans. *Int J Neurosci* 1990;53:53-62.
- 3 Gabriel A, Urhausen A, Kindermann W: Mobilization of circulating leucocyte and lymphocyte subpopulations during and after short, anaerobic exercise. *Eur J Appl Physiol Occup Physiol* 1992;65:164-170.
- 4 McCarthy DA, Macdonald IA, Shaker HA, Hart P, Georgiannos S, Deeks J, Wade AJ: Changes in the leucocyte count during and after intensive exercise. *Eur J Appl Physiol Occup Physiol* 1992;64:518-522.
- 5 Xanthou M: Leucocyte blood picture in healthy full-term and premature babies during neonatal period. *Arch Dis Child* 1970;45:242-249.
- 6 Nikischin W, Weisner D, Oldigs HD: Perinatal glucose metabolism as an indicator for stress and hypoxia during different forms of delivery. *J Perinat Med* 1990;18:209-213.

Announcement

14th International Congress: The Fetus as a Patient

Amsterdam, May 12-16, 1998

Venue: RAI Congress Centre.

Organization: The International Society The Fetus as a Patient.

Chairman:

Prof. Dr. H.P. van Geijn, chairman of the local organizing committee
Department of Obstetrics and Gynecology
University Hospital, Vrije Universiteit
NL-Amsterdam (The Netherlands)

Tel. +31 20 444 22 27, Fax +31 20 444 48 11

Information:

Congress Office SCEM
Ms. M. de Haan
PO Box 21
NL-4196 ZG Tricht (The Netherlands)

Tel. +31 345 57 66 42, Fax +31 345 57 17 81, E-mail SCEM@SCEM.NL.