Reply

P.M. Yuen P.S. Ng D.S. Sahota

We thank Dr. Mahajan and colleagues for their comments on our study [1]. We agree with them that misoprostol is a useful uterotonic drug in developing countries for the prevention of postpartum hemorrhage (PPH). We also concur on their comment on the estimation of blood loss. Clinical estimation of blood loss underestimates true blood loss, whereas the weighing method described in the WHO study could overestimate it as the presence of amniotic fluid could not be excluded.

We have repeatedly stressed this point in our previous studies related to the use of misoprostol in the management of PPH [1, 2]. More importantly, the amount of clinical estimation of blood loss does not reflect the actual hemoglobin change. Over 25% of the women who were defined as having PPH had a peripartum hemoglobin drop of over 20%. That is the reason why we chose to use the peripartum change in hemoglobin level as our primary outcome measure.

References

- 1 Ng PS, Lai CY, Sahota DS, Yuen PM: A double-blind randomized controlled trial of oral misoprostol and intramuscular syntometrine in the management of the third stage of labor. *Gynecol Obstet Invest* 2007;63:55–60.
- 2 Ng PS, Chan ASM, Sin WK, Tang LCH, Cheung KB, Yuen PM: A multicentre randomised controlled trial of oral misoprostol and i.m. syntometrine in the management of the third stage of labour. *Hum Reprod* 2001; 16:31–35.

Dr. Pong Mo Yuen
Department of Obstetrics and Gynaecology
Shatin, N.T. Hong Kong, SAR (China)
Tel. +852 2632 2810, Fax +852 2636 0008
E-Mail pmyuen@cuhk.edu.hk