Preface

Based on the plans formulated by the Japan Research Committee on Long-Term Treatment of Parkinson’s Disease, the third symposium was designed and held in Tokyo on April 23, 1988.

In the morning, the interim report on a 2-year bromocriptine treatment of Parkinson’s disease in Japan was made by Dr. I. Kanazawa. Then, Dr. B.P. Drayer presented beautiful magnetic resonance imaging in parkinsonian and many other movement disorders.

In the afternoon session, the pathophysiological features of Parkinson’s disease were revealed by four well-known authorities. Dr. R.G. Lee presented a critical review of current knowledge concerning the physiological basis of rigidity. Dr. N. Yanagisawa showed that recognition of a light stimulus and initiation of voluntary movements after the stimulus became delayed in patients with Parkinson’s disease. Dr. H. Narabayashi demonstrated the neurosurgical evidence that Vim (nucleus ventralis intermedius) in the thalamus was the key-member of the circuit producing tremor. Dr. H.J. Freund suggested that Parkinson’s disease was characterized by the interference with elementary motor functions such as initiation and execution.

A large number of Japanese clinical neuroscientists attended this symposium and participated actively in the discussion.

I would like to express my sincere gratitude to all the distinguished speakers who gave us excellent presentations. Furthermore, I wish to thank Sandoz Pharmaceutical Company and the staff for providing financial support and their organizational skills and sustained efforts, which made this symposium possible.

Takao Nakanishi