Introduction

'The goal of life is to live in agreement with nature'
Zeno (335–264 BC)

Humanity faces two important challenges when dealing with energy and the environment, issues that are intimately connected: (1) How can all the goods that society needs and will need be produced using energetically efficient and environmentally friendly processes? (2) How can the dependence on fossil energy sources be reduced and the environmental impact of human activities be minimized?

All human activities on Earth produce some type of byproduct. Under normal conditions these byproducts, some known as pollutants, are recycled into the environment. In fact, natural systems have the ability to assimilate some pollutants into the environment and correct ecological imbalances if given enough time. However, if a persistent overload of a pollutant or if the pollutant cannot be integrated into the environment, then the environment has little chance to 'self-clean' itself. This accumulation is especially critical for the increasing number of novel industrial compounds called xenobiotics, which are rarely found in nature.

Large quantities of organic compounds are manufactured and used by industries, agriculture and municipalities. These manmade organic compounds are of tremendous concern. Modern economic activities require transportation and storage of materials used in manufacturing, processing, and construction. Along the way, some of this material can be lost through spillage, leakage, or improper handling. Pesticides, fertilizers, and herbicides are agricultural sources of environmental contamination. Agricultural contamination sources are varied and numerous: spillage of fertilizers and pesticides during handling

and runoff from the loading and washing of pesticide sprayers or other application equipment are examples. Residential wastewater systems can be sources of many categories of contaminants including organic compounds. Improper storage or disposal of household chemicals such as paints, synthetic detergents, solvents, oils, medicines, disinfectants, pool chemicals, pesticides, batteries, gasoline and diesel fuel can contribute to contamination of the environment.

Environmental biotechnology started in the early 20th century with the use of activated sludge and anaerobic digestion; however, the introduction of new technologies from modern microbiology, enzymology and molecular biology has enabled engineers and scientist to tackle the more contemporary environmental problems such as detoxification of hazardous chemicals. Using tools from metabolic and protein engineering, research efforts have been focused on designed microbes and enzymes to function as environmental biocatalysts, in which certain desirable properties are brought together with the aim of optimizing the rates and specificities of biodegradation pathways.

The aim of this symposium is to present an overview of different biological and biochemical transformations of pollutant compounds including xenobiotic chemicals. Biotransformations of aromatic compounds, polychlorinated chemicals, pesticides, dioxins and fuel additives are analyzed and discussed. Critical reviews at three different levels – bioremediation processes, microbial and plant metabolic pathways and enzymatic transformations – are presented.

Hypersaline environments are important for both surface extension and ecological significance. As in all other ecosystems, they are impacted by pollution. Due to their reduced microbial diversity when compared with non-extreme environments, pollutant degradation becomes critical in these environments. Conventional non-extremophilic microorganisms are unable to efficiently perform the removal of organic pollutants at high salt concentrations. A literature survey by LeBorgne and co-workers indicates that both the moderately halophilic bacteria and the extremely halophilic archaea have a broader catabolic versatility and capability than previously thought. A diversity of contaminating compounds is susceptible to be degraded by halotolerant and halophilic bacteria. Nevertheless, significant research efforts are still necessary in order to estimate the true potential of these microorganisms to be applied in environmental processes and in the remediation of contaminated hypersaline ecosystems.

Julia Foght reviews the anaerobic biodegradation of aromatic hydrocarbons, including the carcinogenic and mutagenic polycyclic aromatic hydrocarbons (PAHs). Whereas aerobic biodegradation has been well studied for decades, anaerobic hydrocarbon biodegradation is a nascent field undergoing rapid shifts in concept and scope. A critical review of the known pathways, enzymes, genes, and terminal electron acceptors during the anaerobic biodegradation of aromatic hydrocarbons both in situ and in laboratory studies is presented. This information is essential to understand and manipulate biodegradation processes for potential application in situ.

Polychlorinated biphenyls (PCBs) represent a class of compounds which have been used since 1929 for various industrial and commercial purposes. They can serve as hydraulic fluids, plasticizers and fire retardants. PCBs are highly carcinogenic and are able to produce serious effects on the immune, reproductive, nervous and endocrine systems. Pieper and Seeger review the reductive dehalogenation of PCBs and the metabolic pathway for aerobic PCB-degradation.

The biphenyl dioxygenases are of crucial importance for the successful metabolism of PCBs. Vézina's group examines the sequence diversity of biphenyl dioxygenases, from cultured bacteria and soil microflora, enzymes that catalyze stereospecific dioxygenation of biphenyl and its analogues. The phylogenetic analyses included cultured bacteria from different ecological niches and uncultured bacteria from PCB-contaminated soils; results show that most biphenyl dioxygenases are associated with three clusters. Few diverge from these branch-

es, showing a high degree of diversity for these enzymes in nature. The authors suggest that PCB-degrading soil bacteria have evolved to optimize their PCB-degrading capacity.

Dioxins are two-ring aromatic compounds containing one to eight chlorine atoms, (polychlorinated dibenzo-*p*-dioxins and dibenzofurans). Dioxins are among the most problematic environmental pollutants because of their recalcitrance, lipid solubility, and toxicity. Doubtless, dioxins are some of the most dangerous chemicals ever synthesized. The origin of the public health risk associated with dioxins occurs because they are immunotoxic, developmental disruptors, neurotoxic and carcinogenic compounds, affecting skin, liver, and gastrointestinal systems. Large amounts of highly toxic dioxins enter into the environment as a result of events such as the Vietnam War and the Seveso industrial accident. Clean-up of dioxin-contaminated sites has been difficult due to their toxicity and biological recalcitrance. Yoon-Seok Chang reviews the recent developments in microbial degradation and biotransformation of dioxins. Dioxin degradation has been reported in fungi, where several oxidative enzymes are responsible for dioxin degradation. These enzymes include cytochrome P-450, lignin peroxidase, manganese peroxidase, and laccase. However, successful bioremediation techniques for field-scale application are still under development.

Ligninolytic fungi are good candidates for xenobiotic decontamination through biotechnological processes. These organisms are able to metabolize chemically diverse molecules, including chlorinated phenols, pesticides, PCBs, dioxins, organophosphorus compounds, nitrotoluenes, chloranilines, dyes, and other compounds of environmental concern. Their extracellular enzymatic systems, which are involved in lignin degradation, consist mainly of oxidative enzymes: laccases, lignin peroxidases, manganese peroxidases, and versatile peroxidases. These peroxidases are able to oxidize a wide range of substrates with high redox potential including various xenobiotic compounds such as industrial dyes, polycyclic aromatic hydrocarbons, pesticides, dioxins, chlorophenols and explosives. Ayala and coworkers discuss the scientific challenges for the use of these enzymes in large-scale environmental processes. Three main protein engineering challenges have been identified: (1) enhancement of stability in the presence of hydrogen peroxide, (2) increase of the enzyme redox potential in order to widen the substrate range, and (3) development of heterologous expression for industrial production. Research efforts, mainly using DNA recombinant techniques, are needed

to solve these challenges, designing an environmental biocatalyst able to oxidize a wide range of xenobiotics or pollutants.

Bioremediation of several compounds can be enhanced by expressing bacterial (*Vitreoscilla*) hemoglobin in heterologous hosts. Stark's group present experimental evidence of pollutant degradation with different bacterial strains that have been genetically engineered. They demonstrated that the expression of *Vitreoscilla* hemoglobin can enhance the biodegradation of several aromatic and organophosphorus compounds. Experiments were carried out under conditions that closely resemble those in field applications, such as soil columns, continuous flow reactors and membrane bioreactors. In most cases, particularly in bioreactors, the advantages due to hemoglobin expression were greater under conditions of limited aeration or hypoxic conditions.

In recent years, concern about the environmental presence of methyl *tert*-butyl ether (MTBE), a gasoline additive, has increased. The massive use of this fuel additive, combined with its mobility in water and its low biodegradability, makes it an important and persistent pollutant of groundwater. Maciel and colleagues report on the capacity of *Mycobacterium austroafricanum* to degrade MTBE and its performance in a bioreactor for the bioremediation of contaminated groundwater. The bioreactor was monitored using molecular tools and showed removal efficiencies higher than 99%.

The well-known toxic cyanide ion is also a pollutant. Beyond natural sources, millions of tons of hydrogen cyanide are produced worldwide. Hydrogen cyanide and

cyanide salts are widely used in the synthesis of organic and inorganic chemicals. They are also used in ore extracting processes for the recovery of gold and silver, electroplating, case-hardening of steel, base metal flotation, metal degreasing, dyeing, printing, and photography. There are four general enzymatic pathways for the biodegradation of cyanide involving hydrolytic, oxidative, reductive, and substitution/transfer reactions. Cipollone and coworkers review the biochemical and physiological properties of microbial rhodanese, the best characterized thiosulfate:cyanide sulfurtransferase, in light of its importance in cyanide detoxification by the cyanogenic bacterium *Pseudomonas aeruginosa*. Challenges for the application of a rhodanese-based cellular system in cyanide bioremediation processes are critically discussed.

Finally, the contribution of Kawahigashi and his group shows a vast spectrum of possible environmental applications of transgenic plants. Phytoremediation has successfully detoxified soils contaminated with heavy metals. Plants may enhance the remediation of xenobiotic-contaminated soils. The introduction of genes encoding cytochrome P450 monooxygenases, which are involved in xenobiotic degradation, produce transgenic plants more tolerant to various herbicides and are able to remove atrazine and metolachlor from soil. Transgenic plants appear to have a huge potential for remediation processes.

Society has an imperative goal, even more important than that of preserving humanity: to reestablish Nature's equilibrium. Environmental Biotechnology can certainly be of help.

Rafael Vazquez-Duhalt