Reply

Pravin J. Gupta

Fine Morning Hospital and Research Center, Laxminagar, Nagpur, India

Dear Sir,

I respectfully acknowledge the positive appraisal of my paper by Prof. Francesco F. di Mola and others. My paper describes the ill effects of red chilies after anal fissure surgery [1]. As he has correctly pointed out, the aim of my study was based on our clinical assumption that consumption of red chili increases pain, anal burning and frequency of stool after anal fissure surgery. I have also carried out a similar study in patients during the post-hemorrhoidectomy period and found that consumption of chilies during this period increases postoperative pain, analgesic consumption, bleeding and pruritus [2].

I have mentioned in my discussion that capsaicin, the pungent principle of hot pepper, has the ability to excite and later defunctionalize a subset of primary afferent neurons. It has also been found that chilies cause rectal hyperalgesia, and a significant number of mucosal inflammatory cells and an increase in BrdU-immunoreactive nuclei were detected following mucosal exposure to capsaicin in the colon. I indeed failed to mention the very important findings of di Mola et al. [3] regarding activation of the transient receptor potential vanilloid 1 (TRPV1) by capsaicin. TRP channels are now included among the targets of peptide toxins, showing that chili peppers avert predators by activating TRP channels on sensory nerve fibers to elicit pain and inflammation [4]. TRPV1 might also play an important role in initiation and maintenance of persistent visceral hypersensitivity [5].

Nonetheless, an experimental study has found that capsaicin inhibits catecholamine secretion and synthesis via suppression of Na^+ and Ca^{2+} influx through a vanilloid receptor-independent pathway [6]. Another study on murine Peyer's patch (PP) cells has shown that capsaicin modulates T-cell-immune responses, and its immunomodulatory effects on murine PP cells is partly due to both a TRPV1-dependent and -independent pathway [7]. Capsaicin has been shown to activate and desensitize C-fiber and A-delta sensory nerve fibers. Stimulation causes an acute neurogenic response including vasodilatation, plasma extravasation and hypersensitivity [8], but in another study it was shown that capsaicin produces a dose-dependent vasoconstrictor effect in the mouse knee joint via TRPV1 receptor activation [9].

Looking toward these variable findings with regards to the action of capsaicin, it would be interesting to look out for more conclusive studies about the action of chili pepper on rectum and anal canal.

References

- 1 Gupta PJ: Red hot chilli consumption is harmful in patients operated for anal fissure – a randomized, double-blind, controlled study. *Dig Surg* 2007;24:354–357.
- 2 Gupta PJ: Effect of red chili consumption on postoperative symptoms during the post-hemorrhoidectomy period: randomized, double-blind, controlled study. *World J Surg* 2007;31:1822–1826.

- 3 di Mola FF, Friess H, Königer J, Selvaggi F, Esposito I, Büchler MW, di Sebastiano P: Haemorrhoids and transient receptor potential vanilloid 1. *Gut* 2006;55:1665–1666.
- 4 Siemens J, Zhou S, Piskrowski R, Nikai T, Lumpkin EA, Basbaum AI, King D, Julius D: Spider toxins activate the capsaicin receptor to produce inflammatory pain. *Nature* 2006;444:208–212.
- 5 Winston J, Shenoy M, Medley D, Naniwadekar A, Pasricha PJ: The vanilloid receptor initiates and maintains colonic hypersensitivity induced by neonatal colon irritation in rats. *Gastroenterology* 2007;132:615–627.
- 6 Takahashi K, Toyohira Y, Ueno S, Tsutsui M, Yanagihara N: Capsaicin inhibits catecholamine secretion and synthesis by blocking Na^+ and Ca^{2+} influx through a vanilloid receptor-independent pathway in bovine adrenal medullary cells. *Naunyn Schmiedeberg Arch Pharmacol* 2006;374:107–116.
- 7 Takano F, Yamaguchi M, Takada S, Shoda S, Yahagi N, Takahashi T, Ohta T: Capsicum ethanol extracts and capsaicin enhance interleukin-2 and interferon- γ production in cultured murine Peyer's patch cells ex vivo. *Life Sci* 2007;80:1553–1563.
- 8 Shimizu T: Structure and function of nociceptive neuronal receptors. *Nippon Rinsho* 2005;63:1713–1719.
- 9 Keeble JE, Brain SD: Capsaicin-induced vasoconstriction in the mouse knee joint: a study using TRPV1 knockout mice. *Neurosci Lett* 2006;401:55–58.

Gupta Nursing Home
D/9, Laxminagar, Nagpur (India)
Tel. +91 712 223 1047
E-Mail drpgj@yahoo.co.in