Neuroendocrine-Immune Interactions in Rheumatoid Arthritis: Mechanisms of Glucocorticoid Resistance

Marni N. Silverman Esther M. Sternberg

Section on Neuroendocrine Immunology and Behavior, National Institute of Mental Health, National Institutes of Health, Rockville, Md., USA

Key Words

Glucocorticoids · Glucocorticoid resistance · Corticosteroid binding globulin · Multidrug resistance transporter · 11 β -Hydroxysteroid dehydrogenase · Glucocorticoid receptor · Cytokines · Inflammation · Autoimmune disease

Abstract

Rheumatoid arthritis (RA) is characterized by chronic inflammation of the synovial membrane, leading to joint destruction. Many autoimmune diseases and disease states of chronic inflammation are accompanied by alterations in the complex interactions between the endocrine, nervous and immune systems. Glucocorticoids, an end product of the hypothalamic-pituitary-adrenal axis, are a mainstay treatment for many autoimmune diseases, including RA, because of their potent anti-inflammatory action. However, about 30% of patients with RA fail to respond to steroid therapy. There are various mechanisms that may contribute to the development of glucocorticoid resistance in inflammatory disorders, which will be the subject of this review. In addition, glucocorticoid resistance may be a contributing factor in the development of inflammatory/autoimmune diseases themselves. Therefore, further elucidation of these mechanisms will reveal new targets for therapeutic intervention in the treatment of RA.

Copyright © 2008 S. Karger AG, Basel

Introduction

In the 1940s, Philip Hench discovered that patients with autoimmune disorders, such as rheumatoid arthritis (RA), produced an endogenous substance under 'stressful' conditions that had anti-inflammatory/immunosuppressive properties and hence, ameliorated the symptoms of the autoimmune disease. Isolation and characterization of this endogenous compound by Edward Kendall led to the discovery of the adrenal steroid, cortisone, which along with other glucocorticoids, has become a mainstay in the treatment of autoimmune and inflammatory diseases. Of note, Hench and Kendall shared the Nobel Prize in Medicine for their discovery in 1950.

Although the immunomodulatory effects of glucocorticoids initially were believed to be mediated by pharmacological rather than physiological concentrations, seminal work by Hugo Besedovsky and colleagues in the 1970s and 1980s substantiated a physiological role for glucocorticoids in regulating immune responses. Besedovsky and del Rey [1] were also one of the first to demonstrate that immune system activity could influence the release of glucocorticoids. Since then, many others have provided evidence for the bidirectional communication between the neuroendocrine and immune systems [2, 3]. Today, in contrast to the traditional view of glucocorticoids as immunosuppressive hormones, they are more accurately

conceptualized as immunomodulatory hormones that can stimulate as well as suppress immune function, depending on glucocorticoid concentration, the type of immune response, the immune compartment and the cell type involved [4]. Once glucocorticoids are released into the general circulation, maintenance of appropriate glucocorticoid activity is accomplished by local tissue regulation of glucocorticoid availability and action by factors such as corticosteroid binding globulin (CBG), the multidrug resistance transporter (MDR), 11 β -hydroxysteroid dehydrogenase (11 β -HSD), and ultimately, the glucocorticoid receptor (GR). Altered expression and/or function of these factors have been found in RA and have become the subject of interest for the development of new therapeutic interventions.

Impaired HPA Axis Activity and Other Stress Systems in RA

Conditions of chronic inflammation are often associated with impaired anti-inflammatory stress response systems. In addition to absolute plasma levels of stress hormones (as exemplified below), it is important to study hormone levels in relation to inflammation or steroid hormone shifts, which can indicate preferential production of one hormone over another and influence the progression of chronic inflammatory diseases [5]. In RA, alterations in neuroendocrine function include inadequate ACTH and cortisol secretion (impaired HPA axis function), an increased sympathetic tone at rest but an inadequate response during stress, functional loss of synovial sympathetic nerve fibers concomitant with the presence of proinflammatory sensory fibers, a local beta-to-alpha adrenergic receptor shift and local uncoupling of cortisol and norepinephrine. In addition, a decrease in adrenal androgen production, such as DHEAS, DHEA and androstenedione, has been reported with a preference for cortisol production (although insufficient in relation to sustained inflammation). Taken together, these alterations in stress response systems lead to insufficient regulatory/anti-inflammatory responses to keep inflammation in check and may contribute to the pathology characteristic of RA [6].

Normal plasma levels of ACTH and cortisol in the presence of systemic inflammation is indicative of an inadequate HPA axis response to systemic inflammation. Interestingly, the circadian rhythm of cortisol in RA patients, whose disease activity is relatively low to moderate, is similar to that found in healthy controls [7, 8],

whereas a loss of circadian rhythm, as indicated by a flattened cortisol curve, has been observed in RA patients when the disease is very active [8]. Early morning peaks in plasma proinflammatory cytokine levels (e.g., TNF- α , IL-6) are shifted a couple of hours later (to around 5–7 a.m.) and are of greater amplitude and longer duration in RA patients versus controls. These circadian changes, despite the similarity of the circadian curves for serum cortisol (in amplitude and shape), also indicate inadequate cortisol secretion in relation to inflammation in RA. Elevated proinflammatory cytokines probably account for the increased morning stiffness (due to edema) and pain that are usually experienced by RA patients.

A defective HPA axis has also been associated with susceptibility to autoimmune/inflammatory diseases in several animal models. In regard to RA, the neuroendocrine differences observed between Lewis (LEW/N) and Fischer (F244/N) rat strains are a prime example. Lewis rats are highly susceptible, whereas Fischer rats are relatively resistant, to the development of a wide range of autoimmune diseases in response to a variety of proinflammatory/antigenic stimuli. Injection of group A streptococcal cell wall peptidoglycan polysaccharide (which mimics human RA) into inflammatory-susceptible Lewis rats produces a blunted HPA axis response (ACTH and corticosterone) compared to the exaggerated HPA axis response observed in inflammatory-resistant Fischer rats. Moreover, the replacement or removal of glucocorticoids in Lewis and Fischer rats, respectively, reverses their susceptibility to streptococcal cell wall-induced arthritis [9]. Dysfunctional HPA axis activity in Lewis rats has been shown to be due to the altered expression of multiple factors that regulate the HPA axis, including hypothalamic corticotrophin-releasing hormone, pituitary proopiomelanocortin, CBG and GR [10]. Local factors regulating glucocorticoid bioavailability and actions are discussed below.

Local Factors Regulating Glucocorticoid Bioavailability and Action

Factors Regulating Glucocorticoid Bioavailability

While circulating levels of glucocorticoid hormones are relevant to steroid action, at the cellular level, activity of glucocorticoids is determined by local factors that regulate the access of free hormone to its receptor. Such factors include CBG, the MDR and 11 β -HSD (fig. 1). All of the above have been shown to be altered under conditions of immune activation. Proinflammatory cytokines tend

Fig. 1. Local factors regulating glucocorticoid bioavailability and action: (1) CBG, (2) MDR pump, (3) 11 β -HSD, (4) GR (= GR α) nuclear translocation, (5) GR interaction with other transcription factors [AP-1 (jun/fos), NF- κ B] or MAPK, and (6) ratio of GR α :GR β isoform expression. HSP = Heat shock protein. [Reprinted with permission from Silverman et al.]

to lower CBG levels, decrease MDR expression and/or function, and increase 11 β -HSD-1 expression and reductase activity (and decrease 11 β -HSD-2 expression/activity), thereby favoring an increase in glucocorticoid bioavailability. However, the opposite trend in each of these factors, which would favor a decrease in glucocorticoid bioavailability, may lead to an increased susceptibility to the development of autoimmune/inflammatory diseases.

Corticosteroid Binding Globulin

Only free or unbound glucocorticoids are capable of diffusing across the plasma membrane and activating the GR. Over 90% of circulating glucocorticoids are bound to CBG [11]. Therefore, the relative concentration of CBG is an important determinant of 'free' and available glucocorticoids. For example, decreases in the circulating level of CBG have been associated with evidence of occupation/activation of GR in the spleen of stressed rats [12]. Likewise, several studies have shown that in response to endotoxin administration in rats [13] and in a murine model of systemic lupus erythematosus [14], reduced CBG levels are observed. In addition, a decrease in plasma CBG-binding capacity has been reported in human septic shock and trauma [15–17]. In numerous studies, IL-6 has been shown to be a negative regulator of CBG production, and hence a determinant of cortisol bioavailability [16, 18]. In contrast, two studies examining plasma CBG during viral infection detected no

change in CBG levels in infected subjects compared to controls [19, 20]. It is important to note that even if a change in plasma CBG levels is not detected, local changes in the concentration and/or affinity of CBG during inflammation (for instance, within the microenvironment of the various lymphoid compartments or inflamed tissue) remain a consideration. In some cases, discrepancies exist between the concentration of free glucocorticoid levels in the plasma versus locally inflamed tissue, such as in RA patients who exhibit a greater percentage of bound corticosteroids in the synovial fluid relative to that found in plasma [21]. This observation demonstrates that the synovial membrane plays an important role in the local bioavailability of glucocorticoids in patients with RA.

Multidrug Resistance P-Glycoprotein

Multidrug resistance P-glycoprotein is an ATP-dependent multidrug efflux pump that decreases intracellular concentrations of potentially toxic chemicals (drugs and hormones). It is expressed in both human and rodent tissues, including the adrenal gland, kidney, liver, colon, small intestine, and brain and testis capillary endothelial cells [22]. The MDR pump at the blood-brain barrier, in both mice and humans, transports cortisol and the synthetic glucocorticoid dexamethasone, but not corticosterone, out of endothelial cells lining the brain [23]. In vitro studies have indicated that cytokines, such as IL-1, IL-6 and TNF- α , decrease MDR expression and/or function

in rodent hepatocytes [24, 25], human colon carcinoma cells [26] and human brain endothelial cells [27]. Moreover, rodents treated with LPS or turpentine exhibit reduced hepatic [24, 28–30], intestinal [31] and brain [28] MDR expression/activity. In contrast, humans with autoimmune diseases, such as RA [32, 33], colitis/Crohn's disease [34, 35] and lupus [36, 37], tend to exhibit high lymphocytic MDR expression and/or activity, which positively correlate with disease activity in some cases. However, this increased MDR expression may be secondary to treatment with high-dose glucocorticoids [35, 38]. Increased MDR expression has been found to be more prominent in a subpopulation of colitis [39] and RA [33, 40] patients who are resistant to steroid (or disease-modifying antirheumatic drugs) therapy and may be an underlying cause of their refractory response. Therefore, evaluation of MDR expression/activity may allow prediction of the efficacy of specific drug treatments. In addition, the use of MDR inhibitors (e.g., verapamil, cyclosporin A) may help to overcome treatment resistance or to improve incomplete responses in some RA patients, as has been shown in the case of chemotherapeutic agents [41]. Combined therapy with glucocorticoids and verapamil in a small group of lupus patients was shown to reduce MDR expression [36]; however, the clinical usefulness of this approach remains to be established.

Greater MDR function in immune cells may reduce glucocorticoid availability, thereby enhancing the synthesis/release of proinflammatory cytokines and exacerbating inflammatory responses. One possible reason for the discrepancy between the *in vitro* cytokine/animal studies and studies in humans with autoimmune disorders may be the differential effects of acute versus chronic inflammation on MDR expression. During acute inflammation, cytokines may downregulate MDR expression and increase local glucocorticoid concentrations, thereby limiting local inflammation and further cytokine release. In contrast, an upregulation of MDR expression may develop as a compensatory mechanism during chronic inflammatory conditions, and hence predispose humans toward autoimmune disease. In support of this contention, higher levels of peripheral blood mononuclear cell (PBMC) TNF- α mRNA were reported in RA patients exhibiting greater lymphocytic MDR activity, where reduced amounts of cortisol would be able to act intracellularly to inhibit proinflammatory responses [33]. Both high MDR activity and TNF- α levels were associated with poor outcome in RA. Another possible consideration regarding discrepant results is that changes in MDR expression/function may be cell type specific (e.g.,

PBMCs vs. synovial cells). Indeed, synovial cells express MDR; however, it may be an atypical MDR phenotype [42].

11 β -Hydroxysteroid Dehydrogenase

Another factor regulating the access of glucocorticoids to their receptors in target cells and tissues is 11 β -HSD. There are two isoforms of this enzyme, type 1 and type 2. Whereas 11 β -HSD-1 is a reversible oxidoreductase, 11 β -HSD-2 only exhibits oxidative or dehydrogenase activity [43]. 11 β -HSD-2 breaks down naturally occurring glucocorticoids (but not synthetic glucocorticoids) as they enter the cell, leaving the hormones in the form of inactive metabolites (i.e., cortisol to cortisone in humans; corticosterone to 11-dehydrocorticosterone in rodents). Significant differences in 11 β -HSD-2 activity have been found among immune compartments, and there is a direct correlation between 11 β -HSD-2 activity and the preferential production of Th1 versus Th2 cytokines by T cells residing in particular lymphoid organs [44]. Inhibition of 11 β -HSD-2 activity (which would enhance the available amount of hormone to bind to GR) leads to reduced Th1 responses and enhanced Th2 cytokine production by activated T cells [44].

Recent studies suggest that the bidirectional 11 β -HSD-1 prefers the reductase direction unless cells are disrupted [43]. Therefore, in intact tissues, 11 β -HSD-1 reactivates the inactive 11-keto glucocorticoids (corticosterone/11-dehydro-corticosterone) into their active 11-hydroxy glucocorticoid form (cortisol/corticosterone). Proinflammatory cytokines, such as TNF- α and IL-1 β , have been shown to upregulate 11 β -HSD-1 and/or downregulate 11 β -HSD-2 expression/activity in numerous cell types, including rat glomerular cells [45], human adipose stromal cells [46], and more related to RA, human bone cells, such as osteoblasts [47], human fibroblasts [48] and human and mouse vascular smooth muscle cells [49]. Inflamed colon specimens (in rats and humans) also exhibit elevated 11 β -HSD-1 and reduced 11 β -HSD-2 expression [50], thereby favoring the formation of active glucocorticoids and counterbalancing the proinflammatory effect of cytokines.

In immune cells, 11 β -HSD-1 is induced during the maturation of antigen-presenting cells, such as human macrophages [51] and murine dendritic cells [52]. It also has been shown to play a role in promoting macrophage phagocytosis of apoptotic leukocytes [53]. Moreover, 11 β -HSD-1 mRNA, protein, and (reductase) activity are expressed in murine lymphocytes, where activation of CD4⁺ T cells into Th1 or Th2 cells increases 11 β -HSD-1

activity [52]. In this case, the presence of greater levels of active glucocorticoids would reduce proinflammatory cytokine synthesis in Th1 cells and increase anti-inflammatory cytokine synthesis in Th2 cells. Therefore, changes in the relative activity of 11 β -HSD in immune tissues during inflammation may influence the relative impact of glucocorticoids on immune responses. Thus, the microenvironment of the various immune compartments is a potentially important site for intracrine regulation of neuroendocrine-immune interactions.

Reduced capacity for local reactivation of cortisone to cortisol has been observed in RA synovial cells, as evidenced by a greater ratio of 11 β -HSD-2:11 β -HSD-1-positive macrophages in the synovial tissue (compared to osteoarthritis synovium) [54]. This may be due to the local loss of catecholaminergic activity (loss of sympathetic nerve fibers and β -adrenergic receptor density on RA leukocytes) that would usually inhibit the inactivation of glucocorticoids. In addition, 11 β -HSD-2 was shown to be the second most overexpressed gene in RA and 11 β -HSD-2 protein levels correlated with inflammation scores [55]. Moreover, cortisone-induced inhibition of IL-6 in synovial fibroblasts has been shown to be dependent on 11 β -HSD-1 activity [48] and synovial macrophages from RA patients exhibit a reduced ability of the anti-inflammatory cytokine, IL-10, to induce 11 β -HSD-1 mRNA expression [56]. This defective reactivation of cortisone may be an important factor in the perpetuation of inflammation in patients with RA. The therapeutic value of 11 β -HSD-2 inhibitors has not been evaluated in RA, but may serve as another target of interest.

The Glucocorticoid Receptor

The ultimate effect of glucocorticoids on immune system regulation occurs at the level of the GR. Upon glucocorticoid binding to cytosolic GRs, a conformational change in GR causes heat-shock protein 90 (hsp90) and other ancillary proteins to dissociate from the receptor, and the ligand-bound receptor then translocates into the nucleus. Here, the glucocorticoid/GR complex acts as a transcription factor that regulates gene transcription through binding to glucocorticoid response elements (GREs) in the promoter regions of genes (requires GR dimerization) or through protein-protein interactions with other transcription factors (e.g., NF- κ B and AP-1; does not require GR dimerization). A tissue's sensitivity to glucocorticoid activity can be influenced by a change in (1) GR number or affinity or (2) GR function, including its ability for nuclear translocation, its interaction with other signal transduction pathways and the expression of

particular GR isoforms (fig. 1). Proinflammatory cytokines have been shown to impact a number of these factors.

GR Number and Affinity

There is a large body of data on the impact of cytokines on GR number [57]. However, the results are split into those studies that report an increase in GR number following cytokine administration and those that report a decrease. The discrepancy in results appears to depend on how the receptors were measured. Studies using whole-cell assay binding techniques tend to find an increase in cytokine-induced GR expression, while those using cytosolic receptor binding techniques tend to find a decrease. Few studies have documented changes in receptor affinity. Sher et al. [58] report that lymphocytes (T cells) simultaneously exposed to Th1 (IL-2) and Th2 (IL-4) cytokines (simulating the conditions of steroid-resistant asthma) exhibit a reduced affinity of GR for glucocorticoids. GRs on synovial fluid cells and PBMCs (but not synovial tissue cells) from RA patients have also been shown to exhibit reduced binding affinity [59]. However, no differences in GR binding affinity in PBMCs from RA patients compared to healthy controls have also been reported [60].

The GR number expressed in PBMCs of RA patients may be one factor in assessing glucocorticoid sensitivity and predicting which patients will respond to lower doses of steroid (e.g., prednisone) treatment and, therefore, avoid the unwanted side effects of higher doses. However, studies investigating GR expression in RA patients have given rise to contradictory results. Early diagnosed, untreated female (but not male) RA patients [61] and those with active disease of longer duration (off glucocorticoid therapy for at least 6 months) [60] exhibited reduced GR density in PBMCs compared to healthy controls, but GR density was not correlated with disease activity, suggesting that differential GR expression may not be involved in the pathogenesis of RA. Other studies have shown diminished GR numbers in PBMCs of glucocorticoid-treated patients, while those of untreated RA patients exhibited upregulated GR expression [62, 63], indicating that whether or not RA patients are undergoing drug therapy or are treatment naive can influence the level of GR expression. Interestingly, when measuring mRNA rather than protein, GR α mRNA expression in PBMCs was negatively correlated with disease activity [64]. Discrepant results may be due to the different sensitivities in change of expression levels between mRNA and protein.

Reduced PBMC GR density has not been associated with functional glucocorticoid resistance in the sense that proliferation and cytokine release of lymphocytes of RA patients and healthy controls were inhibited by glucocorticoids to the same extent [65]. On the other hand, when subpopulations of glucocorticoid-resistant versus glucocorticoid-sensitive RA patients are selected, differences in PBMC proliferation can be detected between the two groups. Inhibition of PBMC proliferation after steroid treatment was significantly lower in a glucocorticoid-resistant as compared to a glucocorticoid-sensitive group [66]. Moreover, a positive correlation has been shown between glucocorticoid-induced inhibition of PBMC proliferation and the clinical outcome of glucocorticoid treatment in RA [67]. Therefore, the measurement of steroid sensitivity of peripheral lymphocytes (not necessarily reflected by GR number) may be a useful tool in predicting the therapeutic efficacy of glucocorticoids in RA patients.

Although all nucleated cells in humans have GRs, in RA, it is important to consider possible abnormalities in GR expression/function in the synovium, and hence glucocorticoid sensitivity at the site of inflammation. Indeed, GRs are expressed by multiple cell types in synovial tissue, including lymphocytes, fibroblasts, endothelial cells and smooth muscle cells in the sublining layer [68], suggesting that glucocorticoids directly target the synovium. GR expression on subsynovial fibroblastic cells of RA patients pretreated with glucocorticoids was significantly lower compared to patients who had not received GC treatment; however, less suppression of GR expression was observed in patients on low-dose GC treatment [68], supporting the majority of findings regarding GR expression in PBMCs of RA patients. Further investigation is needed to determine how synovial GR expression may relate to disease activity.

GR Function

Several autoimmune/inflammatory disorders have been associated with impaired GR function, possibly contributing to the excessive inflammation characteristic of these illnesses [69]. Proinflammatory cytokines (i.e., TNF- α , IL-1, and IL-6) and cytokines that mediate lymphocyte growth and differentiation (i.e., IL-2, IL-4) have been found to inhibit GR function [57].

Although the mechanism by which cytokines inhibit GR is unknown, several possibilities have been considered. First, cytokines may influence GR function through their effects on GR translocation from the cytoplasm into the nucleus. For example, TNF- α and IL-1 have been

shown to block dexamethasone-induced GR nuclear translocation [70, 71]. Another possible mechanism by which cytokines may influence GR is through cross talk among signal transduction pathways. For example, downstream in the IL-1 and TNF- α signal transduction pathways are the transcription factors NF- κ B and AP-1 (which consists of jun and fos proteins). GR and NF- κ B/AP-1 have been shown to mutually antagonize each other's transcriptional activity through multiple mechanisms [72–74]. In addition, IL-1 and TNF- α activate mitogen-activated protein kinase (MAPK) signaling pathways [i.e., p38 MAPK and jun N-terminal kinase (JNK)], which have been shown to be inhibitory to GR function [75–78]. It has recently been shown that a major mechanism by which glucocorticoids exert their inhibitory action on MAPK pathways, and therefore inflammation, is by the upregulation of the MAPK phosphatase, MKP-1 [79]. Other cytokines, such as IL-6, IFN- α/β , IFN- γ , IL-12, IL-2, and IL-4, induce janus kinase (JAK)/signal transduction and activator of transcription (STAT) signaling pathways, and STAT proteins have been found to reciprocally influence GR activity by direct protein-protein interactions [72, 74, 80]. Indeed, immune-related transcription factors, such as NF- κ B, AP-1, MAPKs and STATs, are elevated in PBMCs and/or synovial tissue of RA patients [81–85], and may serve as another mechanism by which cytokines can confer a glucocorticoid-resistant state. On the other hand, a deficit in inhibitory signaling molecules, such as MKP-1, may also contribute to reduced glucocorticoid sensitivity in inflammatory/autoimmune diseases [79]. Interestingly, elevated levels of macrophage inhibitory factor, a proinflammatory cytokine that directly counteracts the immunosuppressive actions of glucocorticoids, have been demonstrated in RA and other chronic inflammatory diseases, and has been shown to exert its antiglucocorticoid effects through MKP-1 inhibition [86, 87]. Therefore, therapeutic interventions aimed at antagonizing inflammatory signaling pathways may be another way to increase glucocorticoid sensitivity.

Finally, cytokines may affect GR function by altering the ratio of GR α :GR β isoform expression. Alternative splicing of the human GR primary transcript produces multiple isoforms [88]. The two that have received the most attention are GR α , the classical transcriptionally active isoform, and GR β , which may negatively regulate GR α activity. The GR α isoform binds hormone and activates glucocorticoid-responsive genes, while the GR β isoform fails to bind hormone and activate glucocorticoid-responsive genes and attenuates the *trans*-activation

of the hormone-bound GR α isoform [89]. These findings suggest that when GR β is present in excess in various tissues (reduced GR α :GR β ratio), it can act as a dominant negative inhibitor of GR α activity. In fact, Webster et al. [90] have shown that IL-1 and TNF- α lead to the selective accumulation of GR β protein in cells of lymphoid origin and the development of a glucocorticoid-resistant state. Interestingly, glucocorticoid sensitivity in different immune cell types has been associated with varied degrees of GR β expression [91, 92]. Moreover, increases in GR β levels have been found in immune cells of patients with glucocorticoid-resistant asthma [93, 94], colitis/Crohn's disease [95, 96] and RA [97].

Derijk et al. [98] were the first to demonstrate that a polymorphism in the human GR β gene, which increases its mRNA stability, is associated with RA. Other polymorphisms of the GR have also been associated with RA, whereby those that confer an increased glucocorticoid sensitivity (i.e., N363S and *BclI*) are associated with a decreased susceptibility to develop RA and those related to reduced glucocorticoid sensitivity (i.e., ER22/23EK and GR β -A3669G) are associated with an enhanced predisposition to develop RA [van Oosten et al., unpubl. data]. However, other studies have failed to find an association with GR polymorphisms and RA susceptibility [99, 100]. Screening for GR polymorphisms in RA patients may also aid in the identification of those who will benefit the most from glucocorticoid treatment [for a review of common GR polymorphisms and their associations with disease, see 101, 102]. Alternative translation initiation of GR α mRNA can also lead to the expression of GR isoforms with different transcriptional activity [103]. The unique transcriptional activities and distinct tissue-specific distribution patterns of GR α isoforms could provide a novel mechanism for tissue-specific glucocorticoid responses.

References

- 1 Besedovsky HO, del Rey A: Immune-neuro-endocrine interactions: facts and hypotheses. *Endocr Rev* 1996;17:64–102.
- 2 Silverman MN, Pearce BD, Biron CA, Miller AH: Immune modulation of the hypothalamic-pituitary-adrenal (HPA) axis during viral infection. *Viral Immunol* 2005;18:41–78.

Table 1. Molecular mechanisms of glucocorticoid resistance

↑ CBG	→	↓ free GCs
↑ MDR Pgp expression	→	↓ intracellular GCs
↓ 11 β -HSD-1/↑ 11 β -HSD-2	→	↓ active GCs
↓ GR affinity for GCs	→	↓ GR function
↓ GR translocation into nucleus	→	↓ GR function
↑ Expression of AP-1 and NF- κ B	→	↓ GR function
↑ Expression of MAPKs (p38, JNK)	→	↓ GR function
↑ Expression of GR β isoform	→	↓ GR function

MDR Pgp = Multidrug resistance P-glycoprotein.

Conclusion

A variety of factors have been shown to regulate GC availability or bioactivity. A change in any of them towards reduced glucocorticoid function can lead to the development of glucocorticoid resistance (table 1). Should this occur during chronic inflammation, patients would be expected to be more susceptible to the deleterious, tissue-damaging effects of an overproduction of proinflammatory cytokines, which may increase vulnerability to the development of autoimmune disease. Further investigation is required to assess the safety and efficacy of new therapeutic strategies, including inhibitors of the multi-drug resistance P-glycoprotein, 11 β -HSD-2 and downstream inflammatory signaling molecules (e.g., AP-1, NF- κ B, STAT, MAPKs), which may help to reverse a state of glucocorticoid resistance.

Acknowledgments

The authors would like to thank the National Institute of Mental Health for their continued support and the Prince of Wales Foundation funded NCCAM Director's Fellowship that has been granted to MNS.

- 5 Straub RH, Cutolo M: Involvement of the hypothalamic-pituitary-adrenal/gonadal axis and the peripheral nervous system in rheumatoid arthritis: viewpoint based on a systemic pathogenetic role. *Arthritis Rheum* 2001;44:493–507.
- 6 Straub RH, Dhabhar FS, Bijlsma JW, Cutolo M: How psychological stress via hormones and nerve fibers may exacerbate rheumatoid arthritis. *Arthritis Rheum* 2005;52:16–26.
- 7 Crofford LJ, Kalogeras KT, Mastorakos G, Magiakou MA, Wells J, Kanik KS, Gold PW, Chrousos GP, Wilder RL: Circadian relationships between interleukin (IL)-6 and hypothalamic-pituitary-adrenal axis hormones: failure of IL-6 to cause sustained hypercortisolism in patients with early untreated rheumatoid arthritis. *J Clin Endocrinol Metab* 1997;82:1279–1283.
- 8 Straub RH, Cutolo M: Circadian rhythms in rheumatoid arthritis: implications for pathophysiology and therapeutic management. *Arthritis Rheum* 2007;56:399–408.
- 9 Sternberg EM, Hill JM, Chrousos GP, Kamilaris T, Listwak SJ, Gold PW, Wilder RL: Inflammatory mediator-induced hypothalamic-pituitary-adrenal axis activation is defective in streptococcal cell wall arthritis-susceptible Lewis rats. *Proc Natl Acad Sci USA* 1989;86:2374–2378.
- 10 Eskandari F, Webster JI, Sternberg EM: Neural immune pathways and their connection to inflammatory diseases. *Arthritis Res Ther* 2003;5:251–265.
- 11 Rosner W: Plasma steroid-binding proteins. *Endocrinol Metab Clin* 1991;20:697–720.
- 12 Spencer RL, Miller AH, Moday H, McEwen BS, Blanchard RJ, Blanchard DC, Sakai RR: Chronic social stress produces reductions in available splenic type II corticosteroid receptor binding and plasma corticosteroid binding globulin levels. *Psychoneuroendocrinology* 1996;21:95–109.
- 13 Shanks N, Windle RJ, Perks P, Wood S, Ingram CD, Lightman SL: The hypothalamic-pituitary-adrenal axis response to endotoxin is attenuated during lactation. *J Neuroendocrinol* 1999;11:857–865.
- 14 Shanks N, Moore PM, Perks P, Lightman SL: Alterations in hypothalamic-pituitary-adrenal function correlated with the onset of murine SLE in MRL +/+ and lpr/lpr mice. *Brain Behav Immun* 1999;13:348–360.
- 15 Beishuizen A, Thijs LG, Vermes I: Patterns of corticosteroid-binding globulin and the free cortisol index during septic shock and multiple trauma. *Intensive Care Med* 2001;27:1584–1591.
- 16 Bernier J, Jobin N, Emptoz-Bonneton A, Pugeat MM, Garrel DR: Decreased corticosteroid-binding globulin in burn patients: relationship with interleukin-6 and fat in nutritional support. *Crit Care Med* 1998;26:452–460.
- 17 Ho JT, Al-Musallhi H, Chapman MJ, Quach T, Thomas PD, Bagley CJ, Lewis JG, Torpy DJ: Septic shock and sepsis: a comparison of total and free plasma cortisol levels. *J Clin Endocrinol Metab* 2006;91:105–114.
- 18 Tsigos C, Kyrou I, Chrousos GP, Papanicolaou DA: Prolonged suppression of corticosteroid-binding globulin by recombinant human interleukin-6 in man. *J Clin Endocrinol Metab* 1998;83:3379.
- 19 Lambert M: Thyroid dysfunction in HIV infection. *Baillieres Clin Endocrinol Metab* 1994;8:825–835.
- 20 Miller AH, Spencer RL, Pearce BD, Pisell TL, Tanapat P, Leung JJ, Dhabhar FS, McEwen BS, Biron CA: 1996 Curt P. Richter Award. Effects of viral infection on corticosterone secretion and glucocorticoid receptor binding in immune tissues. *Psychoneuroendocrinology* 1997;22:455–474.
- 21 Feher KG, Zahumenszky Z, Feher T, Banyai B: Protein-corticosteroid interactions in peripheral plasma and synovial fluid of patients with rheumatoid arthritis and osteoarthritis. *Z Rheumatol* 1980;39:176–181.
- 22 Chin KV, Chauhan SS, Pastan I, Gottesman MM: Regulation of mdr RNA levels in response to cytotoxic drugs in rodent cells. *Cell Growth Differ* 1990;1:361–365.
- 23 Karssen AM, Meijer OC, van der Sandt IC, Lucassen PJ, de Lange EC, de Boer AG, de Kloet ER: Multidrug resistance P-glycoprotein hampers the access of cortisol but not of corticosterone to mouse and human brain. *Endocrinology* 2001;142:2686–2694.
- 24 Hartmann G, Kim H, Piquette-Miller M: Regulation of the hepatic multidrug resistance gene expression by endotoxin and inflammatory cytokines in mice. *Int Immunopharmacol* 2001;1:189–199.
- 25 Sukhai M, Yong A, Pak A, Piquette-Miller M: Decreased expression of P-glycoprotein in interleukin-1 β and interleukin-6 treated rat hepatocytes. *Inflamm Res* 2001;50:362–370.
- 26 Stein U, Walther W, Shoemaker RH: Modulation of mdr1 expression by cytokines in human colon carcinoma cells: an approach for reversal of multidrug resistance. *Br J Cancer* 1996;74:1384–1391.
- 27 Mandi Y, Ocsosvski I, Szabo D, Nagy Z, Nelson J, Molnar J: Nitric oxide production and MDR expression by human brain endothelial cells. *Anticancer Res* 1998;18:3049–3052.
- 28 Goralski KB, Hartmann G, Piquette-Miller M, Renton KW: Downregulation of mdr1a expression in the brain and liver during CNS inflammation alters the in vivo disposition of digoxin. *Br J Pharmacol* 2003;139:35–48.
- 29 Piquette-Miller M, Pak A, Kim H, Anari R, Shahzamani A: Decreased expression and activity of P-glycoprotein in rat liver during acute inflammation. *Pharm Res* 1998;15:706–711.
- 30 Sukhai M, Yong A, Kalitsky J, Piquette-Miller M: Inflammation and interleukin-6 mediate reductions in the hepatic expression and transcription of the mdr1a and mdr1b genes. *Mol Cell Biol Res Commun* 2000;4:248–256.
- 31 Kalitsky-Szirtes J, Shayeganpour A, Brocks DR, Piquette-Miller M: Suppression of drug-metabolizing enzymes and efflux transporters in the intestine of endotoxin-treated rats. *Drug Metab Dispos* 2004;32:20–27.
- 32 Borowski LC, Lopes RP, Gonzalez TP, Dummer LA, Chies JA, Silveira IG, Keisermann M, Bauer ME: Is steroid resistance related to multidrug resistance-I (MDR-I) in rheumatoid arthritis? *Int Immunopharmacol* 2007;7:836–844.
- 33 Llorente L, Richaud-Patin Y, Diaz-Borjon A, Alvarado de la Barrera C, Jakez-Ocampo J, de la Fuente H, Gonzalez-Amaro R, Diaz-Jouanen E: Multidrug resistance-I (MDR-I) in rheumatic autoimmune disorders. I. Increased P-glycoprotein activity in lymphocytes from rheumatoid arthritis patients might influence disease outcome. *Joint Bone Spine* 2000;67:30–39.
- 34 Farrell RJ, Murphy A, Long A, Donnelly S, Cherikuri A, O'Toole D, Mahmud N, Keeling PW, Weir DG, Kelleher D: High multidrug resistance (P-glycoprotein 170) expression in inflammatory bowel disease patients who fail medical therapy. *Gastroenterology* 2000;118:279–288.
- 35 Hirano T, Onda K, Toma T, Miyaoka M, Moriyasu F, Oka K: MDR1 mRNA expressions in peripheral blood mononuclear cells of patients with ulcerative colitis in relation to glucocorticoid administration. *J Clin Pharmacol* 2004;44:481–486.
- 36 Diaz-Borjon A, Richaud-Patin Y, Alvarado de la Barrera C, Jakez-Ocampo J, Ruiz-Arguelles A, Llorente L: Multidrug resistance-I (MDR-I) in rheumatic autoimmune disorders. II. Increased P-glycoprotein activity in lymphocytes from systemic lupus erythematosus patients might affect steroid requirements for disease control. *Joint Bone Spine* 2000;67:40–48.
- 37 Tsujimura S, Saito K, Nakayamada S, Tanaka Y: Relevance of multidrug resistance 1 and P-glycoprotein to drug resistance in patients with systemic lupus erythematosus. *Histol Histopathol* 2007;22:465–468.
- 38 Maillefert JF, Maynadie M, Tebib JG, Aho S, Walker P, Chatard C, Dulieu V, Bouvier M, Carli PM, Tavernier C: Expression of the multidrug resistance glycoprotein 170 in the peripheral blood lymphocytes of rheumatoid arthritis patients. The percentage of lymphocytes expressing glycoprotein 170 is increased in patients treated with prednisolone. *Br J Rheumatol* 1996;35:430–435.

- 39 Sambuelli AM, Negreira SM, Gil AH, Huer-
nos SP, Goncalves S, Toro MA, Kogan Z, Ca-
banne A, Camarero S, Bai JC, Lazarowski AJ:
Multidrug resistance gene (MDR-1) expres-
sion in the colonic mucosa of patients with
refractory ulcerative colitis. *Acta Gastroen-
terol Latinoam* 2006;36:23–32.
- 40 Yudoh K, Matsuno H, Nakazawa F, Yoneza-
wa T, Kimura T: Increased expression of
multidrug resistance of P-glycoprotein on
Th1 cells correlates with drug resistance in
rheumatoid arthritis. *Arthritis Rheum* 1999;
42:2014–2015.
- 41 Richaud-Patin Y, Soto-Vega E, Jakez-Ocam-
po J, Llorente L: P-glycoprotein in autoim-
mune diseases. *Autoimmun Rev* 2004;3:
188–192.
- 42 Jorgensen C, Sun R, Rossi JF, Costes J, Rich-
ard D, Bologna C, Sany J: Expression of a
multidrug resistance gene in human rheu-
matoid synovium. *Rheumatol Int* 1995;15:
83–86.
- 43 Seckl JR, Walker BR: Minireview: 11beta-hy-
droxysteroid dehydrogenase type 1 – a tis-
sue-specific amplifier of glucocorticoid ac-
tion. *Endocrinology* 2001;142:1371–1376.
- 44 Hennebold JD, Ryu SY, Mu HH, Galbraith A,
Daynes RA: 11 beta-hydroxysteroid dehy-
drogenase modulation of glucocorticoid ac-
tivities in lymphoid organs. *Am J Physiol*
1996;270:R1296–R1306.
- 45 Escher G, Galli I, Vishwanath BS, Frey BM,
Frey FJ: Tumor necrosis factor alpha and in-
terleukin 1beta enhance the cortisone/corti-
sol shuttle. *J Exp Med* 1997;186:189–198.
- 46 Tomlinson JW, Moore J, Cooper MS, Bujal-
ska I, Shahmanesh M, Burt C, Strain A,
Hewison M, Stewart PM: Regulation of ex-
pression of 11beta-hydroxysteroid dehydro-
genase type 1 in adipose tissue: tissue-spe-
cific induction by cytokines. *Endocrinology*
2001;142:1982–1989.
- 47 Cooper MS, Bujalska I, Rabbitt E, Walker
EA, Bland R, Sheppard MC, Hewison M,
Stewart PM: Modulation of 11beta-hy-
droxysteroid dehydrogenase isozymes by
proinflammatory cytokines in osteoblasts:
an autocrine switch from glucocorticoid in-
activation to activation. *J Bone Miner Res*
2001;16:1037–1044.
- 48 Hardy RS, Filer A, Cooper MS, Parsonage G,
Raza K, Hardie DL, Rabbitt EH, Stewart PM,
Buckley CD, Hewison M: Differential ex-
pression, function and response to inflam-
matory stimuli of 11beta-hydroxysteroid de-
hydrogenase type 1 in human fibroblasts: a
mechanism for tissue-specific regulation of
inflammation. *Arthritis Res Ther* 2006;8:
R108.
- 49 Dover AR, Hadoke PW, Macdonald LJ, Mil-
ler E, Newby DE, Walker BR: Intravascular
glucocorticoid metabolism during inflam-
mation and injury in mice. *Endocrinology*
2007;148:166–172.
- 50 Zbankova S, Bryndova J, Leden P, Kment M,
Svec A, Pacha J: 11beta-hydroxysteroid dehy-
drogenase 1 and 2 expression in colon from
patients with ulcerative colitis. *J Gastroen-
terol Hepatol* 2007;22:1019–1023.
- 51 Thieringer R, Le Grand CB, Carbin L, Cai
TQ, Wong B, Wright SD, Hermanowski-Vo-
satka A: 11 Beta-hydroxysteroid dehydroge-
nase type 1 is induced in human monocytes
upon differentiation to macrophages. *J Im-
munol* 2001;167:30–35.
- 52 Zhang YT, Ding X, Daynes RA: The expres-
sion of 11 beta-hydroxysteroid dehydroge-
nase type I by lymphocytes provides a novel
means for intracrine regulation of glucocor-
ticoid activities. *J Immunol* 2005;174:879–
889.
- 53 Gilmour JS, Coutinho AE, Cailhier JF, Man
TY, Clay M, Thomas G, Harris HJ, Mullins
JJ, Seckl JR, Savill JS, Chapman KE: Local
amplification of glucocorticoids by 11 beta-
hydroxysteroid dehydrogenase type 1 pro-
motes macrophage phagocytosis of apoptot-
ic leukocytes. *J Immunol* 2006;176:7605–
7611.
- 54 Schmidt M, Weidler C, Naumann H, Anders
S, Scholmerich J, Straub RH: Reduced capac-
ity for the reactivation of glucocorticoids in
rheumatoid arthritis synovial cells: possible
role of the sympathetic nervous system? *Ar-
thritis Rheum* 2005;52:1711–1720.
- 55 Haas CS, Creighton CJ, Pi X, Maine I, Koch
AE, Haines GK, Ling S, Chinnaiyan AM,
Holoshitz J: Identification of genes modulat-
ed in rheumatoid arthritis using comple-
mentary DNA microarray analysis of lym-
phoblastoid B cell lines from disease-
discordant monozygotic twins. *Arthritis
Rheum* 2006;54:2047–2060.
- 56 Antoniv TT, Ivashkiv LB: Dysregulation of
interleukin-10-dependent gene expression
in rheumatoid arthritis synovial macro-
phages. *Arthritis Rheum* 2006;54:2711–
2721.
- 57 Miller AH, Pearce BD, Ruzek MC, Biron CA:
Interactions between the hypothalamic-pi-
tuitary-adrenal axis and immune system
during viral infection: pathways for environ-
mental effects on disease expression; in
McEwen BS (ed): *Handbook of Physiology*.
Section 7: The Endocrine System. New York,
Oxford University Press, 2001, vol 4: Coping
with the Environment: Neural and Endo-
crine Mechanisms, pp 425–450.
- 58 Sher ER, Leung DY, Surs W, Kam JC, Zieg G,
Kamada AK, Szefer SJ: Steroid-resistant
asthma. Cellular mechanisms contributing
to inadequate response to glucocorticoid
therapy. *J Clin Invest* 1994;93:33–39.
- 59 Braidman IP, Livingston K, Tomlinson IW,
Jayson MI, Jasani MK: Effect of inflamma-
tory arthritis on dexamethasone receptors in
phagocytes – a possible role for plasma fac-
tors. *Agents Actions* 1981;11:602–604.
- 60 Schlaghecke R, Kornely E, Wollenhaupt J,
Specker C: Glucocorticoid receptors in rheu-
matoid arthritis. *Arthritis Rheum* 1992;35:
740–744.
- 61 van Everdingen AA, Huisman AM, Wenting
MJ, van Reesema S, Jacobs JW, Bijlsma JW:
Downregulation of glucocorticoid receptors
in early-diagnosed rheumatoid arthritis.
Clin Exp Rheumatol 2002;20:463–468.
- 62 Eggert M, Kluter A, Rusch D, Schmidt KL,
Dotzlaw H, Schulz M, Pabst W, Boke J, Ren-
kawitz R, Neeck G: Expression analysis of
the glucocorticoid receptor and the nuclear
factor-kB subunit p50 in lymphocytes from
patients with rheumatoid arthritis. *J Rheu-
matol* 2002;29:2500–2506.
- 63 Sanden S, Tripmacher R, Weltrich R, Rohde
W, Hiepe F, Burmester GR, Buttgerit F: Glu-
cocorticoid dose dependent downregulation
of glucocorticoid receptors in patients with
rheumatic diseases. *J Rheumatol* 2000;27:
1265–1270.
- 64 Onda K, Rimbara E, Hirano T, Oka K, Abe
H, Tahara K, Takanashi H, Tsuboi N, Niit-
suma T, Hayashi T: Role of mRNA expres-
sion of transcription factors in glucocorti-
coid sensitivity of peripheral blood mono-
nuclear cells and disease state in rheumatoid
arthritis. *J Rheumatol* 2004;31:464–469.
- 65 Schlaghecke R, Beuscher D, Kornely E,
Specker C: Effects of glucocorticoids in rheu-
matoid arthritis. Diminished glucocorticoid
receptors do not result in glucocorticoid re-
sistance. *Arthritis Rheum* 1994;37:1127–
1131.
- 66 Sliwinski-Stanczyk P, Pazdur J, Ziolkowska
M, Jaworski J, Kaminska-Tchorzewska E,
Lacki JK: The effect of methylprednisolone
on proliferation of PBMCs obtained from
steroid-sensitive and steroid-resistant rheu-
matoid arthritis patients. *Scand J Rheumatol*
2007;36:167–171.
- 67 Kirkham BW, Corkill MM, Davison SC,
Panayi GS: Response to glucocorticoid treat-
ment in rheumatoid arthritis: in vitro cell
mediated immune assay predicts in vivo re-
sponses. *J Rheumatol* 1991;18:821–825.
- 68 Tohyama CT, Yamakawa M, Murasawa A,
Nakazono K, Ishikawa H: Localization of
human glucocorticoid receptor in rheuma-
toid synovial tissue of the knee joint. *Scand J
Rheumatol* 2005;34:426–432.
- 69 DeRijk R, Sternberg EM: Corticosteroid re-
sistance and disease. *Ann Med* 1997;29:79–
82.
- 70 Adcock IM, Brown CR, Barnes PJ: Tumour
necrosis factor alpha causes retention of ac-
tivated glucocorticoid receptor within the
cytoplasm of A549 cells. *Biochem Biophys
Res Commun* 1996;225:545–550.
- 71 Pariente CM, Pearce BD, Pisell TL, Sanchez
CI, Po C, Su C, Miller AH: The proinflam-
matory cytokine, interleukin-1alpha, reduc-
es glucocorticoid receptor translocation
and function. *Endocrinology* 1999;140:
4359–4366.

- 72 Adcock IM, Caramori G: Cross-talk between pro-inflammatory transcription factors and glucocorticoids. *Immunol Cell Biol* 2001;79: 376–384.
- 73 De Bosscher K, Vanden Berghe W, Haegeman G: Mechanisms of anti-inflammatory action and of immunosuppression by glucocorticoids: negative interference of activated glucocorticoid receptor with transcription factors. *J Neuroimmunol* 2000;109:16–22.
- 74 Smoak KA, Cidlowski JA: Mechanisms of glucocorticoid receptor signaling during inflammation. *Mech Ageing Dev* 2004;125: 697–706.
- 75 Irusen E, Matthews JG, Takahashi A, Barnes PJ, Chung KF, Adcock IM: p38 Mitogen-activated protein kinase-induced glucocorticoid receptor phosphorylation reduces its activity: role in steroid-insensitive asthma. *J Allergy Clin Immunol* 2002;109:649–657.
- 76 Rogatsky I, Logan SK, Garabedian MJ: Antagonism of glucocorticoid receptor transcriptional activation by the c-Jun N-terminal kinase. *Proc Natl Acad Sci USA* 1998;95: 2050–2055.
- 77 Wang X, Wu H, Lakdawala VS, Hu F, Hanson ND, Miller AH: Inhibition of Jun N-terminal kinase (JNK) enhances glucocorticoid receptor-mediated function in mouse hippocampal HT22 cells. *Neuropsychopharmacology* 2005;30:242–249.
- 78 Wang X, Wu H, Miller AH: Interleukin 1 α (IL-1 α) induced activation of p38 mitogen-activated protein kinase inhibits glucocorticoid receptor function. *Mol Psychiatry* 2004;9:65–75.
- 79 Clark AR: MAP kinase phosphatase 1: a novel mediator of biological effects of glucocorticoids? *J Endocrinol* 2003;178:5–12.
- 80 Shuai K: Modulation of STAT signaling by STAT-interacting proteins. *Oncogene* 2000; 19:2638–2644.
- 81 Chikanza IC, Kozaci DL: Corticosteroid resistance in rheumatoid arthritis: molecular and cellular perspectives. *Rheumatology (Oxford)* 2004;43:1337–1345.
- 82 Eggert M, Kluter A, Zettl UK, Neeck G: Transcription factors in autoimmune diseases. *Curr Pharm Des* 2004;10:2787–2796.
- 83 Ivashkiv LB, Hu X: The JAK/STAT pathway in rheumatoid arthritis: pathogenic or protective? *Arthritis Rheum* 2003;48:2092–2096.
- 84 Korb A, Tohidast-Akrad M, Cetin E, Axmann R, Smolen J, Schett G: Differential tissue expression and activation of p38 MAPK α , β , γ , and δ isoforms in rheumatoid arthritis. *Arthritis Rheum* 2006; 54:2745–2756.
- 85 Rottapel R: Putting the brakes on arthritis: can suppressors of cytokine signaling (SOCS) suppress rheumatoid arthritis? *J Clin Invest* 2001;108:1745–1747.
- 86 Aeberli D, Leech M, Morand EF: Macrophage migration inhibitory factor and glucocorticoid sensitivity. *Rheumatology (Oxford)* 2006;45:937–943.
- 87 Flaster H, Bernhagen J, Calandra T, Bucala R: The macrophage migration inhibitory factor-glucocorticoid dyad: regulation of inflammation and immunity. *Mol Endocrinol* 2007;21:1267–1280.
- 88 Zhou J, Cidlowski JA: The human glucocorticoid receptor: one gene, multiple proteins and diverse responses. *Steroids* 2005;70: 407–417.
- 89 Lewis-Tuffin LJ, Cidlowski JA: The physiology of human glucocorticoid receptor beta (hGR β) and glucocorticoid resistance. *Ann NY Acad Sci* 2006;1069:1–9.
- 90 Webster JC, Oakley RH, Jewell CM, Cidlowski JA: Proinflammatory cytokines regulate human glucocorticoid receptor gene expression and lead to the accumulation of the dominant negative beta isoform: a mechanism for the generation of glucocorticoid resistance. *Proc Natl Acad Sci USA* 2001;98: 6865–6870.
- 91 Li LB, Leung DY, Hall CF, Goleva E: Divergent expression and function of glucocorticoid receptor beta in human monocytes and T cells. *J Leukoc Biol* 2006;79:818–827.
- 92 Strickland I, Kisich K, Hauk PJ, Vottero A, Chrousos GP, Klemm DJ, Leung DY: High constitutive glucocorticoid receptor beta in human neutrophils enables them to reduce their spontaneous rate of cell death in response to corticosteroids. *J Exp Med* 2001; 193:585–593.
- 93 Goleva E, Li LB, Eves PT, Strand MJ, Martin RJ, Leung DY: Increased glucocorticoid receptor beta alters steroid response in glucocorticoid-insensitive asthma. *Am J Respir Crit Care Med* 2006;173:607–616.
- 94 Sousa AR, Lane SJ, Cidlowski JA, Staynov DZ, Lee TH: Glucocorticoid resistance in asthma is associated with elevated *in vivo* expression of the glucocorticoid receptor beta-isoform. *J Allergy Clin Immunol* 2000;105:943–950.
- 95 Honda M, Orii F, Ayabe T, Imai S, Ashida T, Obara T, Kohgo Y: Expression of glucocorticoid receptor beta in lymphocytes of patients with glucocorticoid-resistant ulcerative colitis. *Gastroenterology* 2000; 118:859–866.
- 96 Towers R, Naftali T, Gabay G, Carlebach M, Klein A, Novis B: High levels of glucocorticoid receptors in patients with active Crohn's disease may predict steroid resistance. *Clin Exp Immunol* 2005;141:357–362.
- 97 Kozaci DL, Chernajovsky Y, Chikanza IC: The differential expression of corticosteroid receptor isoforms in corticosteroid-resistant and -sensitive patients with rheumatoid arthritis. *Rheumatology (Oxford)* 2007;46:579–585.
- 98 Derijk RH, Schaaf MJ, Turner G, Datson NA, Vreugdenhil E, Cidlowski J, de Kloet ER, Emery P, Sternberg EM, Detera-Wadleigh SD: A human glucocorticoid receptor gene variant that increases the stability of the glucocorticoid receptor beta-isoform mRNA is associated with rheumatoid arthritis. *J Rheumatol* 2001;28:2383–2388.
- 99 Donn R, Payne D, Ray D: Glucocorticoid receptor gene polymorphisms and susceptibility to rheumatoid arthritis. *Clin Endocrinol (Oxf)* 2007;67:342–345.
- 100 Lee EB, Kim JY, Lee YJ, Song YW: Glucocorticoid receptor polymorphisms in Korean patients with rheumatoid arthritis. *Ann Rheum Dis* 2005;64:503–504.
- 101 DeRijk RH, Schaaf M, de Kloet ER: Glucocorticoid receptor variants: clinical implications. *J Steroid Biochem Mol Biol* 2002; 81:103–122.
- 102 van Rossum EF, Russcher H, Lamberts SW: Genetic polymorphisms and multifactorial diseases: facts and fallacies revealed by the glucocorticoid receptor gene. *Trends Endocrinol Metab* 2005;16:445–450.
- 103 Lu NZ, Cidlowski JA: Glucocorticoid receptor isoforms generate transcription specificity. *Trends Cell Biol* 2006;16:301–307.