Reply

T.-M.

Wang

Department of Biology and Anatomy, National Defense Medical Center, Taipei, Taiwan, ROC

Teen-Meei Wang, PhD, Department of Biology and Anatomy, National Defense Medical Center, PO Box 90048-502, Taipei, Taiwan 100 (ROC)

Dear Sir

In response to Dr. Sreenathan's letter, first of all, I would like to make an apology for the mistake we made in figure 1a (i.e. misplaced S,) which has been corrected and shown in the enclosed photo. The cephalad midsagittal diameters of the lumbar spinal canals in Chinese and Indian adult skeletons shown in table 1 (varying from 4.5 ± 0.18 to 8.07 ± 0.19 mm) are correct and measured based on the distance between the convex part of the posterior surface of the vertebral body and the nearest part of the cephalad border of the laminar arch. In addition, the data concerning S, (in fig. 1a) in *Acta Anatomica* (144: 25) have been measured before (unpublished) and shown to range from 12.56 to 18.24 mm.

Fig. 1 a.

Sj=Cephalad mid-sagittal diameter between the convex part of the posterior surface of the vertebral body and the nearest part of the cephalad border of the laminar arch.

© 1994 S.KargerAG, Basel 0001-5180/94/1491-0080 \$ 5.00/0