Preface

D.F.	Donald F.	Newgreen ^a
M.W.	Mark W.J.	Ferguson ^b

^aMelbourne and ^bManchester

Epithelium and mesenchyme constitute the two basic cell phenotypes, and these two states starkly contrast both in the extracellular relationships of the cell to adjacent cells and noncellular structures, and in the intracellular organisation of the cytoskeleton and metabolic apparatus. Yet, despite these fundamental differences, these two contrasting cell states are interconvertible. By definition, the cells of metazoans must cling together: Evolutionarily, early forms may have consisted of epithelial-like structures with specialised, but still epithelial, infoldings. Although this meant that considerable control of the internal milieu was possible, to achieve the vast complexity of form and great range of sizes seen in the animal kingdom required cellular constituents between the epithelial cell layers. An expedient, early adopted in evolution, and used repeatedly in development, is simply the controlled conversion of certain epithelial cells into cells capable of occupying the subepithelial spaces, into cells capable of giving this space new structure and function, that is, into mesenchyme cells.

In the present volume, an overview of the field of epi-thelial-mesenchymal transitions is given by Hay. The following reviews of particular biological systems showing epithelial-mesenchymal transitions refer repeatedly to three elements that differentiate the two cell states. We have therefore assembled specialised reviews on the molecular nature of these three elements: Firstly, the mechanisms whereby cells interact adhesively with their cellular neighbors; details of such cell-cell adhesive molecules are described by Prieto and Crossin. Secondly, the mechanisms whereby cells adhere to their noncellular surroundings; details of the integrin class of extracellular matrix adhesion molecules and their ligands are supplied by Garratt and Humphries. Thirdly, the complex interactive network of structural and signalling linkages between these cell-surface adhesive molecules and the intercellular structural and functional organisation of the cytoskeleton are elegantly presented by Geiger, Yehuda-Levenberg and Bershadsky. These three contributions form, in essence, a reference section for the reviews to follow.

Two archetypal embryonic epithelial-mesenchymal transitions, gastrulation in mammals and neural crest morphogenesis, are then reviewed by Viebahn and by Duband, Monier, Delannet and Newgreen, respectively.

In a following volume new ideas on the molecular control of epithelial-mesenchymal transitions via kringle domain proteases are given by Théry and Stern. Details of another epithelial-mesenchymal transition in embryogenesis are supplied by Markwald, Eisenberg and Sugi in heart morphogenesis. Conversions in the reverse direction, mesenchyme to epithelium, are also known: Davies describes such events in kidney development. Recent studies of implantation of the mammalian blastocyst have allowed this to be reinterpreted as an event similar to classical epithelial-mesenchymal conversion; this is described by Vicovac and Aplin.

The creation of a new cell state, mesenchyme, allowed an enormous step in the order and complexity of the meta-zoan body plan to be realised. However, this creation is a two-edged sword, for if such cellular conversions occur in an unregulated way and are coupled to abnormalities of cell growth, metastasis occurs in previously benign cancers. This most feared of disorders is explored by Birchmeier and Birchmeier.

Finally, the molecular nature of epithelial-mesenchymal transitions can be examined in highly accessible and controlled tissue culture systems. This potentially allows for a greater range and depth of experimental manipulations, which may reveal fundamentals of the process obscured by system-specific details in natural examples. These model systems are described by Boyer, Vallés and Thiery.

The editors sincerely thank all the contributors for these valuable reviews.

© 1996 S. Karger AG, Basel