Introduction

J.

Joachim

Kühn

Institut für Virologie der Universität Köln, Deutschland

Dr. Joachim Kühn, Institut für Virologie der Universität zu Köln, Fürst-Pückler-Strasse 56, D-50935 Köln (Deutschland)

The ability to establish long-lasting, in many cases lifelong, persistent infections has evolved in members of various viral families in order to ensure effective horizontal and vertical spread within the human population. Depending on the viral agent itself as well as on host factors, the majority of persistent infections will cause only a few symptoms or will remain completely asymptomatic; however, they may also be associated with the development of neoplasia or severe organ dysfunctions in a significant percentage of immunocompetent individuals. In the immunocompromised host, the situation dramatically changes, and reactivated persistent viral infections, e.g., by human cytomegalovirus, frequently cause severe, life-threatening disease. Elucidating the viral mechanisms that lead to the establishment and maintenance of the persistent state is crucial for our understanding of the pathogenesis of these clinically important infections and for the development of effective diagnostic strategies.

The laboratory diagnosis of persistent viral infections has to meet special requirements. Thus, quantitation of the extent of virus replication may become necessary for reliable discrimination between anamnestic and persistent or reactivated latent infections.

Furthermore, direct demonstration of organ involvement, e.g., of the central nervous system or certain visceral organs, differentiation of viral genome types, subtypes and individual strain variants as well as monitoring of the effectiveness of antiviral therapies will become increasingly important.

From what has been said above it is easily conceivable that our present routine measures are far from being perfect for the laboratory diagnosis of most clinically important persistent viral infections.

This special issue provides an overview of recent findings regarding the pathogenesis and diagnosis of persistent infections caused by herpes simplex viruses, hepatitis C virus, cytomegalovirus, human papilloma viruses, human polyoma viruses and human T-lymphotropic virus type I. It is hoped that this compilation of reviews will be helpful for clinicians and microbiologists involved in the diagnosis and treatment of persistent viral infections, and will stimulate investigators interested in their molecular biology, immunology and pathogenesis.

KÄHGEII

E-Mail karger@karger.ch Fax+ 41 61 306 12 34 <http://www.karger.ch>

© 1997 S. KargerAG, Basel 0300-5526/96/0393-0139\$ 10.00/0