Introductory Remarks

W.	Wolfgang	Schwarz
----	----------	---------

This issue of Cellular Physiology and Biochemistry presents a series of short reviews on recent developments of structure and function of P-type ATPases. These enzymes represent a class of integral membrane proteins that transport cations across cell membranes. A common characteristic is that they become temporarily phosphorylated at an aspartate residue during the transport cycle. Another feature that allows to distinguish P-type ATPases from other ion-motive ATPases is the fact that vanadate specifically inhibits the enzyme activity.

Figure 1a shows a hypothetical phylogenetic tree of the catalytic subunits of P-type ATPases based on similarities between pairs. These polypeptides are visibly grouped and can be classified with respect to the transported cations. The highest degree of homology is found between the Na⁺/K⁺- and HVK⁺-ATPases. Among the Ca²⁺-transporting ATPases, the pumps found in sarcoplasmic reticulum (SERCA) are closer to the Na⁺/K⁺- and H⁺/K⁺ pumps than to the Ca²⁺ pumps of plasma membrane (PMCA). The species marked in bold print are selected examples for the different groups of pumps that are dealt with in this series of reviews including the bacterial K⁺-transporting ATPase Kdp. Despite the diversity in amino acid sequence, all these ATPases exhibit regions that are highly conserved during evolution and which can, therefore, be believed to be characteristic for P-type ATPases. This includes the region around the acceptor aspartate for the γ -phosphate of ATP (Asp376LysThrGlyThr; see fig. 2b, numbering for the Na⁺/K⁺-ATPase of *Torpedo*) and the region Thr219GlyGluSer (fig. 2a), which is possibly essential for phosphatase activity. Another conserved region is the sequence ThrGlyAsp716Gly; Asp716 as well as the conserved Lys507 and Lys697 are assumed to be in close proximity to the ATP binding site and to be involved in the formation of the binding pocket. An interesting additional feature is a highly conserved region around Pro333, which was suggested to stabilise cation binding.

Several of the P-type ATPases are composed of a single polypeptide of about 100 kD like the Ca²⁺-ATPases of plasma membrane (reviewed by Penniston and Enyedi) or of sarcoplasmic reticulum (reviewed by Inesi et al.). The Na⁺/K⁺-ATPases (Vasilets and Schwarz, Schmalzing and Gloor) and the H⁺/K⁺-ATPases (Klaassen and De Pont) have been

human a1 pig a1 sheep a1 horse a1 rat a1 chicken
a1 Xenopus a Bufo a teleost a

1

H

Z
I

rat a2 chicken a2 Q

H

+

I

human a3 rat a3 chicken a3 ^

+

i

100% Torpedo a
Drosophila a Artemia a

human a pig a

rabbit a rat a X
+

62(26) %

1

s

7N

baker's yeast ATPase 1

chicken 1 28(35 %) rabbit 1

2(2) rat 2(2) rabbit 2(2) human 2(1) rat 2(1) pig 2(1) rabbit 2(1) human 2(2) pig

~

CA

m

O>

rat 3
 Drosophila
 SR
 rat 1A human IB pig IB rat 2 human 4
 r
 -V
 1
 Mzz
 o
 >

23(35)%

baker's yeast fission yeast

-T
 I
 i/W% > 5%
 Streptococcus
 E coli B

o

+

1a

Fig. 1. Phylogenetic tree determined by the program CLUSTAL of PC/Gene for the catalytic subunits of P-type ATPases (a) and for the β - or C-subunit (b). Those species that are particularly addressed in this issue are given in bold, and the relative degree of homology in amino acid sequence is indicated; numbers without brackets represent identity, those with brackets similarity.

Fig. 2. Comparison of amino acid sequences of catalytic subunits of different P-type ATPases. Comparison is presented for three different regions with a high degree of homology and that are, therefore, likely to be involved in forming the ATP binding pocket. The enlarged Asp (D) in b represents the phosphorylation site. The numbers above the arrows refer to the α -sub-unit of the Na⁺/K⁺-ATPase of Torpedo electroplax.

78

Schwarz

Introductory Remarks

62(26)%

62(26)%

61(30)%

human mouse rat

pig

dog

sheep

chicken

Q

Torpedo β β 2 β 2

37(34)%

37(33)%

61(26)%

human mouse rat β 2Xenopus β 3

pig

rabbit rat

7Z,

Artemia β Na⁺/K⁺

12(30)%

E. coli

KdP

Torpedo Na/K-a pig H/K-a rabbit SERCA1 human PMCA 4 E.coli KdPB
219 ↓

II	SACSC--KVDNSSL TGES EPQSRSP	232
IL	QAQGR--KVDNSSL TGES EPQTRSP	242
IL	SIKSTTLRVDQSIL TGES VSVIKHTEPV	194
IL	IQGNDLKIDESSL TGES DHVKK	248
VIEGGAS	VDESAI TGES APVIR--ESG	170 a

Torpedo Na/K-a pig H/K-a rabbit SERCA1 human PMCA 4 E.coli KdPB
Torpedo Na/K-a pig H/K-a rabbit SERCA1 human PMCA 4 E.coli KdPB
376

I
EAVETLGSTSTICS DKTGT LTQNRMTVAHM 391
EAVETLGSTSVICS DKTGT LTQNRMTVSHL 401
PSVETLGCTSVICS DKTGT LTTNQMSVCKM 366
DACETMGNATAICS DKTGT LTMNRMTWQA 4 80
RAVEAAGDVDVLLL DKTGT ITLGNRQASEF 322 b
Λ * Λ * ΛΛΛΛ ***** Λ* ΛΛΛ Λ Λ
716

I
EGC QRQGAIVAV TGDG VNDSPALKK 726
ESC QRLGAIVAV TGDG VNDSPALKK 737
E YLQSYDEITAM TGDG VNDAPALKK 713
KGIDSTVGEHRQWAV TGDG TNDGPALKK 795
RQY QAEGRLVAM TGDG TNDAPALAQ 528 C
ΛΛ^{3/8}Λ ***** Λ**Λ^{1/8}Λ

shown to be αβ-heterodimers with a catalytic α-subunit of about 100 kD and a glycosylated β-subunit (about 30-35 kD for the unglycosylated polypeptide). A third form of P-type ATPases reviewed in this issue is the K⁺-transporting Kdp-ATPase of bacteria that is composed of three major subunits (A, B and C; reviewed by Altendorf and Epstein). Interestingly, while the catalytic subunits of the Ca²⁺- and X⁺/K⁺-ATPases bear both the ATP and phosphorylation sites as well as the cation binding sites, for the Kdp-ATPase ATP binding and phosphorylation occurs on the B-subunits, cation binding on the A-subunit. For the α- or catalytic subunits, the N- and C-termini have been demonstrated to have cytoplasmic orientation; as a consequence, an even number of membrane segments has to be assumed. In the presented reviews, the authors favour 10 segments for all types of pumps except for the B-subunit of the Kdp-ATPase.

The β-subunit of the X⁺/K⁺-ATPases is assumed to have a single transmembrane segment with a cytoplasmic N-terminus. This subunit is believed to play an important role in the correct assembly and function of the enzyme. The C-subunit of Kdp-ATPase has a similar orientation and seems to play an equivalent role and is, therefore, included in the phylogenetic tree (fig. 1b). In spite of these functional similarities, homology with respect to amino acid sequence is only about 10-15% compared with the β-subunits of Na⁺/K⁺- and H⁺/K⁺-ATPases. The function of β-subunits is explicitly addressed in the contribution by Schmalzing and Gloor.

Several of the above-mentioned features are discussed in more detail in the following reviews. We intend this selection to inspire comparison of P-type ATPases, which may help in the understanding of structure-function relationships.

80

Schwarz

Introductory Remarks