Book Review

John Terborgh

Five New World Primates

A Study in Comparative Ecology

Princeton University Press, Princeton, N.J., 1983 260 pp.; US\$ 40.00 (cloth), US\$ 13.50 (paper)

ISBN 0-691-08337-1 (cloth), 0-691-08338-X (paper)

The dominant conceptual framework in field studies of primate ecology and of social behavior and ecology from the mid-1960s to the present has been socioecology, in which many of the basic variables are macro-variables (e.g., mean day range, group home range, proportions of food types in the diet) that are several steps removed from the individual animal living in its habitat. Natural selection operates on behavior, not on morphological or physiological traits directly, and not on summary statistics of the sorts just mentioned. Food is extremely important for survival and reproduction, so the search for and acquisition of food are fundamental in understanding ecology and social behavior. Individual animals search for and acquire food by means of foraging behavior. Thus, a science of foraging ecology that focuses on behavior as a process is preferable to 'feeding ecology', which is often close to 'diet ecology' with relatively little emphasis on process as might be implied by the gerund 'feeding'.

Comparative ecological studies of Old World primates have been essentially socioecological in outlook; they have yielded invaluable basic information on the primate components of communities, and they have generated a large array of interesting hypotheses. Terborgh's research is the first comparative study that is foraging ecology in the sense used here. In addition to being a highly significant analysis of five species of primates, it furnishes conceptual and methodological guidelines for testing hypotheses generated from correlations of macro-variables. The study site of Cocha Cashu in Peruvian Amazonia is inhabited by eleven species of primates in an area where several different habitats are intermixed. The five species studied intensively by the time this book was written were *Cebus apella*, *C. albifrons*, *Saimiri sciureus*, *Saguinus imperator*, and *S. fuscicollis*, ranging in size from 400 g to 3 kg.

Most of *Five New World Primates* concerns the ways that animals with different capacities deal with diverse characteristics of fruit and prey to produce different foraging behaviors. (Here I use 'foraging' in the broadest sense, noting that for Terborgh feeding means plant parts and foraging means insect prey.) The study is placed in perspective by Terborgh's lucid explanation of the basic problems faced by mammals with different diets: in small herbivores the rate of digestion limits ingestion (so they rest a good portion of the day), while in insectivores ingestion rate normally cannot exceed digestion, and searching for prey is the primary constraint. *Omni-vores* are animals that consume some animal prey and some plant parts other than leaves, and include the five primates reported on here. They must compromise: 'A need to obtain protein in the form of insects puts major demands on the time budget, while a need to obtain carbohydrate in the form of fruit largely dictates the use of space.'

Here I can indicate only a few of the significant points of this book. Fruit production and phenology were assessed by monitoring traps made of plastic bags suspended from wire hoops

just above the ground. The results show that different habitats vary in their schedules of fruiting, and that when fruit is scarce it also occurs in fewer patches that are farther apart than when fruit is more abundant. This kind of observation on food dispersion would benefit numerous studies, for example, of locomotion, and was achieved with relatively little effort. The sampling schedule for observations of primates involved periods of about 20 continuous days for each species in each of four seasons, instead of the usual 5 days per month used by many investigators (including this reviewer). The former scheme helps reveal patterns that are not discernible in briefer periods.

The two species of *Cebus* and *Saimiri* spend almost half their day searching for animal prey, far more than Old World monkeys that have been studied. Although it is very difficult to explain why animals range as they do, Terborgh suggests: 'Among our set of species, *Saimiri*, and to a lesser degree *Cebus*, make extended forays for days for which the only discernible purpose is insect foraging. The tamarins, in contrast, appear to do most of their insect foraging while en route between fruit trees.'

228

Book Review

With regard to plant foods (mostly fruit), *Saimiri* and *Cebus* have much more diversified diets than the tamarins. Both species of *Cebus* use palm resources a great deal, particularly *C. apella*, and they deal with the tough nuts of *Astrocaryum* in different ways: *C. apella* has a more massive masticatory apparatus and can crack the nuts readily between its teeth, whereas *C. albifrons* must search for nuts that have been attacked and weakened – but not yet completely consumed – by beetle larvae. During the dry season *C. albifrons* and *Saimiri* specialize on figs to a considerable extent. Fig fruit at Cocha Cashu occurs in large but rare patches, distributed irregularly in space and time, probably explaining the huge home ranges of *C. albifrons* (about 150 ha) and *Saimiri* (over 250 ha). Old World primates of comparable body size (or group weight) have much smaller home ranges and the differences cannot be explained readily. Terborgh speculates that perhaps fruit is less abundant or more highly seasonal in the New World.

Saimiri forms mixed-species associations with both species of *Cebus*, and Terborgh is able to show convincingly that it is *Saimiri* that actually joins *Cebus* and benefits more from the association than does *Cebus*. The benefits are assistance in finding ripe fruit crops and predator detection. The evidence for the role of predation in primate sociality provided in this book is a major contribution to the current debate

about the adaptive significance of grouping in primates. The two species of *Saguinus* associate, but the pattern is very different from that of the larger species. A permanent alliance of one family group of each species of tamarin jointly defends a common territory against similar mixed groups. Other benefits – and perhaps costs – of this highly unusual arrangement are difficult to unravel, and one can only hope that Terborgh plans more work on the problem.

Finally, Terborgh attempts a comparison with Old World primate communities, noting that the range of body size is much greater in New World monkeys, and that the species at Cocha Cashu consume more nectar and insects, and less leaves, than the higher primates that have been studied in the Old World. Some of these differences likely arise from contrasts in climate and vegetation, and from historical differences, e.g., there is no foregut fermenter among New World monkeys. A more satisfying interpretation of differences between communities must await application of Terborgh's sort of foraging ecology to diverse Old World and New World primates.

Five New World Primates is certain to prove extremely useful to seasoned investigators, and its concise and lucid exposition of major issues in primate ecology makes it invaluable for undergraduate courses in primate biology, animal ecology, and animal behavior.

John G.H. Cant, New Haven