Special Section

Am. J. Nephrol. 1986;6:469

Medical Art

D.	Daniel	Casellas
----	--------	----------

-ΛinMH

$$1/8 1/8 1/8 1/8 f_7/8 y^{\wedge \wedge * 3/4}_j \dot{w}$$
$$\therefore \therefore 5/8$$
 $\hat{A}E$
$$^{\wedge}E) \text{ illitfKP} \gg \text{IBBBBBp}\text{lllllll}^{\wedge}\text{MBB}\S^{1/8}_{1/8}$$

$\frac{1}{8} \cdot \frac{1}{8} = \frac{1}{64}$

■ ..■■.■'■■ ..

$${}^1_8\text{S}'\text{R}\phi^{*}\text{''''''}\ddot{\text{A}}\text{J}..$$

■:--...-■' ■' ■■ ■

$$\frac{7}{8} \wedge'' \ddot{A} \frac{1}{8} \& SB \gg t^{\wedge},$$
 $\frac{7}{8} *$

“■■’■-’ :: ?

 $\leq \frac{7}{8}B$

-...,,,* , TM ■ , “7₈-3₈7₈.c³₈|7₈]r S

«vJÉÉ^H

$$5^{1/8} \sim 1$$
$$1/8, 1/8, 1/8, 3/8, \dots,$$
$$\wedge./^{\circ}\mathbf{S}^{1/8\wedge 1/8^3/8^5/8^3/8}_{\mathbf{C}}^{1/8^7/8^7/8}$$

HP!

■ ■ ■ ■ ■

WÈÈÈÈ

BHm^mHhe

Scanning electron micrograph of the glomerular vascular pole as seen from inside Bowman's capsule of the rat kidney. The entry point of the afferent and efferent arteriole was exposed by removing the glomerular tuft. The narrow area between the vessels marks the transition between extra- and intraglomerular mesangium. $\times 2,000$.