Preface

N.	Normal	Ofsthun
T.R.		Shockley

McGaw Park, Ill

During peritoneal dialysis, fluid loss from the peritoneal cavity reduces net ultrafiltration, on average, by 40–50%. Lack of detailed knowledge about the pathways for this fluid loss has led to considerable scientific debate. This special issue of Blood Purification contains the proceedings of an industry-sponsored symposium entitled 'Lymphatic and Non-Lymphatic Fluid Loss from the Peritoneal Cavity' which was held following the annual meeting of the American Society of Nephrology in Baltimore in November of 1991. Clinical nephrologists, physiologists, lymphologists, engineers, and other scientists were brought together to discuss fluid loss from the peritoneal cavity. Papers were solicited on the following topics: (1) identification of lymphatic and nonlymphatic pathways for fluid loss, (2) methods for measuring fluid loss, (3) dependence of fluid loss on other parameters (intraperitoneal pressure, state of consciousness, drugs, etc.), and (4) effect of fluid loss on net ultrafiltration in peritoneal dialysis.

As might be expected when convening scientists from several disciplines, participants brought with them not only differing experimental and analytical methods, but also differing research objectives, viewpoints, and terminology. For example, most nephrologists who practice peritoneal dialysis (PD) have historically been concerned with total fluid loss from the peritoneal cavity because net ultrafiltration affects the efficacy of PD therapy. As a result, a major goal of PD clinical research has been to understand ultrafiltration kinetics sufficiently to prevent or treat ultrafiltration failure. PD clinicians have often used the term 'lymphatic absorption rate' to describe the total rate of fluid loss from the peritoneal cavity, and have not distinguished among the various pathways out of the peritoneal cavity.

While many clinicians have focused on macroscopic effects such as the dependence of calculated fluid loss rates on intraperitoneal volume and dialysate tonicity, others have focused on fluid loss at the microscopic level. For example, physiologists and engineers have studied fluid movement at the tissue level (i.e., through the interstitium into capillaries and lymphatic vessels) with the hope of elucidating the mechanisms for fluid and solute transport from the peritoneal cavity.

Similarly, lymphologists have focused on fluid flow into and through lymphatic vessels. By cannulating lymph vessels in animals, they have measured flow rates through lymphatics that drain the peritoneal cavity, thereby increasing our understanding of the role lymph-

113

phatics play in fluid loss during PD. Unfortunately, cannulation techniques cannot be applied to humans. Other researchers have turned to mathematical modeling to understand fluid and solute transport in peritoneal dialysis, depicting the peritoneal 'membrane' as a membrane permeable to certain solutes. While these models adequately simulate PD and provide estimates for fluid loss from the peritoneal cavity through the various pathways, they cannot precisely delineate the experimentally observed pathways for this transport.

One of the primary accomplishments of this symposium was the frank discussion of the terminology used to describe fluid movement during PD. Participants agreed that any term which includes the word 'lymph' (e.g., lymph flow, lymphatic drainage, lymphatic absorption rate) should be used only when describing the fluid flowing through lymphatic vessels, and the term 'fluid loss' should be used when describing the total fluid loss from the peritoneal cavity via the various pathways. Much confusion could be avoided if investigators studying fluid movement during PD would make a conscious effort to eliminate unorthodox use of the term 'lymph'.

To give the uninitiated reader the background necessary for assimilating the material submitted by individual participants, we have written a short tutorial. This tutorial describes the pathways for fluid loss from the peritoneal cavity and discusses the techniques for estimating flow rates through lymphatic and non-lymphatic pathways.

As editors of these proceedings, we would like to thank Dr. Lee Henderson and Baxter Healthcare for sponsoring this symposium. We hope that the monograph accurately reflects the content of individual presentations and the discussion that followed. Finally, we would like to thank all the participants for their contributions to the symposium, for the value of this monograph lies in the strength of the individual papers.