Introduction

W.

Rafflenbeul

Abteilung Kardiologie, Medizinische Hochschule, Hannover, FRG

Prof. Dr. med. W. Rafflenbeul, Abteilung Kardiologie, Medizinische Hochschule Hannover, Postfach 61 01 80, D-3000 Hannover 61 (FRG)

This volume represents the Proceedings of the International Symposium on Mononitrates held at the Royal Society, London, in June 1986. The papers presented reflect the spectrum of knowledge that has accumulated in recent years on mononitrates in the management of patients with coronary artery disease or heart failure.

Over many years of successful clinical application, organic nitrates have become an established therapeutic regimen in patients afflicted with angina pectoris. Nitrates unfold their antianginal properties predominantly by unloading the heart following venodilation. In addition, dilation of large epicardial coronary arteries and particularly attenuation of stenosis severity following nitrate application may enhance oxygen delivery to the ischemic myocardium [1, 2].

Some unfavorable characteristics of the widely used dinitrates, like variable bioavailability, short half-life and particularly the extensive first-pass metabolism, ignite the search for more consistent substitutes [Tau-chert and Jansen, this volume]. In this regard the vascular-active metabolites of isosorbide dinitrate, the mononitrates isosorbide 2-mononitrate and particularly isosorbide 5-mononitrate (IS 5-MN) are superior to the parent substance because they exhibit almost complete oral absorption, a relatively long half-life and a low intersubject variability of pharmacokinetics [Chasseaud, this volume]. These properties favor the mononitrates particularly for long-term therapy of symptomatic coronary artery disease.

However, during this Symposium, major interest was focused on its acute as well as on its long-term antianginal mechanism and efficacy. After acute application, 20 mg of IS 5-MN reduces systolic and diastolic blood pressure and the rate-pressure product as well as end-diastolic and end-systolic left ventricular volumes. A persistent reduction of preload could be demonstrated after 2 months' therapy with 20 mg IS 5-MN three times daily. Increasing doses of slow-release IS 5-MN show a dose-dependent effect on reduction of ST-segment depression as well as frequency of anginal attacks. With a single high-dose (50 mg), slow-release IS 5-MN capsule daily, the pronounced anti-ischemic effect could be maintained over a 12-month period without obvious signs of tolerance development.

Continuous Holter monitoring becomes an established procedure to demonstrate ischemic attacks – symptomatic or asymptomatic –

2

Rafflenbeul

tomatic – during daily life activities in patients with coronary artery disease. In addition, the efficacy of an antianginal treatment regimen can be documented. Although a substantial decrease

of anginal attacks after application of mononitrates can be registered with Holter monitoring, only half of the patients became completely free of isch-emic episodes. A similar pattern of response was observed with other antianginal drugs like the calcium antagonist nifedipine. In some of those patients only partially responsive to mononitrates, additional nifedipine may further reduce the frequency of isch-emic episodes detected by Holter monitoring.

Mononitrates not only prevent myocardial ischemia elicited by effort, but are also capable of abolishing coronary vasoconstriction provoked with ergonovine. Coronary vasoconstriction, particularly in coronary artery stenoses, is regarded as a major patho-physiological mechanism responsible for spontaneous angina pectoris.

In summary, the data presented in this volume clearly demonstrate that the antianginal activity of mononitrates is particularly beneficial for patients with chronic angina pectoris. Long-term therapy with this new compound alleviates subjective and objective signs of effort-induced myocardial ischemia. Based upon this experience, the mononitrates are characterized as a alternative to conventional antianginal drugs in the treatment of symptomatic coronary artery disease.

References

Rafflenbeul, W.; Urthaler, F.; Russell, R.O.; Lichtlen, P.; James, T.N.: Dilatation of coronary artery stenosis after isosorbide dinitrate in man. Br. Heart J. 43: 546(1979).

Rafflenbeul

W.; Lichtlen

P.R.: Quantitative coronary angiography: evidence of a sustained increase in vascular smooth muscle tone in coronary artery stenosis. Z. Kardiol. 72: suppl. 3 p. 87 (1983).