Introduction

L.H.

Opie

Heart Research Unit, Department of Medicine, University of Cape Town, Observatory, Cape Town, South Africa

Gone are the days when congestive heart failure belonged to the realm of poorly understood, poorly treated, and generally neglected cardiac diseases. In view of the significant advances recently developed in the diagnosis and management of heart failure, the present Symposium is highly appropriate.

The first session is devoted to basic approaches to heart failure. Prof. Dalla Volta carefully analyses the multiple causes of myocardial dysfunction in heart failure. His important point is that not all types of heart failure have the same etiology and according to the differences in etiology there are differences in mechanical function which, in turn, should be reflected in differences in the therapeutic approach. Dr. Francis goes on to discuss the importance of the extramyocardial features of heart failure including the changes in catecholamines and the renin-angiotensin system. It is generally felt that such changes may contribute to the adverse overall effects of congestive heart failure. Dr. MacNee discusses right ventricular function in cor pulmonale. Frequently, the right ventricular component to heart failure is ignored, but it may be most important from

the symptomatic and therapeutic point of view. Thereafter, Drs. Fletcher and Bulpitt discuss the quality of life in congestive heart failure. No longer is it deemed adequate simply to achieve an improved hemodynamic picture in heart failure; the important point is whether the patient can achieve more physically and feels better. Although very few studies have had as their end-point objective and careful quality of life measurements, it is clear that this pattern of thinking is going to become increasingly important.

The pharmacological principles of treatment of heart failure are discussed by Prof. Kulbertus from Belgium. Besides diuretic therapy, the most important ways to go are the inotropic and the vasodilator avenues. An important and somewhat debatable question is whether the angiotensin-converting enzymes act purely as vasodilators or also as specific inhibitors of the harmful vicious circle involving renin, angiotensin and peripheral vasoconstriction. Dr. Taylor then considers how to treat acute heart failure. Here the whole approach is different because the hemodynamic changes become all important and, frequently, the therapy is highly effective. Thereafter, Prof. Hugenholtz makes a

2

Introduction

clear case that early thrombolytic treatment can prevent later ischemic congestive heart failure. The final session of the meeting is devoted to controversies in treatment of congestive heart failure. Prof. Poole-Wilson argues that digitalis is more dead than alive and should seldom be used. Dr. Franciosa argues the positive case for vasodilators and feels that they are undoubtedly

useful. Prof. Hampton considers the case for the use of angiotensin-converting enzyme inhibitors and although there are powerful data supporting their use and benefit, it is not entirely clear when they should be introduced in the step-care of congestive heart failure. It may be that an increase in diuretic therapy is as effective as the addition of ACE inhibitors. Thereafter, Prof. Bussmann examines the question of the use of ibopamine, an inodilator which has theoretically attractive properties for use in congestive heart failure. He considers whether the hemodynamic changes achieved during acute therapy are sustained during chronic therapy. Finally, the paper by Opie assesses whether the 'ino-dilators' might be the way ahead in the treatment of congestive heart failure. Among the inodilators, ibopamine has attractive qualities and theoretically provides a beneficial hemodynamic profile together with an added diuretic component. With the ino-tropic effect, arterial vasodilation, and diuretic effects combined, it is argued that this agent deserves further evaluation for use in congestive heart failure.