Editorial

J.

Tooke

Exeter, UK

For many microcirculationists the driving force behind their endeavours is the goal of understanding the patho-physiology of human disease processes. The relaunch of the International Journal of Microcirculation: Clinical and Experimental as the official journal of the European Society for Microcirculation provides a timely opportunity to reflect on whether such observations on the microcirculation are being translated into clinical practice.

At a general level it can be argued that the description of microvascular derangement in common disease processes has raised the general awareness of this part of the circulation. In hypertension for example, the view is becoming accepted that it is insufficient to correct bra-chial artery pressure alone and there needs to be consideration as to whether a particular therapy corrects micro-vascular derangement, particularly the structural remodelling that occurs. In insulin-dependent diabetes means are emerging to correct the capillary hypertension that characterises this condition. Direct observation of white cell activation in reperfusion injury and an understanding of the nature of this process are leading to trials of antioxi-dants.

Perhaps the most productive feature of clinical physiological observation is the opportunity such functional analysis provides to propose plausible molecular or cellular patho-genetic mechanisms [1]. Arguably such a reductionist approach, i.e. examination of those possibilities that fit the physiological facts, is more likely to be fruitful than unfocused examination of myriad mechanisms in isolated cells that may or may not be involved in the intact organism. This assertion may be illustrated by considering our understanding of the development of human diabetic microangiopathy: in vitro studies have revealed increased blood viscosity and reduced blood cell deforma-bility which understandably have been suggested as causes of reduced microvascular blood flow. However,

direct microvascular observations confirm the fact that microvascular pressure and flow are increased in most tissues early in diabetic life, despite the adverse blood Theological measurements. In fact, the challenge in diabetes is to clarify why hyperglycaemia provokes precapillary vaso-dilation in so many microvascular beds and how shear stress and/or microvascular hypertension is transduced by endothelial cells, resulting in the production of extracellular matrix proteins such as type-IV collagen. The answers to these questions and many like them will come from the targeted application of the newer sciences of molecular and cellular biology, and already the first tangible therapeutic benefits from such an approach are being realised in other disease states. For example, attempts to block the overproduction of nitric oxide in septic shock are being made and monoclonal antibodies against cell adhesion molecules are being tried in attempts to correct transplant rejection.

The future then holds prospect of specific therapies for diseases that have an impact upon the microcirculation: the acquisition of such treatments is likely to be accelerated if physiologists and

pharmacologists establish close working relationships with both clinicians and molecular biologists.

Such a marriage of basic and applied skills is not only the ethos of the European Society for Microcirculation, but is also captured in the title of the International Journal for Microcirculation: Clinical and Experimental, which aspires to publish the key papers describing the product of such endeavours in the coming years.

Reference 1 Tooke JE: Microvascular physiology and its clinical relevance with special reference to diabetes. Q J Med 1992;83:567-573.