Foreword

Disease of the urinary tract of the child has been recognised and described since the days of Hippocrates more than two thousand years ago. His idea that the cause of stone in infants was wine in the diet of nurse and child may provoke merriment, but the grin should be wry. Little indeed is known, even in 1973, concerning the pathogenesis of the urolithiasis which is widespread in the children of the Middle East, Africa and India. The problem of enuresis has puzzled physicians over the century, but the post hoc bell and pad of today differs little from the proper hoc pig's bladder of yesterday. The classical facies of the disease now called the idiopathic nephrotic syndrome of childhood (INS) must have been known since before the days of Hippocrates, but the earliest written description seems to be that of Roelans in 1484. By 1722 (a century before Bright snatched the headlines) Zuinger had linked the natural history of INS from the oedematous nephrotic through to the prodigious thirst of terminal renal failure. Study of tubular defects are relatively modern. The classical 'inborn errors' described by Garrod at the turn of the 20th century included study of cystinuria. The simplicity of the early works on this topic has now gone and extended to an almost incomprehensible galaxy of polysyllabic tubular anomalies known today.

As the 20th century entered its third decade an ever increasing interest arose in various aspects of the disorders of the urinary tract. This was the age of adventure. Just as in the 19th century the skill, daring (and dare we say recklessness) of surgeons led to thymectomy of out-patients to 'cure' rickets, so did Edehbol's bilateral decapsulation of kidneys for nephrotic syndrome become the 'in' thing of the 1920s. It is a matter of continuing astonishment that the figures published from Glasgow by Campbell in 1929 for decapsulation (carried out on grossly oedematous children long before sulphonamides)

68

Foreword

showed a 2-year recovery rate in 13 of 26 children. Having personally traced one of these children, thereby meeting and investigating a fecund man of 49 whose kidneys were decapsulated 47 years ago, one could not but admire both his scars and continuing health! It is interesting that Campbell realised that the stress of operation was the likely cause of improvement and not the decapsulation. From Glasgow, Findlay in 1922 drew attention to the problems and dangers of pyelonephritis in the newborn. The problem remains today and, if better understood, is far from solved. By now the pace was quickening in many parts of the world. Fanconi in Switzerland, Smith in the United States and McCance in England were in the forefront in the study of function and physiology which began to rank alongside the study of disease and the dramatic surgical tours de force.

With the 1939-1945 war came a sharp change in the pattern. The European lead in scientific medicine, paediatrics and nephrology dwindled away and the running was taken up in North America. A range of new names such as Heymann and Barnett sprang to the fore as the nephrotic syndrome became the main focus of attention. Quite suddenly interest in paediatric nephrology seemed to expand all over the world. By 1966 many younger men were coming into the field of

paediatric nephrology in a variety of centers but the anomalous situation had arisen that, whilst many European paediatric nephrologists were well known at meetings in the United States, and many paediatric nephrologists in the USA were familiar with the men of Europe, inside Europe there was little contact between paediatric nephrologists from differing countries, let alone East and West.

In 1966, Harmen Tiddens of Utrecht came to Glasgow where in an atmosphere of straight malt (spiritus frumenti) the European Society for Paediatric Nephrology was conceived. 23 European countries were invited and sent representatives to Glasgow where the inaugural meeting was held in September 1967. The Foundation Council included Prof. H. Bickel, Germany; Prof. E. Gautier, Switzerland; Prof. N. Hallman, Finland; and Prof. J. Winberg, Sweden; in addition to the founders (Prof. H. A. W. Tiddens and myself). One year later, the American Society of Paediatric Nephrology was formed followed by the Japanese Paediatric Nephrology Society. In parallel with these events another International Group was developing. In 1965, Henry Barnett was invited to give a lecture to the British Paediatric Association. In discussion afterwards the idea of the International Collaborative Study of Kidney Disease in Children sprang into being. The original Directors included Prof. N. Hallman of Helsinki, Prof. H. A. W. Tiddens of Utrecht and myself from Europe; Prof. Kobayashi of Japan, Prof. G. Gordillo of Mexico, and Prof.

Foreword

69

H.L. Barnett as Chairman. A central office was set up with Drs. C. Edelmann, I. Greifer and M. Abramowicz at the Albert Einstein College of Medicine, New York, USA. The concept of many centres on an international basis agreeing to conform to protocols (in order to obtain results in the study of uncommon illnesses more rapidly) had often been tried, but this time it worked! All clinics were asked to provide retrospective data from their records. Those who did so competently, on 541 cases, were accepted. Those who did not were out!

For the remainder a controlled double-blind prospective study of azo-thioprine treatment for children with INS was the first task. The criteria, management and treatment were laid down for further trials including cyclo-phosphamide treatment of INS. Each biopsy specimen was read 'blindly' by four pathologists in France, England and USA (2). A standard terminology was developed and described in the Lancet in 1970. A set of 21 teaching slides illustrating the classification was prepared on 35-mm slides, together with descriptive literature by Prof. J. Churg, New York, USA and these were widely distributed throughout the USA, Europe and elsewhere. Already, 400 children with INS are being followed prospectively with histology 'agreed' and the outcome is being vetted in detail. The prognostic significance of focal segmental glomerulosclerosis was recognised by the ICSKDC in 1970 and was re-discovered very rapidly at many centres. When cyclophosphamide gonadal toxicity became a problem the framework of ICSKDC together with the National Kidney Research Foundation of New York rapidly set up a meeting in New York in June 1972, published the best advice available and set up a register of cyclophosphamide gonadal toxicity for children.

Nor did the momentum cease. In December 1968 in Guadalajara, Mexico, Prof. Gustavo Gordillo held the first World Symposium of Paediatric Nephrology to be followed by the second, organised by Prof. Pierre Royer in 1971 in Paris, France. The third will be in Washington, USA, in 1974 with Dr. Phil Calcagno as President.

The development of National, Continental and International Societies of Paediatric Nephrology seemed to set fair the barometer for further progress in paediatric nephrology. This indeed is true

although basic problems still underly. Horizontal integration with the remainder of paediatrics still vies with vertical integration with the nephrology of adults.

By 1970 things changed again as haemodialysis and transplantation became practicable for children and even infants. The threat arose, and remains, that paediatric nephrologists might be divided into three types: those working on parenchymatous disorders, those working on urinary tract infections and

70

Foreword

those working on chronic renal failure. The same problems beset nephrology for adults. Will such tripartite state of nephrology proceed to the development of further vertical streaming at the expense of a more general paediatric and renal approach ? Nobody knows, but since the only thing that is constant is change, let us hope with Candide, that it will be for the best. Tout se change; toujours la même chose?

Gavin C. Arneil, Glasgow