General Discussion

S.M.

Shalet

S.M. Shalet, MD, Christie Hospital and Holt Radium Institute, Department of Endocrinology, Wilmslow Road, Manchester M20 9BK (UK)

Shalet: I think that it would now be appropriate to return to the earlier theme of the session. There is general agreement that the brain tumour children lose a great deal of potential growth partly due to radiation-induced GH deficiency. GH therapy in the brain tumour children is less controversial now and the area for discussion concerns the optimum time to start GH treatment. More confusion and controversy have been generated over the GH requirements of the ALL child. Some groups have found no adverse effect of chemotherapy and CNS irradiation on final height. Other groups, including our own, have noted a modest adverse effect on growth, but as final height is unknown in the majority of these children the possibility of impaired pubertal growth may mean that final height loss is substantial in a proportion of children. Finally, there is a third group of ALL children, illustrated by the Australian experience, in whom there is a much greater retardation of growth.

In the first group there is clearly no indication for GH therapy; in the second group the exact requirements are ill-defined, but I believe that only a minority of these children require GH therapy. However, the Australian workers in the final category have treated the majority of their ALL children with GH.

I do not feel that these different results are as conflicting as they appear. If one remembers the old literature on GH secretion in the ALL child, there was great debate over whether or not radiation caused any abnormality of GH secretion. The debate was resolved when it was understood that changes in the radiation schedule regarding fractionation, which appear minor, had a profound influence on the results of GH tests. Similarly, if we look more critically at the radiation schedules and chemotherapy protocols it may be possible to understand the differences in height loss observed in various centres.

Kulin: One must not overly focus on small deficits in growth in children receiving cancer therapy. Behavioural effects must be evaluated carefully in mild statural compromise. Treatment of selected patients may produce more harm than gain psychologically. The cause of precocious puberty in these patients is of interest and suggests that facilitatory and inhibitory control of gonadotrophin secretion (or GnRH control) exists in man. Radiation appears to preferentially affect the inhibitory influence and thereby allow facilitatory effects to predominate. Further investigations in these individuals may, therefore, aid our still very unclear understanding of the cause of pubertal onset in man.

Law: When considering the administration of GH to children who have been treated for malignancy by irradiation and/or chemotherapy, one should consider the incidence of malignancy in other members of that family. The same caution should be taken in children considered for non-conventional GH therapy. After all, GH is a potential oncogen and acromegalic patients have an increased incidence of intestinal polyposis. Therefore, I advocate taking a careful family history in all the above-named patients.

Sanders: With respect to GH treatment in children with malignancy one has to keep in mind the real but unknown incidence of secondary malignancies. There has been an association of brain tumours and acute leukaemia and acute leukaemia with brain tumours. When an agent such as GH is available and it does not appear to increase the incidence of leukaemia in children with hypo-pituitarism then GH should be used to treat these radiation-induced GH-deficient children and data collected very carefully.

Shalet: Just to change the direction of the discussion it has been mentioned that the hypothalamus is more vulnerable to radiation damage than the anterior pituitary. Nonetheless, the pituitary may itself be directly damaged by irradiation. This is important when considering the potential use of a GRF depot preparation, when it becomes available, in children with radiation-induced GH deficiency. I would welcome Dr. Albertsson-Wikland's comment on this suggestion as her group has shown a decline in the GH response to GRF with time in radiation-induced GH deficiency. This might mean that the time scale for pituitary damage is rather different to that of the hypothalamus. Alternatively, the somatotrophs may remain intact but simply require priming with GRF.

Albertsson-Wikland: The mean GH response to a GRF test in CNS-irradiated children is about half of that in short healthy children. There is a reduced GH response to GRF with time after CNS irradiation. However, after CNS irradiation for ALL the pattern of GH secretion remains pulsatile and is characterised by distinct pulses of low amplitude with mostly very low trough levels in between the peaks. This implies that endogenous GRF and somatostatin are still present in irradiated ALL children, but does not distinguish between a hypothalamic and pituitary site of damage. Similarly, inadequate GH secretion during puberty could be explained by a failure of GH response to the action of oestrogen at either the hypothalamus or pituitary. If I have to speculate, then I believe that the somatotrophs of the irradiated child will respond to GRF if adequately primed, but we need more data in this area.

General Discussion

91

Czernichow: Intact vasopressin secretion is against hypothalamic sensitivity to X-ray therapy.

Shalet: I completely agree with your observation but not the conclusion. No case of radiation-induced diabetes insipidus has ever been reported. However, I interpret this observation to mean that the GRF-secreting neurones of the hypothalamus are more radiosensitive than those of the supra-optic and paraventricular nuclei.

van Vliet: The diminished GH response over time to GRF testing is not peculiar to irradiation-induced GH deficiency, so it is not an argument to postulate that irradiation can have effects on both GRF neurons and pituitary somatotrophs. The decreased responsiveness to exogenous GRF may result from 'disuse atrophy' of the pituitary somatotrophs after long-standing endogenous GRF deficiency and not necessarily from direct radiation-induced damage to the pituitary.

Stanhope: Dr. Shalet, as an adult endocrinologist perhaps you could help us with some information about the oncological effect of GH. What is the incidence of tumours in acromegaly?

Shalet: There is a small but significantly increased incidence of a variety of tumours in patients with acromegaly. However, there is no increase in the incidence of leukaemia in acromegalic patients.