Introduction

X.

Xavier

Bertagna

Prof. Xavier Bertagna, Clinique des Maladies Endocriniennes et Métaboliques, Hôpital Cochin, 27, rue du Faubourg-St-Jacques, F-75674 Paris Cedex (France)

Since a young American student discovered the mechanism of proinsulin synthesis by a human islet cell tumor, peptide hormone-secreting tumors have become a field of research where interactions between basic scientists and clinical investigators are the source of considerable progress in endocrinology. We thought the time had come for all actors in this area to share their most recent results during these 32^{es} Journées Internationales Henri-Pierre Klotz d'Endocrinologie Clinique.

The molecular and cellular aspects of peptide hormone production have been thoroughly investigated: their fine mechanisms, and increasing complexity, have been unravelled and the general pattern of the synthesis and secretion which applies to all peptide hormones has been elucidated. Those who will have assimilated this pattern will be offered the means to understand some peculiar clinical features – or lack of clinical features -associated with tumors expressing abnormal molecular forms of a given hormone and to select the most contri-butive tool for the evaluation of their patients. It is now an unavoidable task for endocrinologists, especially in an IRMA era, to become perfectly familiar with all the possible ways through which endocrine tumors can tailor their peptide products.

Perhaps more importantly, clues have recently emerged which shed a new light on the possible mechanisms responsible for the formation of the tumors themselves. They largely evolve from the breakthrough in recombinant DNA technologies which led to the discovery of oncogenes and antioncogenes; some endocrine tumors are associated with in situ gene alterations such as loss of heterozygosity, like in the retinoblastoma. The most fascinating endocrine model in man is probably MEN II (Sipple syndrome) where linkage studies have precisely located the gene of susceptibility to cancer;

already probes are available which, in some informative families, can predict the risk of a medullary thyroid carcinoma, and thus avoid the repeated tests, and anxiety, of the previous screening strategy using calcitonin as a tumor marker. The exciting results in this peculiar endocrine disease even have implications for understanding tumor formation in general; it is a fine example of how close collaboration between basic scientists and clinicians can be fruitful and bear important promises.

A new experimental tool has been developed in trans-genic mice which allows the production of tumors in a given cell type. This technique (site-directed oncogene-sis) creates an 'artificial' model where the steps from endocrine cell hyperplasia to neoplasia mimic those which occur in some 'natural' endocrine diseases as with the thyroid C cells in MEN II. This spectacular approach, which is not restricted to oncogenesis, opens up new avenues in endocrinology; for example, it has permitted to establish cell lines derived from a specific endocrine cell type: it is from such an islet cell tumor line that a CREB (cAMP response element binding) protein was eventually cloned. It was quite intriguing to observe that this ubiquitous intermediate in cAMP

response has sequence homology with other transactivating factors involved in cell proliferation such as the Jun and Fos oncogenes.

Along with these most recent advances in our basic understanding of peptide hormone-secreting tumors, the Journées also focused on more clinically oriented topics. We had chosen to discuss pancreatic islet cell tumors and PTH hypersecretory syndromes.

The ontogeny of the endocrine pancreas and the bio-synthetic pathways for each of its various peptide products have provided a new insight into the general organization of this highly complex endocrine organ. Pan-

8

Introduction

creatic islet cell tumors are characteristically multipotent-secreting tumors. They are a challenge to the physician and their treatment requires a multidisciplinary approach using surgery, pharmacology, chemotherapy, and chemoembolization.

PTH hypersecretory syndromes are among the most common endocrine disorders due to peptide hypersecretion. It was of interest to review the state of the art for their management, the new IRMA and ICMA (Immuno-Chemiluminometric-assay) approaches for PTH measurements, and the consensus, or lack of consensus, on treatment. As a final and most recent example of a successful interaction between clinicians and basic scientists, we were offered the history of the discovery of a

new hormone, the PTH-related protein, a peptide hormone, the existence of which was long suspected on clinical grounds and which was eventually purified and fully characterized in a cell line derived from a human bronchial tumor. It provided the long sought explanation to the many hypercalcemic syndromes associated with malignancies, and already appears as a potential marker of some lung cancers.