Introduction 1

H.		Frisch ^a
S.		Severi ^b

^aDepartment of Paediatrics, University of Vienna, Austria; ^bDepartment of Paediatrics, University of Pavia, Italy

Growth hormone (GH) therapy has been available for over 40 years, but until the development of recombinant-derived GH 8 years ago, supplies were limited and the size and frequency of dose were, therefore, restricted. This symposium provided the opportunity to review the success, and state-of-the-art use of GH therapy for two particular indications – GH deficiency (GHD) and Turner syndrome. Including speakers from Europe, USA and Japan, different experiences of GH therapy were brought together, including studies using GH extracted from the pituitary as well as recombinant-derived GH. The results clearly show that modern standards of GH therapy, using recombinant GH, have brought final height in GHD very close to target height, and that considerable benefit may be derived from GH therapy in Turner syndrome.

Final Height in Growth Hormone-Deficient Patients

Reports on final heights in GHD patients have shown that children treated with the older regimen of intramuscular pituitary-derived GH, 3 times/week, have an unsatisfactory response. In contrast, modern GH therapy using recombinant-derived GH, given as daily subcutaneous injections, has demonstrated a significant improvement in outcome, with final height now approaching target height. However, the optimum time for induction of puberty in patients who are gonadotrophin deficient and for stopping GH therapy still need to be clarified. This will only be possible when larger numbers of GH-treated patients reach their final height.

Final Height in Turner Syndrome

Until recently, it was thought that the benefits of growth hormone (GH) therapy in Turner syndrome and other non-GHD causes of short stature were, at the best, limited. However, the only real measure of the success of GH therapy is that of final height. In the case of Turner syndrome, final height cannot be related to the general population, but must be compared to growth curves determined for the condition. In addition, the interactions between GH therapy and other therapy for these patients must be considered. Dr. Takano, using GH as monotherapy, reported final heights between +1.6 and +2.0 SDS, and pointed out that an increase in dosage would not necessarily increase the final height. Dr. Massa, who added ethinyl oestradiol to the GH treatment in patients older than 12 years, reported final heights of 152.8 cm for GH-treated patients compared with 147.0 cm for untreated patients with Turner syndrome. The response between patients was quite variable, and highlighted the need for an extensive review of patients receiving GH therapy to identify responders from non-responders. In their talk, Drs. Attanasio and Rekers-Mombarg reported better final heights in those patients who had a lower growth velocity before therapy. It was concluded that the optimal time for initiation of growth-promoting therapies, dose of GH and combination with anabolic steroids and oestrogens need to

be defined. Final height may be greatly improved for some patients with Turner syndrome, but the results vary between patients.

© 1995 S. Karger AG, Basel

0301-0163/95/0434-0121

\$8.00/0