Renal Perfusion with Non-Blood Media

L.O.

Simpson

Department of General Practice, University of Otago, Dunedin, New Zealand

Dr. L.O. Simpson, University of Otago, Medical School, PO Box 913, Dunedin (New Zealand)

Dear Sir,

An extraordinary aspect of the paper by Raz et al. [1] was their aim of exploring the effects on renal hemodynamics of perfusion with a non-blood medium. By making the assumption that the physical nature of a physiological solution can be equated with that of blood, the writers create a nonsense scenario and whatever their results imply they relate only to kidneys perfused with physiological solutions. In contrast to the Newtonian nature of physiological solutions, blood is a non-Newtonian fluid with thixotropic characteristics. This means that its viscosity is lowest in vessels where flow is rapid and highest where flow rate is slow. Such characteristics are of importance in renal function.

Glomerular filtration causes haemoconcentration so that the viscosity of post-glomerular blood is higher than that of pre-glomerular blood. In the peritubular plexus reabsorption of water and solutes lowers blood viscosity to pre-glomerular levels. When pre-glomerular blood viscosity is raised, post-glomerular blood viscosity will be hyperviscous. As a result, flow in the peritubular plexus will be impaired and may reduce the absorptive efficiency of tubular epithelial cells to the extent that an abnormal filtrate leaves the nephron. Because of the problems of flow of hyperviscous blood in the peritubular plexus (which in effect is an increase in peripheral resistance) this may stimulate an upstream vasodilation so that capillary blood flow rates may be restored to normal. Should this occur, intraglomerular capillary pressure will increase, filtration fraction will rise and there will be a further increase in post-glomerular blood viscosity.

Therefore, perfusion media which do not undergo the post-glomerular changes in viscosity and rheology typical of blood must be of little relevance to in vivo kidney function. For the same reason, 'haemodynamic' theories of glomerular filtration which do not recognise the haemorheological consequences of filtration must be inadequate to account for the following observations.

Solerte and Fioravanti [2] and Gordge et al. [3] have reported linear relationships between blood viscosity and proteinuria. Furthermore, lowering of blood viscosity with pen-toxifylline resulted in the loss of proteinuria [4]. In polycythemia, blood viscosity is raised commensurately with hematocrit and the associated proteinuria is not due to renal pathology. Successive phlebotomies which lowered blood viscosity also suppressed proteinuria [5].

Such observations indicate that while perfusion studies of isolated kidneys may provide interesting laboratory exercises, their results are of little relevance to in vivo renal function.

References

Raz E, Kaminski N, Brezis M: Effects of perfusion pressure and renal flow upon albumin excretion in isolated perfused kidneys. Nephron 1990;56:396-398.

Solerte SB, Fioravanti M: Hemodynamic alterations in long-term insulin-dependent diabetics with overt nephropathy: role of blood hyperviscosity and plasma protein changes. Clin Nephrol 1987;28:138–143.

Gordge MP, Faint RW, Rylance PB, Neil GH: Abnormal blood rheology in progressive renal failure: a factor in non-immune glomerular injury? Nephrol Dial Transplant 1988;3: 257–262.

Solerte SB, Ferrari E: Diabetic retinal vascular complications and erythrocyte filterability: results of a two-year follow-up study with pentoxifylline. Pharmatherapeutica 1985;4:341–350.