Effect of Erythropoietin Administration on Thyroid Functions of the Patients Undergoing Regular Hemodialysis

N.		Seyrek
S.		Paydas
Y.		Saglikler

Cukurova University Medical Faculty, Department of Nephrology, Adana, Turkey

Prof. Saime Paydas, Associate Professor, Cukurova University, Faculty of Medicine, Department of Internal Medicine, TR-01330 Adana (Turkey)

Dear Sir,

Impaired hypothalamic-hypophyseal-gonadal regulation in uremic patients may induce endocrine disturbances [1]. It has been suggested that administration of erythropoietin (Epo) may improve these endocrinological changes as well as correct anemia in patients with chronic renal failure [2]. In the present study, we investigated the influence of Epo administration on the thyroid function tests in patients undergoing regular hemodialysis (HD).

Fifty-nine clinically euthyroid patients, 31 males and 28 females, with an average age of 38.1 ± 14.8 years (range 15-60) on regular HD, were studied. Patients with goiter, known thyroid disease or with severe systemic illness were excluded. The underlying diseases were glomerulonephritis (n = 17), tubulointerstitial nephritis (n = 11), hypertension (n = 23), autosomal dominant poly-cystic kidney disease (n = 3) and unknown (n = 5). Patients were dialyzed for 4 h 2-3 times/week, receiving subcutaneous Epo 150 IU/kg/week (for 6 months) and oral ferrous preparations. Serum hematocrit (Hct), total protein, albumin, triiodothyronine (T₃), thyroxine (T₄) and thyroid-stimulating hormone (TSH) were taken immediately before HD. Hematocrit, total protein and albumin were measured by autoanalyzer, T₃ and T₄ were analyzed by using radioimmunoassay, and TSH was analyzed by ELISA. Results of the measurements of thyroid function were compared between patients receiving and patients not receiving Epo (controls).

T₃ and T₄ were below the normal range in 69 and 5% of the study group, respectively. TSH was above the normal range in 6.7% of the study group. Of the patients who were receiving Epo, T₃ was low in 21 (70%), T₄ was low in 1 (3%) and TSH was elevated in 2 (6.6%). On the other hand, T₃ was low in 20 (69%), T₄ was low in 2 (6.8%) and TSH was elevated in 2 (6.8%) of the patients who were not receiving Epo.

Epo therapy-associated well-being of patients on HD has been due not only to the correction of the anemia, but also to an associated correction of endocrine abnormalities. Lundin et al. [3] have reported that an Epo-associated increase of hair thickness

Student's t test and Pearson's correlation were used for statistical analysis.

Of the 59 patients, the mean Hct, total protein, albumin, total T₃, total T₄ and TSH were $19.7 \pm 4.1\%$, 7.9 ± 1.1 g/dl, 4.3 ± 0.9 g/dl, 53.1 ± 25.5 ng/dl, 6.1 ± 1.6 µg/dl and 1.8 ± 1.6 mIU/ml, respectively. The results of the patients in respect to receiving Epo are summarized at table 1.

Serum T₃, T₄ and TSH were similar in the two groups and there was no statistical difference between the two groups. No significant correlation was demonstrated between Hct and T₃, T₄, TSH in patients who were receiving Epo. But a correlation was found between albumin and T₃ in this group ($r = 0.3863$, $p < 0.05$).

Table 1. Characteristics of the patients

KARGER

E-Mail karger@karger.ch Fax + 41 61 306 12 34

© 1996 S. Karger AG, Basel 0028-2766/96/0724-0714\$ 10.00/0

and change of quality of fingernails might be an influence of Epo on thyroid function. Although decreased serum levels of T₃ and T₄ have been observed in HD patients, no difference has been found in respect to receiving Epo administration or not in HD patients in our study [1]. So, Epo did not seem to improve these abnormalities despite a partial correction of the anemia.

In our study, we found that T₃ and T₄ were below the normal range in 69 and 5 % of the patients on regular HD, respectively. The decrease in serum level of T₃ was more significant.

Extrathyroidal reduction in conversion of T₄ to T₃ in uremia may explain this situation. The serum levels of TSH were within normal limits except in 3 patients in our study group. There was only a correlation between albumin and T₃ in patients who were receiving Epo.

In conclusion, our data confirm that he-modialized patients often have asymptomatic thyroid disturbances. We also suggest that Epo does not improve these abnormalities, despite it correcting anemia in patients undergoing regular HD. Finally, the influences of uremia and HD on thyroid hormones have to be studied regularly and if it is required, thyroid hormone preparations have to be given.

References

- 1 Lim VS, Fang VS, Katz AI, Refetoff S: Thyroid dysfunction in chronic renal failure: A study of the pituitary-thyroid axis and peripheral turnover kinetics of thyroxine and triiodothyronine. *J Clin Invest* 1977;60:522-534.
- Gaffin E, Oliveira DBG, Raggatt P, Evans DB: Assessment of the thyroid function of patients undergoing regular hemodialysis. *Nephron* 1993;65:568-572.
- Lundin AP: Quality of life: Subjective and objective improvements with erythropoietin therapy. *Semin Nephrol* 1989;9(suppl 1):22-29.
- Effect of Epo Administration on Thyroid Functions of the Patients Undergoing Regular Hemodialysis *Nephron* 1996;72:714-715 715