Introduction

C.

Grassi

Department of Pneumology, University of Pavia, Istituto Forlanini, Pavia, Italy

Prof. C. Grassi, Istituto Forlanini, I-27100 Pavia (Italy)

The recent explosion of basic knowledge that has resulted in a revolution in lung biology and pulmonary medicine is fundamental for the understanding of the pathophysiology of chronic bronchitis and emphysema. This meeting is devoted mainly to selected topics concerning the pathophysiology of emphysema. Some of the basic biochemical mechanisms of emphysema are also involved in the induction of cancer, and this meeting will also touch this difficult and intriguing area.

We now have evidence that cigarette smoking is the principal identifiable cause of emphysema in humans. We do not want to repeat here the catastrophic figures related to the smoking habit; a lot of publications are available on this subject and it is supposed that anyone who is deeply involved in 'respirology' will know them. The latest comprehensive report has been published by E. Freeding in the New England Journal of Medicine, no. 8, August 1985.

I am confident that this meeting will show us how the greatest achievements of the last few years have enabled us to move the basic research in lung injury into the clinical domain, which may, potentially, lead us to new frontiers of prophylaxis and therapy of chronic bronchitis and emphysema.

The post-mortem pathological evidence of emphysema was the only diagnosis possible only a few years ago. Now it has been shown that the diagnosis of emphysema is possible in life, in the early stage of the disease, i.e., at the beginning of those biochemical and cellular events which represent the basis of the progress we have witnessed over the last two decades, and which will open the doors to new therapeutic approaches.

Judging from the reports published by the specialized and unspecialized medical journals which deal with biological problems related to chronic bronchitis and emphysema, it seems widely accepted that the disease is due to the destruction of the elastic fibres of the alveolar structures. This destruction is the consequence of an imbalance between elastolytic and anti-elastolytic activities in the lung. This imbalance is mainly related to the oxidative inactivation of the antielastolytic system, which in turn results from the imbalance in another system, between oxidants and antioxidants.

Grassi

Cigarette smoking seems to be the principal identifiable cause of lung injury through this mechanism. In fact, it increases the oxidant burden in the lungs. All of us know that

the oxidant burden is represented mainly by oxygen-centered reactive species, also referred to as free radicals, which seem to oxidize both a **me**-thionine residue essential to the activity of antiproteases, and reduce glutathione, which could lead to the inactivation of thiol-dependent enzymes that are among the major antioxidant tissue defences in vivo.

Once formed, the oxygen-reactive species can enter into a wide variety of toxic reactions. Aggressive oxygen species react, among other things, with:

DNA, by initiating and promoting **mu**-tagenesis and carcinogenesis,

proteins, by depressing the enzymatic defense mechanism,

lipids, by disrupting membrane integrity, resulting in cell damage or death.

The prevention of oxidative cell damage by antioxidants should be regarded as an obvious approach not only in emphysema, but also in drug toxicity and other pathophysiological conditions including those related to cancer induction. In this context, N-acetyl**c**ysteine should be seen as a fundamental drug. This simple thiol-containing aminoacid is a precursor of glutathione and has been shown to be a potent antioxidant and free-radical scavenger in both biological and non-biological free-radical-generating systems. This drug, when given orally in a dosage of 600**mg**/day, has proved to protect lungs from the damage of air pollution and cigarette smoking.

This meeting will extensively take into consideration all the latest scientific data and reasons which prospectively point to N-acetylcysteine for the antioxidative management of emphysema, while waiting for more specific treatments for this disease.