Introduction

New Perspective for Broncho-alveolar Lavage

C.F.		Donner
P.		Howard

Dr. C.F. Donner, Divisione di Pneumologia, Centro Medico di Riabilitazione, I-28010 Veruno, Italy

After a decade of great interest in bronchoalveolar lavage (BAL) as a potential tool in the clinical evaluation of interstitial lung diseases, enthusiasm has faded because of poor specificity of the method and a number of difficulties in technique standardisation [1].

As a therapeutic tool performed with large volumes of saline, BAL is rarely applied in conditions such as alveolar proteinosis, cystic fibrosis and removal of massively inhaled foreign particles. In diffuse diseases, BAL can be considered of diagnostic value only in unusual conditions, such as alveolar proteinosis, histiocytosis X and berylliosis. In environmental lung diseases BAL is useful in detecting or assessing exposure to specific agents [2]. In a clinical setting, BAL fluid cell analysis in non-immunocompromised patients with interstitial lung disease showed a clear contribution to a precise diagnosis in an important percentage of patients [3].

BAL has proven to be, and still remains, an invaluable research tool, which has widened the field of interest of pneumology into the field of molecular cell biology. It is mainly due to BAL if lung specialists are becoming increasingly confident with topics such as cell receptors, cytokines, second messengers and genes. Such aspects were the subject of the 1st International Workshop 'Bronchoalveolar lavage: new insights in research and clinical applications' held in Veruno, Italy, in April 1989 [4].

Recently BAL has regained prominence in the clinical setting. Clinicians now recognise BAL as a useful means of diagnosis of infections in critically immunocompromised patients, and of atypical pneumonia in the normal host. In conditions of suspected primary or metastatic lung cancer BAL can give definitive information, particularly when invasive approaches cannot be used and standard procedures have failed to demonstrate neoplastic tissue. New applications continue to be found for bronchial diseases such as asthma and chronic obstructive pulmonary disease. Since BAL can be safely performed in patients with mild stable airway obstruction, lavage of large airways has become a widely used method of study of asthma, chronic bronchitis and emphysema [5]. This requires a clear understanding of the distinction between BAL and bronchial lavage (BL). The first recovery is predominantly bronchial in origin and should be examined separately from the subsequent washes. For more selective evaluation the use of balloon-tipped or double-lumen catheters to wash specific bronchi are under evaluation.

Assessment of inflammation in bronchial airways and correlation with the clinical and functional data could be of great value for a better understanding of pathogenesis and therapy of the disease. BAL/BL before and after pharmacological therapy may afford important clues to drug efficacy and mechanism of action. Recent trials with alpha-1-antiprotease substitution treatment in PiZZ emphysematous patients, and drug induced variations in cellular and mediator profile after localised specific bronchial challenge in asthma are examples.

This workshop considered these new applications of BAL and BL. Three sessions devoted respectively to

asthma, chronic obstructive pulmonary disease and lung cancer and preneoplastic lesions, all focused on some fundamental issues, namely (1) a comparison between the classical means of investigation, such as histopathology, and BAL/BL in both clinical and purely investigative fields; (2) the pathogenic relevance of the biochemical and cellular data from BL and BAL fluid analysis, and (3) therapeutic implications (e.g. bronchoalveolar modifications after antituberculous therapy and BAL in the evaluation of substitution treatment of alpha-1-antitrypsin deficient destructive lung diseases).

This symposium report contains the important contributions presented at the meeting, which it is hoped will stimulate much future research into BAL and BL. At the same time we are more confident that expectations from airway lavage can be precisely defined, contributing to a clearer appreciation of the relative merits of this technique amongst those available to modern pneumologists.

References

Klech H, Pohl W (eds): Technical recommendations and guidelines for bronchoalveolar lavage (BAL). *Eur Respir J* 1989;2:561-585. Helmers RA, Hunninghake GW: Bronchoalveolar lavage in the non-immunocompromised patient. *Chest* 1989;96:1184-1190. Stoller JK, Rankin JA, Reynolds HY: The impact of bronchoalveolar lavage cell analysis on clinicians' diagnostic reasoning about interstitial lung disease. *Chest* 1987;92(5):839-843.

Workshop Report: Workshop on Bronchoalveolar Lavage: new insights in research and clinical applications. *Eur Respir J* 1990;3: 354-378.

Reynolds HY: Bronchoalveolar lavage. *Am Rev Respir Dis* 1987;135:250-263.

Donner/Howard

Introduction