Chairman's Summary

J.

Dent

Gastroenterology Unit, Royal Adelaide Hospital, Adelaide, Australia

J. Dent, Gastroenterology Unit, Royal Adelaide Hospital, North Terrace, Adelaide SA 5000 (Australia)

The presentations in these Proceedings speak for themselves. They illustrate a remarkable development in the therapeutic uses of acid inhibition, which would have been difficult to predict only a few years ago.

The success of omeprazole in the treatment of reflux oesophagitis, and its potential in the management of non-steroidal anti-inflammatory drug-induced ulcer and acute gastrointestinal bleeding are likely to be dependent upon the marked reduction of aggressiveness of gastric juice. This reduction is, on the whole, probably due to the relative inactivity of pepsin at the gastric pH seen during omeprazole therapy.

The effects of gastric acid inhibition with omeprazole given in combination with oral amoxycillin has opened up a new area of therapy for the management of *Helicobacter pylori*-associated disease, particularly duodenal ulcer. It appears inevitable that the results on the eradication of *H. pylori*, as summarized in these Proceedings, will lead to a major shift in strategies for the management of chronic duodenal ulcer.

The views of the Faculty on patient monitoring during long-term omeprazole therapy, with regard to the issue of safety, illustrate the significant advances that have been made in defining the effects of long-term inhibition of gastric acid secretion with omeprazole. Endo-scopy, biopsy and measurement of serum gastrin levels are being performed on a regular basis in patients treated with long-term omeprazole, who have not been included in clinical trials or surveillance programmes, in the belief that they would provide useful information about the safety of omeprazole in individual patients.

Examination of the logic behind these practices before and during the symposium led the Faculty to support overwhelmingly the view that no formal screening needs to be performed within 5 years of starting continuous omeprazole therapy. In addition, it was concluded that if serum gastrin levels were measured and gastric biopsy was performed in an individual patient, the information obtained could not be interpreted usefully for that patient. Obviously, this does not mean that structured surveillance programmes are not needed – the success of long-term inhibition of acid secretion is such that it will be used increasingly frequently, so it is important that its long-term safety is examined critically.

To date, the most important side-effect of omeprazole is a very happy one. Questions regarding the long-term safety of gastric acid inhibition have provided an enormous stimulus to improve understanding of the biology and structure of the gastric mucosa. Some of this increasing knowledge has been highly relevant to topics covered in these Proceedings. I hope this stimulus will continue for a long time to come.

Karger AG. Basel

0012-2823/92/ 0517-0122 \$ 2.75/0