Foreword

R.C.

Robert C.

Heading,

Edinburgh, UK

Early in the 14th century, William of Occam promulgated the principle, since known as Occam's razor, which states: 'Entities are not to be multiplied beyond necessity'. In fact, it was not William of Occam who originated this idea but the Frenchman Durand de Saint-Pourçain, who advised his fellow theologians, scholars and philosophers that their theories and arguments should not be more complicated than the facts required. However, it is Occam's name that remained attached to the concept. Now, some 650 years on, the same idea has been expressed by the inelegant acronym KISS ('Keep it simple, stupid!') but the concept seems to have a fundamental validity which has enabled it to stand the test of time.

Applied to the scientific principles of observation, hypothesis and experimentation in the biological domain, Occam's razor leads to the sequence of iterative review and refinement of our ideas of biological function, necessitated by the fact that our hypotheses must be the simplest (approximately) which fit the facts. Experimentation generates new facts, which are then reconciled with a revised hypothesis. Nevertheless, as every clinical scientist and physician (and politician and journalist) will recognize, interpretation of the facts' may be critically determined by their order of presentation and the weighting given to each.

The integrity of the gastroduodenal mucosa has been viewed for decades as being preserved only when the balance between aggressive and defensive factors favours the latter. Many a physiology lecturer possesses that familiar

slide showing a schematic see-saw, with aggressive factors for mucosal disruption depicted at one end and defensive ones at the other. No similar concept has been necessary in respect of the oesophageal mucosa, because, in contrast with the stomach and duodenum, there was no experimental data to challenge the existence of an absolute and direct association between exposure to erosive acid + pepsin, and the occurrence of mucosal damage. With hindsight, this may seem naive but the application of Occam's razor permitted the simple hypothesis, even though it might be inconsistent with biological common sense.

Experimental observations have now led us to recognize that there is a considerable complexity to relationships between damaging influences of the luminal content and the occurrence of oesophageal mucosal damage. The factors which together resist the development of injury are very properly described as the processes of oesophageal mucosal defence. Explanation of these processes and careful evaluations of their respective contributions are timely and, as collated in this supplement, are extremely welcome.

The high scientific reputation of the authors of these papers will be well known to most oesophagologists, but I am confident that, like myself, readers will recognize that is not just the individual contributions that are admirable, but the way in which they together build a comprehensive picture of the way in which the oesophagus protects itself and reacts to

potential injury. Some novel experimental techniques have been applied to the subject, and await

1

adoption by other laboratories where ideas will no doubt be further advanced.

Most living tissues seem to have mechanisms which protect them against injury, react to minimize damage, and institute repair if damage occurs. The oesophagus is no exception and we must do our best to understand these mechanisms if we wish to claim knowledge of how the inflammation, ulceration and even metaplastic change in the oesophageal squamous epithelium occurs. These are important matters.

Professors Scarpignato, Galmiche and Giulì are to be congratulated on assembling a distinguished group of investigators whose individual contributions to this supplement are all excellent, and who, together, provide a body of expertise and knowledge on this subject which brings the reader fully up to date. New experimental observations made during future years will no doubt cause theories to be modified – Occam's razor still applies – but we now belatedly recognize that oesophageal muco-sal defence, like gastroduodenal mucosal defence, is important to mucosal integrity. It is easy with hindsight to express incredulity at our failure to recognise this previously.

My congratulations to the authors and editors. I have thoroughly enjoyed reading this supplement.

2

Heading

Foreword