Lipid Peroxidation in Iron Deficiency Anemia

Ş. Sinasi Özsoylu

Department of Pediatrics, Hematology Unit, Faculty of Medicine, Hacettepe University and Hacettepe Children's Hospital, Ankara, Turkey

Prof, Şinasi Özsoylu, MD, Department of Paediatrics, Faculty of Medicine, Hacettepe University, Hacettepe Children's Hospital. TR-06100 Ankara (Turkey)

I have read with interest the paper by Bartal et al. [1] in a recent issue of the Journal. The authors stated that 'lipid peroxidation was much higher in iron-deficient cells than in control cells'. However, the malonyl-dialdehyde (MDA) levels were given as na-nomoles per gram hemoglobin (Hb). In iron deficiency anemia, more cells are required for 1 g Hb than in normal cells. MDA is not related to Hb but to the cell membrane. Therefore, MDA levels should be expressed per red blood cell rather than per gram Hb.

When I determined catalase activity in iron deficiency anemia [2], which was the first demonstration of an iron-containing enzyme deficiency, it could be shown when expressed per red cell but not with Hb level, as later stated by Acharya et al. [3].

It is surprising to see that some studies are not appreciated in the literature despite priority and several reinforcements [2, 4, 5].

References

Bartal M, Mazor D, Dvilansky A, Meyerstein N: Iron deficiency anemia: Recovery from in vitro oxidative stress. *Acta Haematol* 1993;90: 94-98.

Özsoylu Ş: Decreased activity of iron-containing enzyme catalase in human iron deficiency anemia. *Turk J Pediatr* 1964;6:200-210. Acharya J, Punchard NA, Taylor JA, Thompson RPH, Pearson TC: Red cell lipid peroxidation and antioxidant enzymes in iron deficiency. *Eur J Haematol* 1991;47:287-291. Özsoylu Ş: Detection of iron containing enzyme deficiency anemia. *J Pediatr* 1973;82:168. Özsoylu Ş: Note to the paper. *Acta Haematol* 1986;76:181.

Acta Haematol 1994;91:170

M. Bartal, D. Mazor

A. Dvilansky, N. Meyerstein

Department of Physiology, Faculty of Health Sciences, Ben-Gurion University of the Negev, Beer-Sheva, Israel

Reply

We do agree that malonyldialdehyde (MDA) reflects lipid peroxidation and is not related to hemoglobin (Hb). In their detailed paper about auto-oxidation of human red blood cell lipids, Stocks and Dormandy [1] outlined the TBA reaction and calculated MDA in nanomoles per gram Hb. Later, Acharya et al. [2] emphasized that iron deficiency anemia cells were more sensitive to

oxidation when MDA levels were expressed per gram Hb. Our paper supports these findings. Unfortunately, we could not quote Prof. Özsoylu's paper in the Turkish Journal of Pediatrics (1964), as this journal is not available in our library.

References

Stocks J, Dormandy TL: The autoxidation of human red cell lipids induced by hydrogen peroxidase. *Br J Haematol* 1971;20:95.

Acharya J, Panchard NA, Taylor JA, Thompson RPH, Pearson TC: Red cell lipid peroxidation and antioxidant enzymes in iron deficiency. *Eur J Haematol* 1991;47:287.

Accepted:	Dr. M. Bartal	©1994 S. Karger A
April	Department of Physiology, Faculty of 0001-	
	Ben-Gurion University of the Negev	% 8.00/0
	POB 653	