Correspondence

Accepted: October 18, 1984

Acta haemat. 1985;73:61

Unusual Features of Brucellosis Complicating Beta Thalassaemia major

M.A.		Babiker
A.M.		Kambal
E.F.		Ashong

Riyadh

Dr. M.A. Babiker, A.M. Kambal, E.F. Ashong, Department of Paediatrics, College of Medicine, PO Box 2925, Riyadh (Saudi Arabia)

Hypersplenism complicating β -thalassaemia major is usually treated surgically. We are reporting hypersplenism with other rare features of brucellosis in a thalassaemic child that resolved with medical treatment. The child is a 5-year-old boy who needed a fresh unit of washed packed cells every 4–5 weeks to keep his haemoglobin above 9 g/dl. 6 months ago he presented with pallor and backache 15 days after the last transfusion. His spleen was 3 cm bigger than before. The patient had maculopapular skin rash and a small tender cystic mass in his scalp. The laboratory tests revealed pancytopenia (table I). Blood culture, Coombs' test and titres for toxoplasma, cytomegalo-virus, and coxsackievirus were negative. Tuberculin and monospot tests were negative. The bone marrow was hypercellular. Table I shows the haematological data obtained and the treatment given at each 2-week-ly follow-up examination. At the second visit the symptoms and signs persisted, and there was no change in the viral titres or Coombs' tests. At the third visit the brucella abortus antibody titre was 1:5,280. He was started on cotrimoxazole 40 mg 12 hourly. At the fourth visit he had no backache, but the other

Table I. Haematological data and treatment

Visit Hb RBC WBC Platelets Treatment No. g/dl 1012/1 10V1 10V1

signs persisted. At the fifth visit the haematological data improved, but the scalp swelling increased, and the skin rash persisted. Rifampicin 20mg/kg/day was added. After 4 weeks the rash and swelling disappeared, and the pancytopenia resolved. The brucella titre declined to 1:640. Over the next 3 months he remained well. He is currently back on his 4- to 5-week transfusion regimen.

Comment

The diagnosis of hypersplenism was based on clinical observations and routine tests. We could not do sequestration studies. Hypersplenism was reported in very few cases of brucellosis [1]. Skin rash was seen in less than 5% of cases [2]. Soft-tissue abscesses occurred very rarely in abattoir workers [3]. In conclusion, a medically treatable cause of hypersplenism is worth looking for before surgery is considered. Brucellosis should be looked for where it is endemic.

References

Lynch, E.C.; McKechnie, J.C.; Alfrey, C.P.: Brucellosis with pancytopenia. Ann. intern. Med. 69: 319–322 (1968).

Spink, W.W.: The nature of brucellosis, pp. 45–190 (University of Minnesota Press, Minneapolis 1956).

Buchanan, T.M.; Faber, L.C.; Feldman, R.A.: Brucellosis in the United States, 1960–1972: an abattoir-associated disease. I. Clinical features and therapy. *Medicine* 53: 403–413 (1974).