Preface

After a transient ischemic attack or minor stroke the early risk of recurrence is high and ranges up to 10–15% within the following weeks. Currently fluctuating clinical signs and symptoms, high prognostic scores (ABCD, ABCD² and ABCD-I) and very early signs of ischemia in diffusion-weighted MRI studies are poor prognostic markers [1, 2]. Although the long-term risk, both of coronary and cerebral events, is much lower with approximately 2% per year, stroke subtypes associated with large- and small-vessel disease are particularly prone to subsequent focal and global deterioration such as vascular dementia.

At present, treatment of risk factors such as hypertension, diabetes mellitus, lipid or homocysteine metabolic disorders as well as modification of lifestyle are of utmost importance to reduce both early and long-term risks of vascular events, dementia and death [3]. Antithrombotic drugs, such as aspirin, clopidogrel and aspirin + extended-release dipyridamole decrease the relative risk of cerebral and myocardial infarction as well as vascular death in patients with previous events by 15–25% at best, failing to do so in 75–90% of all patients [4]. Furthermore, stroke reduction seems to be different in men and women, cardio- and cerebrovascular ischemia, patients with different combinations of risk factors and for various subtypes of ischemic stroke: again patients with large- and small-vessel disease of either extra- or intracranial manifestation seem to benefit more from antiplatelet agents than patients with cardiogenic, undetermined or rare conditions of stroke.

Evidence-based recommendations for antithrombotic drug treatment result from former large randomized clinical trials using the aforementioned agents in rather

similar designs: however, very early high risk of recurrence, in particular after transient and fluctuating clinical signs/symptoms, had not been addressed and classification of stroke subtypes was mainly based on the old, sometimes modified TOAST classification [5]. More recently designed classifications are based on a wider spectrum and combinations of stroke phenotypes and have introduced a level of diagnostic evidence to recognize completeness, quality and timing of evaluation [6]. In addition, they include more sophisticated diagnostics, which are currently available and widely used today: (a) vascular ultrasound and echocardiography, (b) diffusion-weighted and T₂*-weighted MRI studies for the detection of early focal ischemia or bleedings, (c) ultrasound, MR and CT perfusion and angiography studies and (d) specific laboratory tests. They also address hemorrhagic aspects, which may illustrate potential side effects better, such as primary and secondary hemorrhages, or late ischemic subsequences, like inflammation, apoptosis, reorganization and recovery.

The articles presented in this supplement issue to *Cerebrovascular Diseases* surround a new ongoing, large randomized clinical trial comparing the efficacy and safety of terutroban versus aspirin in secondary prevention of cerebrovascular and cardiovascular events of ischemic origin in patients after transient ischemic attacks or stroke (PERFORM Study) [7]. Terutroban, a TP receptor antagonist, has – like other currently used drugs – considerable antithrombotic effects, but also antivasoconstrictive, antiatherosclerotic and anti-inflammatory effects, which may increase the armamentarium against progression of large- and small-vessel atherosclerosis in the hope of reducing secondary events, including demen-

tia [8]. This ongoing study is unique with regard to the extensive study of many of these aspects so far experimentally investigated only, and hence truly forms a modern translational randomized clinical trial. A series of specific metabolic parameters potentially active at different levels of atherogenesis are determined in most patients recruited. Furthermore, a refined subtype classification adapted to current technologies of investigation has been specially introduced and, finally, large ancillary

studies to identify vascular and brain tissue changes over time using best available ultrasound and MR technologies are accompanying the main study. This promising, up-to-date randomized clinical trial rather soon will hopefully answer many questions still open today with regard to *if, why, how, and in whom* best prevention of recurrent stroke, other vascular events and dementia can be achieved.

M.G. Hennerici

References

- 1 Pendlebury ST, Rothwell PM: Risk of recurrent stroke, other vascular events and dementia after transient ischaemic attack and stroke. *Cerebrovasc Dis* 2009;27(suppl 3): 1–11.
- 2 Chatzikonstantinou A, Willmann O, Jäger T, Szabo K, Hennerici MG: Transient ischemic attack patients with fluctuations are at highest risk for early stroke. *Cerebrovasc Dis* 2009;27:594–598.
- 3 European Stroke Organisation (ESO) Executive Committee; ESO Writing Committee Collaborators. Guidelines for management of ischaemic stroke and transient ischaemic attack 2008. *Cerebrovasc Dis* 2008;25:457–507.
- 4 Boussier MG: Antithrombotic agents in the prevention of ischemic stroke. *Cerebrovasc Dis* 2009;27(suppl 3):12–19.
- 5 Adams HP, Bendixen BH, Kapelle LJ, Biller J, Love BB, Gordon DL, Marsh EE: Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. *Stroke* 1993;24:35–41.
- 6 Amarenco P, Bogousslavsky J, Caplan LR, Donnan GA, Hennerici MG: Classification of stroke subtypes. *Cerebrovasc Dis* 2009; 27:493–501.
- 7 Hennerici MG, on behalf of the PERFORM Study Investigators: Rationale and design of the Prevention of Cerebrovascular and Cardiovascular Events of Ischemic Origin with Terutroban in Patients with a History of Ischemic Stroke or Transient Ischemic Attack (PERFORM) Study. *Cerebrovasc Dis* 2009;27(suppl 3):28–32.
- 8 Chamorro A: TP receptor antagonist: a new concept in atherothrombosis and stroke prevention. *Cerebrovasc Dis* 2009;27(suppl 3):20–27.