Summary and Conclusions

J.		Hirsh
----	--	-------

Hamilton, Ont., Canada

Dr. J. Hirsh, Hamilton Civic Hospitals Research Centre, 711 Concession Street, Hamilton, Ont. L8V 1C3 (Canada)

Orgaran is a low-molecular-weight (LMW) heparinoid composed of heparan sulfate (80%), dermatan and chondroitin sulfate. It has a mean molecular weight of approximately 6,000 U and an anti-factor Xa/anti-factor IIa ratio of greater than 20/1. Compared to standard heparin, it has minimal inhibitory activity on platelet function and produces much less bleeding in experimental bleeding models for equivalent anticoagulant effects. Most of the anti-factor Xa activity is accounted for by a high affinity fraction of heparan sulfate, and the relatively modest anti-factor IIa activity is provided by derma-tan sulfate. In experimental models, the anti-thrombotic activity of Orgaran is more persistent, and the hemorrhagic effects are less persistent than standard heparin; findings which suggest that Orgaran could have therapeutic advantages over standard heparin as an anti-thrombotic agent. Interestingly, not all of the antithrombotic activity of Orgaran is accounted for by its high affinity fraction.

The clearance of Orgaran is much slower than heparin and the LMW heparins. The half-life of anti-factor Xa activity is 25 h and that of anti-factor IIa activity 4.3 h. The bio-availability after subcutaneous injection is excellent, close to 100%, although absorption is relatively slow.

Steady-state anti-factor Xa levels are achieved with twice daily subcutaneous injections. A large proportion of the heparinoid is excreted in the urine.

Orgaran is the glycosaminoglycan of choice for treatment of patients with heparin-induced thrombocytopenia (HIT). It has minimal cross-reactivity in tests for heparin-associated antibody with plasma from patients with HIT (12% compared to 80% for LMW heparins. Orgaran has been used to treat over 160 patients with presumed HIT with good clinical results.

Clinically, Orgaran has been shown to be effective in the prevention of venous thrombosis in patients having elective hip surgery (where it is more effective than the combination of standard heparin/dihydroergotamine), surgery after hip fracture (where it is more effective than warfarin and dextran) and in patients with stroke. The risk reduction compared to untreated controls is very impressive and in the order of 70%. Results of a pilot study suggest that Orgaran can be used safely in selected patients with thrombotic stroke.

Thus, Orgaran represents a real and important addition to contemporary antithrombotic therapy.