findings were acquired on 1.5-T scanners and that this entails some limitations. High b-value DWI requires a hardware upgrade because most scanners cannot generate a 40-mT/m gradient field [6]. Despite the lower MR signal intensity and lower signal-to-noise ratio on high b-value images [8], 3-T DWI is at least as good as, and in some ways superior to, 1.5-T DWI for assessing hyperacute stroke [9].

We postulate that one reason for missed acute ischemic stroke might be the conspicuity of acute ischemic lesions on diffusion-weighted images, which is related to the b-value. In some cases (as in this one) higher b-values might offer an opportunity for earlier lesion detection. We found that DWI_{4,000} performed on the 3-T scanner was superior for the identification of ischemic lesions. Our findings suggest that the PWI-DWI mismatch time window is shorter than was previously thought. Studies are underway in our laboratory to determine the optimal b value for the assessment of acute stroke. Based on our findings, we suggest that high b-value DWI is useful for the selection of thrombolytic therapy in patients with acute stroke.

References

- 1 Kane I, Sandercock P, Wardlaw J: Magnetic resonance perfusion diffusion mismatch and thrombolysis in acute ischemic stroke: a systematic review of the evidence to date. *J Neurol Neurosurg Psychiatry* 2007;78: 485–491.
- 2 Mintorovitch J, Moseley M, Chileuitt L, et al: Comparison of diffusion and T₂-weighted MRI for the early detection of cerebral ischemia and reperfusion in rats. *Magn Reson Med* 1991;18:39–50.
- 3 Gonzalez R, Schaefer P, Buonanno F, et al: Diffusion-weighted MR imaging: diagnostic accuracy in patients imaged within 6 hours of stroke symptom onset. *Radiology* 1999;210:155–162.
- 4 Saur D, Kuchinski T, Grzyska U, et al: Sensitivity and interrater agreement of CT and diffusion-weighted MR imaging in hyperacute stroke. *AJNR Am J Neuroradiol* 2003;24:878–885.
- 5 Lam WW, Wong KS, Rainer TH, et al: Assessment of hyperacute stroke like symptoms by diffusion-weighted images. *Clin Imaging* 2005;29: 6–9.
- 6 Meyer J, Gutierrez A, Mock B, et al: High b-value diffusion-weighted MR imaging of suspected brain infarction. *AJNR Am J Neuroradiol* 2000;21:1821–1829.
- 7 Burdette J, Elster A: Diffusion-weighted imaging of cerebral infarctions: are higher b-values better? *J Comput Assist Tomogr* 2002;26: 622–627.
- 8 Cihangiroglu M, Uluğ AM, Firat Z, et al: High b-value diffusion-weighted MR imaging of normal brain at 3 T. *Eur J Radiol* 2007.
- 9 Kosior RK, Wright CJ, Kosior JC: 3-Tesla versus 1.5-Tesla magnetic resonance diffusion and perfusion imaging in hyperacute ischemic stroke. *Cerebrovasc Dis* 2007;24:361–368.

Kaoru Kurisu, MD, PhD
Department of Neurosurgery
Graduate School of Biomedical Sciences
Hiroshima University
1-2-3 Kasumi, Minami-ku, Hiroshima, 734-8551 (Japan)
Tel. +81 82 257 5225, Fax +81 82 257 5229
E-Mail kuka422@hiroshima-u.ac.jp

Cerebrovasc Dis 2009;27:617–618

DOI: [10.1159/000218074](https://doi.org/10.1159/000218074)

Large Carotid Thrombus

M.G. Delgado^a, P. Vega^b, V. Mateos^a, J. Bogousslavsky^c

Departments of ^aNeurology and ^bRadiology, Hospital Central de Asturias, Oviedo, Spain; ^cDepartment of Neurology, Genolier Swiss Medical Network, Valmont-Genolier, Glion-sur-Montreux, Switzerland

A 76-year-old female with high blood pressure and atrial fibrillation as cerebrovascular risk factors was admitted because of sudden left hemiparesis and disarthria with a right periorbital headache. Transcranial Doppler ultrasound showed increased velocity in the right carotid siphon. Her cranial CT was normal. She was anticoagulated because of carotid siphon stenosis with low-molecular-weight heparin (0.6 ml subcutaneous nadroparin calcium every 12 h). An angio-CT showed calcified plaque in the right carotid siphon with a large thrombus (fig. 1A, B), resulting in a high-grade stenosis. We suggested the thrombus formation was due to a cardiac embolus stopped in the right

Fig. 1. Intra- and extracranial angio-CT. Coronal and lateral projections showed a large thrombus in the right internal carotid artery (arrows, red in the online version).

Color version available online

siphon stenosis and a retrograde enlargement mechanism. The patient was under anticoagulation. An angioCT at 10 days showed complete thrombus resolution. At discharge, she had slight left ataxic hemiparesis (NIHSS = 4). At 1 year she had an mRS = 1.

This case indicates that large thrombi may be dissolved by anticoagulant therapy with low-molecular-weight heparin, although its lytic activity has never been demonstrated. The short recanalization time was probably due to an acute unorganized thrombus.

References

- 1 Hirsh J, Levine MN: Low-molecular-weight heparin. *Blood* 1992;79: 1–17.
- 2 Sandercock PA, Counsell C, Kamal AK: Anticoagulants for acute ischaemic stroke. *Cochrane Database Syst Rev* 2008;4:CD000024.

Montserrat González Delgado
Department of Neurology
Celestino Villamil s/n.
ES-33006 Oviedo (Spain)
Tel. +34 985 10 80 00, ext. 36 206, E-Mail mglezdelgado@yahoo.es