In this edition of *Ophthalmologica*, we have many interesting papers crossing multiple disciplines in ophthalmology with important clinical signs and information to help us take care of complicated patients.

Kim et al. [Ophthalmologica 2009;223:222–227] looked at a vexing problem secondary to the increasingly common practice of combining phacoemulsification with posterior chamber intraocular lens insertion at the same time as a pars plana vitrectomy. This combination is often more efficient for the patient when there is a cataract present, and helps visualization for the retinal surgeon. While the results have been overwhelmingly excellent, there has been a surprising incidence of iris posterior synechia formation to the intraocular lens, which has been reported to be in the range of 7–30% of cases. Exactly why this is more common than without vitrectomy has been a bit of a mystery.

Kim et al. looked at 153 eyes of 153 patients treated by a combination of phacoemulsification and vitrectomy, and then performed a multivariate analysis of a host of possible causative factors. The most important risk factor was anterior chamber fibrin deposition on the first postoperative day. Clearly, the combination of both procedures must result in an increased liberation of inflammatory cytokines and, if you see fibrin, the patient is 6 times more likely to develop synechiae. Clearly, we need to do a better job of trying to block this level of postoperative inflammation.

Interestingly, they also found that if C3F8 gas was used (vs. balanced salt solution), the risk ratio went up 4-fold, clearly suggesting this gas must also promote some inflammatory pathways. Also, the use of a single-piece intraocular lens was 2.7 times more likely to cause this synechial formation. Exactly why a single-piece lens may be more likely to cause this is not quite clear. There may be a mechanical effect from the bulk of the lens or inflammation may be induced as well. Nonetheless, these are some clear clues to help us unravel this particular problem.

A key and common measurement is determining the intraocular pressure. Goldmann applanation tonometry (GAT) has been the gold standard for decades. Now we know that GAT gives a falsely high reading with increased corneal thickness and a falsely low reading with thinner corneas. Finding a device, therefore, not affected by corneal thickness and, hopefully, better able to tell us how to treat patients with glaucoma has been the subject of many studies.

The paper by Roszkowska et al. [Ophthalmologica 2009;223:244–249] specifically looked at dynamic contour tonometry (DCT) as a possible alternative to GAT when considering corneal thickness. They show that DCT is a technique that appears to be independent of corneal thickness. Whether we will all move in this direction only time will tell; however, this paper clearly indicates that knowing what the intraocular pressure actually is would be better served by this newer technology in comparison to our old gold standard GAT.

Better ways of maintaining success with filtering procedures has been an ongoing concern since filtration surgery was born. Inhibition of fibroblast growth has been suggested as one avenue.

α -Tocopherol certainly is known to inhibit fibroblast growth, and therefore Golblum et al. [Ophthalmologica 2009;223:228–232], in a randomized clinical trial, compared placebo with 300 mg/day of α -tocopherol acetate, and found after 1 year of follow-up that this was unfortunately no better than placebo. Let us not forget that negative studies are often as important as a positive result in that they both give us clues about where to go in the future and help us from continually making the same mistakes over and over again.

Adenoviral conjunctivitis can be a miserable condition. Not only does the process often last for weeks, it can result in prolonged visual loss due to corneal complications and is highly infectious. Patients are very desirous of some form of treatment.

There has been no specific antiviral agent established for the treatment; however, cidofovir, zalcitabine, and stavudine have shown anti-adenoviral activity in vitro. The problem is concern about toxicity, in that cidofovir is known to cause lacrimal duct obstruction.

Inoue et al. [Ophthalmologica 2009;223:233–238] looked at toxicity in healthy albino rabbits with utilization of drops 4 times a day for 14 days. They looked at clinical scores and histopathology. Indeed, they documented that cidofovir showed a significant narrowing of the lacrimal canaliculus as a form of scarring response, as well as significant infection of the eyelids and conjunctiva. It does not make much sense to treat a condition where redness of the eyelids and conjunctiva are a problem with a drug that is as bad as the disease. Histology showed that there was an allergic reaction ongoing with eosinophilia, which is probably the core root cause of this problem.

Although zalcitabine and stavudine induced some eyelid and conjunctival infection, there was no change in the lacrimal pathways, and it appears as though they may be promising candidates for treatment of this miserable condition.

Our final paper takes us to the field of ocular oncology: specifically, melanoma. No matter what treatment is undertaken for choroidal melanoma, the big ongoing concern is the risk of metastasis, for which treatment to date is still not very good. Having some type of a serum marker for metastatic disease would be a huge plus when following these patients.

Haritoglou et al. [Ophthalmologica 2009;223:239–243] report very promising results for osteopontin (OPN) as a potential new marker, and they compared it to melanoma inhibitory activity (MIA) for the screening and detection of metastatic uveal melanoma. They found that OPN was much higher in patients with metastatic disease compared to those without detectable metastases, and, while MIA also correlated strongly with metastatic disease, there was no correlation between MIA or OPN levels in regard to choroidal tumor heights. This means that we could have 2 independent markers, both of which may be helpful, in determining metastatic disease. This certainly warrants additional studies that could help us with this vexing problem.

So, a fascinating array covering a spectrum of ophthalmology is presented within, with practical relevance to how we take care of our patients.

Randall J. Olson, Salt Lake City