Panel Discussion 1

J.		Smyth
----	--	-------

Roila: Medical Oncology – Perugia. I would like to ask Dr. Brown about the 2 studies comparing 8 versus 32 mg bolus doses. The results in terms of complete protection from vomiting are contrasting. Before you said that one should choose the dose of ondansetron on the basis of patients' prognostic factors. I disagree with this because I think that with these results, probably the best dose is always 32 mg. Have you any comments on this, Dr. Brown?

Brown: I think it's true to say that we can't find a full explanation as to why the 2 studies are different. We've looked very carefully at the patient characteristics, the sex ratio, the emetogenicity of the chemotherapy regimens etc., and there is no obvious explanation as to why there is a difference between the 2 studies. So at the moment, all I can tell you is that we have got 2 studies, they show somewhat different results, both of them show that single doses are effective, and I agree with you that there is a suggestion that 32 mg may be somewhat better than 8 mg. Whether all patients should receive 32 mg I think is a different matter. But I dare say if we took a straw poll of the people in the audience here we would get a very diverse set of opinions on that, as with so many other things in oncology relating to doses of drugs.

Koslow: Walter Koslow, Medical Oncology, Australia. I'd like to ask either Dr. Tyers or Dr. Brown if they could elucidate the pharmacology of the drug. We've had a number of statements saying that the half-life does not correlate with the effect we've seen. Now this is rare for drugs but does occur for some. And also we've seen a number of dosing schedules administered; 4-hourly, 6-hourly and 12-hourly. And also that intravenously the drug can be given once a day with useful effect. And so can it be given once a day orally?

Tyers: The half-life of ondansetron is about 3-3.5 h, and yet the effect seems to last for 24 h. So that's the clearest example that half-life is not relevant to the duration of the drug's action.

Brown: Yes, you're right in saying that the early studies which looked at giving repeated infusions every 2, 4, 6 and 8 h, didn't show any obvious schedule dependency in terms of efficacy, and now that we know that single doses are as effective as repeated short infusions I think we have the explanation for that. The question about whether you could give once-daily oral dosing is something that we've discussed quite a lot. I can see quite clearly why theoretically that would be a good idea, but we have to bear in mind the fact that a number of patients will continue to be somewhat nauseated, some of them will continue to have some emesis despite general good control, and that if a patient vomits the only dose they've had in the day then you will lose control totally. So to some extent having twice-daily administration is a safety factor with the oral dose.

Smyth: I would like to ask Dr. Tyers: When you were summarising the fact that there are both central and peripheral receptors, is it known that the sensitivity to antagonism from ondansetron is equivalent at those 2 receptor sites?

Tyers: Theoretically it should be and there is certainly a lot of evidence that ondansetron will get into the brain very effectively, so there is no reason to believe they're any different.

Anonymous: I have a question for Professor Jürgens. It has long been believed that ondansetron has a major role during conditioning prior to bone marrow transplantation. However, we observe refractory vomiting episodes during the re-infusion of bone marrow. I am curious to know if you have the same experience.

Jürgens: Well, whether vomiting seen during the re-infusion of the bone marrow is due to late effect from the previous conditioning or is due to preservatives associated with the re-infusion of previously frozen marrow is very difficult to track down. I do agree that this emesis is very difficult to control, and most centres rather than use single agents, including ondansetron, use combinations of anti-emetics.

Green: Michael Green from Melbourne, Australia. A question I'd like to address to Ian Olver but I'd like Dr. Brown to comment on it as well. The issue of oral ondansetron has been discussed, but because of the methodological problems that Ian Olver has raised with the carryover effect from an intravenous injection, I don't feel as though you've really addressed whether or not oral ondansetron actually has a true anti-emetogenic effect as compared to, let's say, a standard dose of metoclopramide. The De Mulder study suggests there is no difference between oral metoclopramide versus oral ondansetron, for delayed emesis.

Olver: Well, I think that there is a methodological problem, as you suggested, the carryover effect, and I think the difficulty is where you don't have this correlation between the pharmacokinetics of the drug and the duration of action. I don't believe that the direct-comparison answer has been satisfactorily resolved yet, just from those basic principles.

Brown: We agree with you that we haven't entirely resolved this issue yet and indeed at this moment we're about to embark upon 2 very large placebo-controlled studies to try and resolve the issue of exactly what the activity of ondansetron is in delayed emesis. There is clear evidence from the early work looking at doses of 16 mg 3 times a day in comparison with placebo, that at that dose at least ondansetron is active. How it compares with metoclopramide is still an unresolved question, because in the one study in which there was a comparison (the De Mulder study) there were problems with the carryover from the first 24-hour period, and we're not entirely sure that there wasn't a design fault in looking at that second period. So I agree with you basically that the problem is not yet resolved, although I think there is already adequate evidence that 5-HT₃ receptor antagonists do have activity in delayed emesis. Now exactly what delayed emesis is remains to be seen, and if you look at those patterns of emesis that we've shown you, I think probably what we're seeing is that the maximal emetic stimulus due to 5-HT occurs in the first 8-10 h after cisplatin (although it would be over a longer period after cyclophosphamide) and that there is some continued 5-HT-mediated emesis

going on for a number of days after that, although clearly at that stage other mechanisms are relatively more important than they would be in the first few hours after chemotherapy.

Borshi: Professor Borshi, Budapest, Hungary. I have a question for Professor Jürgens. You showed us that emesis control was somewhat worse in children than in adults with ondansetron. Now might that be related to the effect of dosage, that you limited the dosage to a maximum of 8 mg per child. Have there been any systematic dose-finding studies in children with ondansetron?

Jürgens: The dose limit of 8 mg should not have affected this result because it's comparable to a single dose given in adults, which is 8 mg. And increasing the dose has not been associated with a marked increase in effect. However, in view of data shown today showing that a single dose of 32 mg is very effective, we should now go back in our paediatric scales and find an adequate dose-finding scale, taking into account that we sometimes do use more aggressive combinations, and also taking into account that sometimes highly emetogenic compounds are given in short

infusions over one hour, or as long-term infusions over a series of 3-5 days and the ideal dosing might be different, depending on dosing schedule.

Aya: Aya, from Germany. I have a question to Professor Jurgens. You showed an interesting slide relating to TBI. We have found that the highest frequency of emesis is after the second or the third fraction of TBI. Do you have an explanation for this different frequency of emetic episodes? And do you think it is better to adjust the dose of ondansetron for fractionated TBI to give different doses for different fractions?

Jurgens: Well, the data that was presented here was the data from 15 patients who received a conditioning regimen, and the data was collected from various centres. So it only gives you a very vague idea that compared to routine and customary used anti-emetic regimens it seems to be that there is a benefit of using ondansetron. I cannot answer your question whether the different scheduling of radiation, 1-day fractionation compared to several-day fractions, requires different scheduling of daily dosage of ondansetron, but it is an area that one has to look into.