Introduction

M.		Marty
----	--	-------

Ondansetron, the first highly selective 5-HT₃ receptor antagonist to reach the market, is now approved (as ondansetron hydrochloride dihydrate) for use in over 70 countries worldwide. After 3 years of clinical use and over 4 million patient treatments ondansetron has established an important role in cancer therapy and represents a real advance in the management of chemotherapy- and radiotherapy-induced emesis. Since the submission of the clinical dossier to regulatory authorities in April 1989, clinical experience with ondansetron has grown considerably. For example, it is now well established that a single intravenous dose of ondansetron is as effective as the dose regimens used in the initial clinical studies [1] and that twice daily oral dosing is as effective as a 3-times-a-day schedule [2]. It is therefore timely to review more recent clinical trial data and the global experience with ondansetron to help us to use this agent in an optimal manner and to look ahead to the future potential of ondansetron in cancer therapy.

References

1 Brown GW, Paes D, Bryson J, Freeman AJ: The effectiveness of a single intravenous dose of ondansetron. *Oncology* 1992;49:273-278.

2 Dicato MA, Kaasa S, Campora E, Bleiberg H, Warner P, Vindovoghel A, Cunningham D, Liebhard A, Upadhyaya BK: Efficacy of twice daily versus three times daily oral ondansetron in the prevention of chemotherapy induced emesis: A randomised, single blind, multicentre study. *Clin Oncol* 1992;4:275-279.