Introduction

M.E.

Michel E.

Marty

Department of Medical Oncology, Hôpital Saint Louis, Paris, France

The symposium 'New Directions in Anti-Cancer Chemotherapy' focuses on the current use of and new indications for megestrol acetate (MA), a well-known hormonal agent with expanding indications both in cancer and AIDS, as well as new data pertaining to Taxol¹ (pacli-taxel). MA – an orally active progesterone derivative – is a safe and efficacious agent in the treatment of hormone-responsive metastatic breast cancer. The main data arising from its worldwide use are reviewed by Dr. M. Espie. Due to its known side effects – the main one being weight gain – MA is mostly used in second-line hormonal therapy of advanced breast cancer, where its activity and safety are well documented.

The weight gain observed in approximately 20% of patients treated with MA is obviously not dependent upon tumor type, but upon the drug's pharmacodynamic effects. Thus, recent studies have addressed the question of effective dosages to treat cachexia in cancer (Loprinzi et al) and AIDS (Von Roenn) patients: In both cases a dose of 800 mg/day proved effective in enhancing appetite with ensuing weight gain. Quality of life studies correlate with weight gain. Thus, MA can add to the control of quality of life in outpatients with advanced malignancy or AIDS.

This symposium focuses also on paclitaxel, which possesses a unique mechanism of action: Promotion of tubulin assembly and inhibition of its depolymerization. Initially, limited supply of the drug – extracted from the bark of the Pacific yew – was considered a major limitation to paclitaxel use. The supply, however, no longer limits its use. The activity of paclitaxel was initially documented in the treatment of advanced ovarian carcinoma after the failure of platinum-based chemotherapy. This symposium focuses on the activity of paclitaxel in advanced breast cancer. The optimal dose – as defined from a large randomized trial – appears to be 175 mg/m², although higher doses have been used (often with colony-stimulating factors) (Spielmann). Paclitaxel is the most active agent used so far – at least comparable to doxorubicin. This provides new avenues for the treatment of patients who have failed anthracycline-containing adjuvant chemotherapy, and also opens the field to more active combination regimens. Ongoing studies have established the activity of paclitaxel in the treatment of non-small cell lung cancer and head and neck carcinomas. Thus, current data clearly show that this agent enlarges the current therapeutic armamentarium for breast cancer patients, who should benefit from new paclitaxel-based combinations.

¹ Taxol is a trademark of Bristol-Myers Squibb Company.