Changes in the Production of Nitric Oxide and Superoxide by Inflammatory Cells in Liver Cirrhosis

E.		Masini ^a
L.		Mugnai ^a
M.		Foschi ^b
G.		Laffi ^b
P.		Gentilini ^b
P.F.		Mannaioni ^a

^aDepartment of Preclinical and Clinical Pharmacology, and ^bClinica Medica II, School of Medicine, University of Florence, Italy

Key Words

Neutrophils
Monocytes superoxide
Nitric oxide
Liver cirrhosis

Abstract

Superoxide (O_2^-) and nitric oxide (NO) production by polymorphonuclear leukocyte (PMNs) and monocytes in patients with liver cirrhosis were evaluated. PMNs obtained from cirrhotic patients were less effective than those from controls in producing O_2^- after stimulation with opsonized zymosan, while they were more effective in producing NO, as shown by the inhibition of platelet aggregation and by the increase in cGMP content. NO synthase activity was higher in leukocytes from cirrhotic patients than in controls. A correlation was found between the cardiac index and the observed changes in the inflammatory cells.

Correspondence to: Dr. Emanuela Masini, MD, Department of Preclinical and Clinical Pharmacology, Florence University, Viale Morgagni 65, I-50134 Florence (Italy)

Introduction

Materials and Methods

Alterations of polymorphonuclear leukocyte (PMN) and monocyte (M) functions in chronic liver diseases contribute to the augmented susceptibility to infections in cirrhotic patients. It has been proposed that the excessive release of vasodilatory substances could be involved in the hyperdynamic circulation, characteristic of cirrhotic patients. However, clear evidence has not yet been obtained [1]. An altered synthesis of lipid mediators and increased synthesis of nitric oxide (NO) have been proposed recently to explain the augmented susceptibility to infections and the derangement in systemic hemodynamics occurring in cirrhotic patients with ascites.

The aim of this study was to evaluate superoxide O_2^- and NO production from PMNs and Ms from cirrhotic patients with ascites and hyperdynamic circulation.

Twelve cirrhotic patients with ascites and ten healthy volunteers were participated in the study. Diagnosis of cirrhosis was made by clinical history, clinical examination, laboratory findings and liver biopsy. Hemodynamic evaluation was performed by color Doppler ultrasound. PMNs and Ms were isolated after dextran sedimentation on a Lymphoprep gradient and then isolated from the leukocyte-rich plasma with an elutriation system using a Beckman J-6N/E centrifuge with a

JE-6B elutriation rotor and Sanderson chamber [2]. The production of O₂ from PMNs was determined by the reduction of cytochrome c [3], while NO production from PMNs and Ms was evaluated using their capacity to inhibit thrombin-induced platelet (PLT) aggregation and increase cGMP content in PLTs coincubated with PMNs or Ms. NO synthase (NOS) activity was evaluated by measuring [¹⁴C]-citrulline formation from [³H]-arginine.

©1995 S.Karger AG, Basel 1018-2438/95/1073-0197 \$8.00/0

Fig. 1. Correlation between cardiac index and NOS activity PMN (a) and M (b) monocytes from cirrhotic patients.

NOS activity (log dpm/μg protein) PMNs

NOS activity (log dpm/μg protein) Ms

Results and Discussion

The results show that PMNs from cirrhotic patients have a reduced capacity to produce O₂ compared to controls (97.3 ± 6.2 and 43.7 ± 4.3 nmol/10⁶ PMNs, respectively, $p < 0.05$), after stimulation with opsonized zymosan. In contrast, PMNs and Ms from cirrhotic patients were more effective than those from controls in inhibiting PLT aggregation. Preincubation of PMNs for 60 min with NG-monomethyl-L-arginine (L-NAME) significantly reduced the inhibitory action on thrombin-stimulated PLT aggregation. When PLTs were coincubated with PMNs as well as Ms of cirrhotic patients, cGMP levels were significantly higher in leukocytes from cirrhotics than from controls (13.52 ± 4.68 and 3.28 ± 1.52 fmol/10⁶ PMNs respectively, $p < 0.05$). NOS activity was significantly higher in cells isolated from cirrhotic patients, and a linear correlation was found with the patients' cardiac index (fig. 1).

In PMNs isolated from cirrhotic patients, these results indicate an impaired capacity to generate O₂ in response to different stimuli, and an increased generation of NO. Our data confirm the augmented susceptibility to infections in these patients, due to the reduced production of O₂. The data also suggest a role for NO in the pathogenesis of the hyper-dynamic circulation associated with cirrhosis.

References

- Bosh J, Gines P, Arroyo V, Rodes J: Hepatic and systemic hemodynamics and the neurohumoral system in cirrhosis; in: Epstein M (ed): The kidney in Liver Disease. Baltimore, Williams & Wilkins, 1988, pp 286-305.
- Almici C, Mangoni L, Carlo-Stella C, Rizzoli V: Human peripheral blood mononuclear cell subfractioning using counterflow centrifugation-elutriation. *Haematologica* 1991;76:89-93.
- Laffi G, Carloni V, Baldi E, Rossi ME, Azzari C, Gresele P, Marra F, Gentilini P: Impaired superoxide anion, platelet-activating factor, and leukotriene B₄ synthesis by neutrophils in cirrhosis. *Gastroenterology* 1993;105:170-177.

Masini/Mugnai/Foschi/Laffi/Gentilini/ Mannaioni
NO and CK in Liver Cirrhosis