Introduction

F.

Fraschini

Department of Pharmacology, University of Milan, Milan, Italy

P-Lactam antibiotics are the group of antibacterial agents most commonly used for treating serious infections. The skilful manipulation of the penam and cephem structures underlying these substances has resulted in molecules exhibiting an extremely high intrinsic activity and a broad spectrum of action unimaginable 10 years ago. These favourable characteristics are supplemented by the well-documented tolerability of the penicillins and cephalosporins, which makes these antibacterials the most suitable in absolute terms.

One of the factors historically limiting the use of the first antibiotics of this series – and the one that proved most difficult to modify – was the short elimination half-life. This made it necessary to give several doses each day, which, particularly in the case of injectable preparations, is a serious obstacle to good compliance. This often leaves no alternative to hospitalizing the patient, and also has a serious impact on the economic aspect of treatment (medical and paramedical staff costs, plus the cost of materials).

Ceftriaxone (Rocephin®) is a third-generation methoximinocephalosporin with a high degree of stability in respect of

bacterial (3-lactamases and with a broad spectrum of action covering all the Entero-bacteriaceae, Haemophilus influenzae, non-enterococcal streptococci, methicillin-sensitive staphylococci and most anaerobes. It is partially active against Pseudomonas species.

The main characteristic distinguishing ceftriaxone from the other third-generation cephalosporins is its long half-life (8 h) which means that it can be given as a single daily dose even in the treatment of extremely serious infections. This unique pharmacokinetic property is attributable to the high rate of binding to plasma proteins, coupled with extremely low total clearance.

The combination of these two properties is responsible for the high and long-lasting plasma concentrations that produce a gradient highly favourable to the diffusion of the agent through the tissue.

Numerous authors have reported the persistence of high concentrations of microbiologically active substance for over 24 h in various regions of the body.

It will be shown – perhaps for the first time in the case of (3-lactam antibiotics -

2

Fraschini

that the bacteria present in the septic foci of peripheral tissue are exposed to stable and continuously inhibitory concentrations of the antibacterial agent, whereas the use of traditional (3-lactams results in fluctuations between peak concentrations and (often subinhibitory) troughs. Now that the therapeutic value of ceftriaxone has been confirmed by several years of use, it is appropriate to reassess critically some of the variables of antibiotic treatment, such as dosage schedule, total duration of therapy for the individual infections compared with traditional re-

gimens, and the significance of in vitro evaluation parameters compared with in vivo parameters. This day of study will bring together some of the clinical experience gained in this new approach to the treatment of various types of infections – including bronchopulmonary infections, meningitis, biliary tract and ENT infections – while taking into account the economic and management advantages of single daily dosing of ceftriaxone as compared with multiple dosing of the reference antibiotics.