Discussion

M.		Monset-Couchard
C.		Amiel-Tison

Chairman: Dr. Rih

1. Dr. Klaus's paper

Dr. Raiha: Has anybody anything to say about bag ventilation?

Dr. Minkowski : It would be useful to have some more details about the technics of this method.

Dr. Amiel : Before bagging, a tube is placed in the stomach to avoid distension and no oral feeding is given during the ventilation period. Nurses are trained to appreciate which pressure they are giving, on a model with a manometer. The desirable pressure is around 30-35 cm H₂O. The rhythm is about 30/min, every 5 out of 20 min, sometimes more if no improvement is observed. The results are appreciated on blood gases and the weaning period is guided by blood gases too.

Dr. Stern : The only problem is that it requires personnel!

Dr. Geubelle: Are the nurses taught to try to inflate the baby when inspiration is starting ? If the infant is expiring and the nurse is inflating, there will be some resistance and I don't know what will happen.

Dr. Stahlman : This type of procedure should be reserved for small communities hospitals where no intensive care unit is available and without possibility of sending the baby to a big center.

Dr. Minkowski: I agree, it should be an auxiliary method or a replacement method.

Dr. Strang: I think it represents one trend that we might see grow, or even two trends: one is to ventilate more babies earlier and the other is not to ventilate babies at all! I would say that the best method for HMD babies is a well-controlled environmental temperature, a very well controlled pH and a 100 percent inspired oxygen. And keep it all the way until recovery . . . , that is one view.

Dr. Martin-Bouyer : This method is surely interesting because it allows an early artificial ventilation and requires just a nurse to use it. But it would be good to know the criteria leading to start the procedure.

Dr. Amiel : The results are excellent on score-III infants (PO₂ less than 50 mm Hg, or two apneas): 16 survivors out of 25 infants.

Dr. Swyer : But the important things to know are what were their weight and gesta-tional age!

Dr. Amiel: Mean birth weight was 1700 g for the whole group.

Dr. Canet : Is it really prudent to get repeated large oscillations in PO₂ in a newborn baby?

2. Dr. Swyer's paper

Dr. Stern: I would agree, it is better to have X-rays going to the baby rather than the baby going to the X-rays. That is an error that a number of people have made in the construction of intensive care units.

1 Morning session: 19. 4. 69.

120

Discussion

Dr. Swyer : We have a portable unit which goes to the baby at the bed-side. We also have an amplifier on the floor and complex radiology is carried out on the floor.

Dr. Geubelle: In our experience, nurses enjoy reading figures rather than looking at a moving needle, for heart rate for example. Don't you think that digital records might be useful for nurses ? Most of these apparatus are built for technicians and not for nurses I

Dr. Swyer: It is a question of extending to a digital reader. And there is a real educational problem for nurses and for doctors, in all monitoring systems.

Dr. Stahlman : We tend to think of monitoring as mechanical problems. It is equally important to monitor a baby by chemistry, PaCO_2 , anemia, hypoglycemia; in addition, I would like to suggest that in future, we are going to have the ability to evaluate clotting factors.

Dr. Strang: I would like to propose something controversial. When it comes to biochemical monitoring, with fixed apparatus, and when it comes to X-rays, there is in the course of HMD a silent period: at the beginning, it is necessary to estimate the severity of the disease to eliminate a diaphragmatic hernia, and so on; but having obtained these data, on the second and the third day of the illness, repeating the testing is really not a need and we have babies that we literally do not touch ! We don't change them, we don't touch them for 2 or 3 days. Blood gases can be drawn through the catheter without touching the baby.

Dr. Stern: As for the silent period, hypocalcemia should be detected and not taken for part of the disease.

Dr. Strang : What about the use of the prolonged QT interval ? A very useful thing in diagnosing hypocalcemia.

3. Dr. Stern's paper

Dr. Raiha: I would like to ask Dr. Stern about the number of cases of kernicterus. I have not seen a case of kernicterus in our unit for years (no clinical cases, nor anatomical ones)!

Dr. Stern : History of kernicterus: only the larger babies exhibited the classical signs of kernicterus. The smaller babies just had some apnoeic spells and died. This, therefore, requires first-class pathologic examination.

4. Dr. Larroche's paper

Dr. Strang : Here are infants who die. And the problem is: To what extent are the changes due just to primordial changes in the vessels ? And, should you see these changes in survivors ? It may not be the same. However, your last two pictures (late deaths) raises the problem of late changes for peripheral vascular accidents between 15 and 20 years ?

Dr. Stahlman: 317 patients had catheters; 214 lived, 103 died and went to autopsy. In those survivors, there were only two complications, one with symptoms and the other one was picked up by accident. One had oesophageal varices with bleeding and one had complete occlusion of the right iliac artery, picked up on aortography. In those babies who died, there are 11 complications with one obviously leading to death. The mean duration of catheters was 48.8 hours.

Dr. Swyer: It has been shown that with side hole catheter, the dead-space tends to clot. Another comment will be about the discrepancy between the clinical signs and the frightening pathological findings: this is terribly important.

Dr. Minkowski : Dr. Larroche, can you set the problem of how long these lesions have been organized, showing that they are not just agonic artefacts ?

Dr. Larroche : When the clots are organized with very solid mass of fibrin and cellular reaction and, when the aspect of the intima is not one of recent lesion, these are the only anatomical criteria.

Dr. Stern : Our own policy is not to use umbilical vein at all.

5. Dr. Stahlman's paper

Dr. Stern: Dr. Stahlman said that an incidence of about 60% of intracranial hemorrhages was observed (slide). These are the autopsy findings of our first 91 cases of our

Discussion

121

respirator series, and we had autopsies on all the 37 deaths. All had HMD. The incidence of intracranial hemorrhage is about 62%. These are not traumatic hemorrhages, they are intraventricular and probably anoxic in nature. When we stepped to the negative pressure respirators, it had been suggested by a number of people that the rapid pressure changes might cause hemodynamic changes in cerebral circulation; however, our findings of no evidence of difference in intracranial hemorrhages in HMD who died, of comparable birth weight, without being in the respirator, are very encouraging. It shows that the respirator in itself does not account for the intraventricular hemorrhage.

Dr. Stahlman: It has been interesting to see that using positive pressure, we have had the same incidence of intracranial hemorrhage with hyaline atelectasis. Somebody said that we were prolonging death rather than life and that may be the explanation for the increasing rate of complications, were it hemorrhage, or infection, or pulmonary fibrosis, or whatever.

Dr. Swyer: Data about clotting are scanty in premature infants. We agree entirely with Dr. Stahlman that there are multiple defects in clotting factors in sick infants but, in fact, they seem to occur with equal frequency in quite normal prematures.

Dr. Stahlman: There is no such thing!!!

Dr. Strang : We think that one form of hemorrhage seen clinically in the newborn infants is pulmonary hemorrhage. We have analysed pulmonary hemorrhages clots and found that they have a very low hematocrit and that the distribution of protein molecules in the substance is such that its presence can only be explained by filtration. And it suggests that most of these cases are in fact hemorrhagic pulmonary edema.

Dr. Stahlman: At the present time, we are investigating the use of heparin in selected babies with a control basis.

Dr. Pouillance: In the neonate and particularly the premature infant, on what basis do you make the diagnosis of intravascular coagulopathy, without mentioning etiology ? We studied some newborns without respiratory distress, without bleeding and, in the three first days, we found prothrombin levels less than 50% (around 30%), platelets counts around 50 000/mm³ or less, and fibrinogen around 2 g. It is difficult to imagine how it is possible to diagnose intravascular coagulopathy by comparison with this type of basic data.

Dr. Stahlman: In our own series, this is not so. In Segal's series (in Journal of Pediatrics, about 18 months ago), the platelet counts were not depressed. As far as fibrinogen is concerned, our fibrinogen levels in our own babies were exactly what they are in adults. I don't believe that a depressed fibrinogen below 1.80 would be surely abnormal in a newborn baby. For the prothrombin time (completely independant of this syndrome) in a normal baby, the data were in very narrow span. The standard deviation was extremely narrow. It becomes a secondary problem in consumption coagulopathy: 100% of babies who died had abnormal prothrombin time and 84 to 94% had abnormal thromboplastin time.

Dr. Pouillance: Once a disseminated intravascular coagulopathy is identified, it becomes necessary to use heparin. What are your doses and route?

Dr. Stahlman: We are following the thromboplastin time and the various clotting times to decide whether or not our babies are heparinized. A given fixed dose of heparin is impossible; we try to double the thromboplastin time as an evidence of full heparinization. The dosage we are using at the present time is 4 mg/kg every 6 hours, which is a normal adult heparinization dose, initially as a push, then as a continuous intravenous infusion. Using fibrinogen would just provide more fibrinogen for clotting and accentuate the lesions. If the fibrinogen is extremely low and if there is active bleeding, heparine should be administered first.

Dr. Strang : I don't think it is very dangerous to heparinize a baby and, many of us have been using heparinized blood for exchange transfusion all the time. I never saw one of these babies bleed.