Reply

E.		Rongioletti ^a
R.		Betti ^b
C.		Crosti ^b
A.		Rebora ^a

Departments of Dermatology, University of ^aGenoa, and ^bSan Paolo Hospital, University of Milan, Italy

Franco Rongioletti, MD, Clinica Dermatologica, V.le Benedetto XV, I-7-16132 Genova (Italy)

Lumping is a difficult exercise, certainly more difficult than splitting. That is mainly because the 'lumper' of rare entities bases its activity on reviewing literature rather than on personal observations. Unfortunately, case reports are not accurate all the times.

We included mosaic acral keratosis (MAK) in the marginal papular acrokerato-dermas (MPAK) because we considered MAK as being a form of acrokeratoelastoidosis (AKE) without elastorrhexis. In fact, Jacyk's patients with MAK [1] looked identical to one of the original cases of AKE, as the picture in Costa's paper clearly shows [2] and Rook's textbook suggests [3]. In addition, one of Jacyk's patients had minute papules over the fingers (no better indication is provided); in cases of AKE, papules also affect the dorsa of the feet and ankles as they do in MAK patients; and both Jacyk's patients had palmoplantar hyperkeratosis, like most of the AKE patients [4, 5].'

We now understand that 'none of the patients... had lesions along the borders of the hands and feet'. As we took marginality as the distinctive character, we agree that MAK cannot be included in MPAK.

As AKE is likely to be a familiar disorder, we included MAK in the hereditary group. We wonder, however, whether the number of reported cases is enough to speculate on etiology, especially so as the family history may be silent even in AKE.

We classified keratoelastoidosis marginalis and digital papular calcinosis as acquired forms and reported UV light as a likely cause of them (see tables 1 and 2 in our paper [6]). Of course, we agree with Jacyk on this point. Nonetheless, even excluding the cases of MAK, we maintain that the other forms can be classified on a clinicopathological basis. Keratotic papules along the borders of hands and feet are the common clinical feature that justifies their aggregation under the title of MPAK. The main distinctive criterion is the presence or lack of elastorrhexis for the hereditary forms and solar elastosis for the acquired ones.

References

Jacyk WK, Smith A: Mosaic acral keratosis. Clin Exp Dermatol 1990;15:361-362.

Costa OG: Acrokeratoelastoidosis. Arch Dermatol Syphilol 1954;70:228-231.

Griffiths WAD, Leigh IM, Marks R: Disorders of keratinization; in Champion RH et al (eds):

Rook's Textbook of Dermatology, ed 5. Oxford, 1992, pp 1381-1382.

Hight AS, Rook A, Anderson JR: Acrokeratoelastoidosis. Br J Dermatol 1982; 106: 337-344.

Shbaklo Z, Jamaledine NF, Kibbi AG, et al: Acrokeratoelastoidosis. Int J Dermatol 1990; 29:333-336.

Rongioletti F, Betti R, Crosti C, Rebora A: Marginal papular acrokeratodermas: A unified nosography for focal acral hyperkeratosis, acrokeratoelastoidosis and related disorders. Dermatology 1994;188:28-31.

© 1995 S. Karger AG, Basel 1018-8665/95/1902-0179 \$ 8.00/0