In Reply

B.M.

Czarnetzki

Basel

Prof. Dr. med. B.M. Czarnetzki, Clinical Research, Dermatology, F. Hoffmann-La Roche Ltd., CH-4002 Basel/Switzerland

Sir.

Thank you very much for your comments on the vehicle used in your clinical trial of MC 903. We therefore both agree that formulations in petrolatum give the most favorable results for all presently available vitamin D derivatives.

Regarding the time until healing, it is true that you described some clinical changes already after 1 week of therapy. These changes were only minimal, however, and clearing of lesions was not evident, even on histology, by 6 weeks of treatment, with overall improvement not even reaching a score of 2 [moderate improvement; see fig. 5, ref. 1] in your patients. Therefore, healing with MC 903 should be regarded as slow which is the case with most presently available therapies for psoriasis.

To investigate further the relationship between circulating anti-mitochondrial antibodies and lichen planus, we studied the sera of 30 patients with lichen planus seen consecutively in the Departments of Dermatology in Sheffield and Chesterfield (UK). The anti-mitochondrial antibodies were detected by indirect immunofluorescence using a rat kidney substrate, bound antibody being localised with FITC conjugated swine anti-human immunoglobulin. A positive result was obtained in only one of the 30 patients, a 59-year-old man with no clinical or biochemical evidence of liver disease.

It appears that there is no general increase in the frequency of anti-mitochondrial antibodies in patients with lichen planus and our study offers no support for an immunopathological association between idiopathic lichen planus and primary biliary cirrhosis. We concur with the suggestion that the immunopathology of lichen planus in patients with primary biliary cirrhosis is likely to resemble that in patients with graft-versus-host disease [7]. The pathogenesis is likely to be different in patients with idiopathic lichen planus who form the great majority seen in dermatological practice.

Reference

Kragballe K, Beck HI, Søgaaard H: Improvement of psoriasis by a topical vitamin D3 analogue (MC 903) in a double-blind study. *Br J Dermatol* 1988;119:223-230.

Dr. A.J.G. Mc Donagh MB, MRCP F. Leyva-Leon BSc. MB D.J. Gawkrödger MD. MRCP A. Milford Ward MD. FRCPATH Depts. of Dermatology & Immunology Royal Hallamshire Hospital Glossop Road Sheffield S10 2JF (UK)

Lichen planus Is Not Commonly a Skin Marker of Primary Biliary Cirrhosis

Sir.

Lichen planus is a common skin disorder accounting for about 1% of referrals to dermatology clinics [1]. Most cases are of the 'idiopathic' variety but a similar reaction pattern may occur as a result of an adverse drug reaction or in graft-versus-host disease. There is controversy

concerning a possible association of lichen planus with liver diseases. Although a high incidence of liver diseases other than primary biliary cirrhosis is reported in a recent study of lichen planus patients from Italy [2], previous studies have been inconclusive in this respect [3]. The best evidence for an association between liver disease and lichen planus comes from studies on the minority of subjects in whom lichen planus is associated with primary biliary cirrhosis [4,5]. These patients have circulating anti-mitochondrial antibodies but not all patients with a positive test for anti-mitochondrial antibodies have primary biliary cirrhosis. Such antibodies have been detected in patients with lichen planus in whom there was no evidence of liver disease [6].

References

- Black MM: Lichen planus and lichenoid eruptions; in Rook A, Ebling FJG, Wilkinson DS, et al (eds): Textbook of Dermatology. ed4. Oxford. Blackwell. 1986. pp 1665-1685.
- Cottoni F, Solinas A, Piga MR, et al: Lichen planus, chronic liver diseases and immunologic involvement. Arch Dermatol Res 1988; 280(suppl):S55-S60.
- Sarkany I: The skin-liver connection, in Exp Dermatol 1988;13: 151-159.
- Graham-Brown RAC, Sarkany I, Sherlock S: Lichen planus and primary biliary cirrhosis. Br J Dermatol 1982;106:699-703.
- Powell FC, Rogers RS, Dickson ER: Lichen planus, primary biliary cirrhosis and penicillamine. Br J Dermatol 1982;107:617.
- Triger DR, Charlton CAC, Milford Ward A: What does the anti-mitochondrial antibody mean? Gut 1982;23:814-818.
- Epstein O, Thomas HC, Sherlock S: Primary biliary cirrhosis is a dry gland syndrome with features of chronic graft-versus-host disease. Lancet 1980;i: 1166-1168.