In Reply

K.		Welp ^a
U.		Gieler ^a
M.		Ständer ^b
H.C.C.		Friederich ^a

^aDepartment of Dermatology University-Clinic Marburg, Deutschhausstrasse 9, D-3550 Marburg (FRG); ^bPsoriasis-Clinic, Am Bade 4, D-444 Bad Bentheim (FRG)

K. Welp, U. Gieler, M. Ständer, H.C. Friederich, Department of Dermatology University-Clinic Marburg, Deutschhausstrasse 9, D-3550 Marburg (FRG), Psoriasis-Clinic, Am Bade 4, D-444 Bad Bentheim (FRG)

Sir.

We have read with great interest the letter by Beer et al. about the concomitance of psoriasis and atopic dermatitis (AD). We agree with the authors' opinion, that there are some interesting immunological comparisons. In Europe as well as in the US psoriasis is seen in approximately 1-2% of the population [1].

Incidence data for AD range from 0.7 to 2.4 depending upon the age as well as the ethnic group studied [2]. In a national screening survey performed in the US the prevalence of AD was estimated to be from 7 (adults) to 19 (children) per 1.000 [3]. In England a survey revealed a 3.1 % prevalence among children under the age of 5 [4]. As in psoriasis AD tends to become even more frequent in the northern countries of Europe [5].

Since both diseases occur at comparatively high frequencies especially in the younger age group, it would be reasonable to suspect that

in a considerable number of patients both skin diseases are simultaneously present. The calculated rates of simultaneously occurrence of both diseases should amount from 0.05 to 0.30% among the Caucasian population. However, the rate of concomitance which should be statistically expected in both diseases has not yet been proven.

Within a patient sample of 1,065 psoriatics attending an inpatient psoriasis clinic within half a year we found 18 (1.7%) cases with an additional diagnosis of AD confirmed by histological investigation. This finding was within the expected range. A positive history of the disease among family members could be found within 44% of all cases with psoriasis. 39% with AD and 33% with rhinitis allergica or asthma bronchiale. Besides typical psoriatic lesions and atopic eczemas (which were also examined histologically) the following marked symptoms were found in addition: pruritus within 100%; white dermographism within 89% (16 patients); a positive sign of the so-called Auspitz phenomena within 83% (15 patients); rhinitis allergica or asthma bronchiale within 72% (13 patients). In 89% of all patients (16 patients) the IgE level was increased with a geometrical mean of 888 U/ml. The HLA-Cwb was positive within 50% (9 patients). Since we focused on patients with typical and fully developed symptoms and did not include patients with minimal or mixed symptoms the frequency of concomitant forms could be supposed to be much higher. A possible explanation of the low concomitance of both diseases found until now might be that in an acute state of disease psoriatic lesions might dominate AD

and makes an exact diagnosis more difficult. Itching and seborrhea as the symptoms of a therapy thought to be too 'aggressive' might represent the first signs of an additional AD. With regard to the incidence of both diseases it should be expected that a lack of concomitance should rarely be found.

Therefore, in a more thorough documentation this combination should be found more frequently.

References

Christophers E, Krueger G: Psoriasis; in Fitzpatrick et al (eds): Dermatology in General Medicine. New York, McGraw Hill, 1987, pp 461-491.

Christophers E: Anlagebedingte Hauterkrankungen. Hautarzt 1984(supplVII):3-6.

Hanifin JM: Atopic dermatitis. J Clin Immunol 1984;73:211-222.

Walker RB, Warin RP: Incidence of eczema in early childhood. Br J Dermatol 1956;68:182-187.

Christophers E: Leukocytes and skin diseases: The conflicting patterns of inflammation in atopic dermatitis and psoriasis. 85th Annu Meet Jap Dermatol Ass, Kyoto Kaikan, 1986.