Papulonecrotic Tuberculides on Glans penis

C.	Chikako	Nishigori
----	---------	-----------

Kyoto

Chikako Nishigori, Department of Dermatology, Faculty of Medicine, Kyoto University, Shougoin, Sakyo-ku, Kyoto (Japan)

In Reply

Dear Sir,

We are very much obliged to Dr. Kumar and Dr. Sharma for their interesting cases and pertinent comments on our paper [1]. As we mentioned in our paper, we used the term 'penis tuberculides' in the case in which the lesion is principally limited to the penis. In this sense, their 3 cases are consistent with 'penis tuberculides', however, in the 4th case the diagnosis of papulonecrotic tuberculides is more proper because the penile lesion is a part of the widespread papulonecrotic tuberculides.

Although Dr. Kumar and Dr. Sharma cited cases of penis tuberculides from Asian countries and Spain and mentioned unimportance of genetic roles, there still remains a question why such cases have scarcely been reported so far from other countries such as England, France, Germany or the United States. Although the report by Wong et al. [2] from Hong Kong was interesting, they did not describe if the eruption was almost exclusively limited to the penis or not. That is the reason why we did not cite their paper.

References

Nishigori, C; Taniguchi, S.; Hayakawa, M.; Ima-mura, S.: Penis tuberculides; papulonecrotic tuberculides on the glans penis. *Dermatologica* 172: 93-97 (1986).

Wong, K.O.; Lee, K.P.; Chiu, S.F.; Tuberculosis of the skin in Hong Kong (a review of 160 cases). *Br. J. Derm.* 80:424-429 (1968).