Opening Remarks

H.		Schmidt
----	--	---------

Dermato-Venerologisk Afdeling, Odense Sygehus, Odense, Danmark President of the International Symposium of Bifonazole, Copenhagen

Prof. Dr. H. Schmidt Odense Sygehus Dermato-Venerologisk Afdeling DK-5000 Odense (Danmark)

iVi% Hjorth should have presided at this event. I am not Niels Hjorth, not in any mean or way. Niels Hjorth whose name is on the front page of the programme fell ill, and consequently was unable to preside here. He is now in good recovery, but very sorry that he could not take his chair at this meeting. Most of you present know Niels Hjorth and are acquainted with his energy and vivacity. It is a very great pity, that he could not attend, and I know that all of us will send him best wishes for a rapid recovery.

There has been a change in the mycolog-ical diseases seen and diagnosed by European dermatologists during the last 40–50 years. Thus, before the Second World War cattle ringworm was not only more common in our communities, but the poor patients suffering from this mycosis were often over-treated by iodine solutions followed by dermatitis provoked by the treatment. The same was the case for tinea pedis in which disease contact dermatitis often followed the treatment with antimycotics.

These dermatitis cases following treatment of fungi diseases were especially common in Germany, where many patients were treated with organic mercurials. In Denmark we used enormous amounts of Arning's tincture before the griseofulvin era, and this tincture was a serious sensitizer due to its content of benzoin and paraben esters. During that time nail mycoses e.g. were not accessible for treatment, whatsoever, and the treatment for mycosis of hair was extremely bad. Epilation of scalp hair by means of X-ray was not only inhuman due to its cosmetic result, where the patient lost all the hair, but even not without risk. It is known from the textbooks of those days, how desperate the situation in reality was – the available treatment did after all not interfere with nature's own cure – if any.

With the introduction of griseofulvin this state changed to a great extent. Thus, I remember Prof. Harvey Blank later on to an Academy meeting in Chicago told that when griseofulvin was introduced on the American market in the fifties, he was able to lecture over the effect of the new drug in 15 patients at the Academy meeting 5 months later – today, he added, at least 3–4 years might elapse before the product was allowed for human beings due to the restrictions and claims set by FDA.

Of course treatment failures were also

2

Schmidt

registered with griseofulvin, but this was to a certain extent alleviated with the introduction of ketoconazole, in the late seventies.

Concerning dermatophytoses not involving nails or hair, imidazoles represented a great step forward. Introduction of this group of drugs was so effective that most dermatophytoses will be cured in general practice. Thus, those patients who are not cured are either therapeutic defaulters or misdiagnosed – which happens now and then. And you can't blame the pharmaceutical industry for our wrong diagnoses.

In Denmark, the number of mycoses treated in hospitals is going down – in England a recent survey has shown increasing numbers of examinations, but a rather steady number of patients suffering from anthropophilic fungi infections. These ciphers might be taken as an expression for efficiency of the imidazole and its derivatives used in general practice as a first choice treatment. However, no statistics are available from the practice branch, it is pure guess if the epidemic has been brought down due to modern therapy or if the decrease has occurred for other reasons.

Whatever is the reason to the decrease, no doubt exists that modern treatment has made life easier for those suffering from dermatophytic lesions. Formerly mycoses such as favus was a disease of bad hygiene seen i.a. in Greenland and in poorer circles in Europe – at the present time mycotic skin disorders are rather expression of welfare e.g. caused via the introduction of bathing facilities and public swimmingpools.

In order to speak a little further on individual mycotic infection it should be mentioned, that *Trichophyton rubrum* infections for instance, were nearly unknown in Europe before World War II, and it was supposed to have been introduced by American soldiers. During the Second World War *Trichophyton rubrum* infections increased, all over Europe, an increase which was probably real even if the diagnoses with the typical change in palms and soles may have passed unrecognized earlier, thus to quote Goethe: 'Was du weisst, das siehst du' (What you know you do observe, too).

During later years foreign workers coming to work in Europe have imported some exotic fungi, e.g. *Trichophyton violaceum*, regarded as a curiosity and not a danger to our hemisphere. Also European tourists travelling to the countries far away may import fungi of seldom sorts back to their homelands, and there giving rise to spread.

Industrial dissemination of fungus infections have been rather well studied, but despite this we still know very little about reinfections from e.g., shoes, socks, or stockings – this may be regarded a challenge to both doctors and the pharmaceutical industry. Those patients who suffer from dermatophytosis and who consequently get a modern and effective therapy, have good reasons to be pleased, which will probably appear also from the lectures to be given during the days to follow.

I. I wish all of you welcome to Denmark
to Copenhagen
and to this Bayer symposium on bifonazole.