Touch: A Biological Regulator of Growth and Development in the Neonate

S.M.		Shanberg ^b
T.		Field ^a
C.M.		Kuhn ^b
J.		Bartolome ^b

^bDuke University Medical Center, Durham, NC 27710 ^aUniversity of Miami, Miami, FL 33136

Mammalian development is markedly influenced by various environmental stimuli. However, those stimuli which are provided by the mother appear to be most critical for survival and growth. Evidence from this and other laboratories demonstrates that disruption of the mother-infant relationship elicits very specific physiologic and behavioral responses in both young animals and human infants which can severely affect their development. The consequences of interrupting this interaction range from changes in body temperature, heart rate, locomotor activity, and suppression of neuroendocrine and physiological responses following short periods of separation to retardation of growth and behavioral development during long-term separation. Each of these responses is initiated by loss of selective sensory cues, i.e. maternal olfactory, gustatory and thermal signals promote specific locomotor responses in developing rat pups. We have shown that even short-term separation of rat pups from their mother (MS) initiates a complex adaptive physiological response in the pup that includes: (1) a decrease of ornithine decarboxylase (ODC), the rate-limiting enzyme in the biosynthesis of polyamines, key regulators of growth and differentiation, (2) a reduction in DNA synthesis, (3) an inhibition of growth hormone (GH) secretion, (4) an increase in the secretion of corticosterone, and, (5) a suppression of tissue ODC responsiveness to exogenously administered GH, prolactin and insulin. This pattern of adaptation to maternal absence is not related to changes in nutritional factors or temperature regulation but results from a lack of a specific type of touching of the pup by the mother and occurs only during the first 18 to 20 days post-natally.

Tactile stimulation in the form of firm stroking with a brush at a frequency that mimics maternal licking patterns reliably prevents or reverses these effects of MS in the pup. The ineffectiveness of kinesthetic and vestibular stimulation in restoring normal physiology in the rat pup strongly supports the hypothesis that it is the tactile component of the intervention that produced the beneficial effects. Our identification of a specific type of touch as a neonatal growth requirement led us to test supplemental tactile stimulation in isolated very-premature (28–32 weeks gestation) human babies. It is well documented that a significant proportion of these babies do not gain weight in the same manner as term babies or normally growing fetuses of equal gestational age. The results of our study were marked. With supplemental stimulation, premature babies averaged a 47% greater weight gain/day, were significantly more active and alert during sleep/wake

behavioral observations, and showed more mature habituation, orientation, motor and range of state behavior on the Brazelton scale than non-supplemented control infants. The enhanced development also was reflected in a six-day shorter hospital stay at a cost savings of about \$3,000 per infant. Our more recent study confirms the positive effects of the tactile intervention on both weight and developmental gains, and demonstrates a significant enhancement in sym-patho-adrenal maturation in the premature babies.

A variety of studies from different laboratories indicate that endogenous opioids may have an important role in the regulation of development. Recently, we found that central but not peripheral injection of beta-endorphin (bE; an endogenous CNS opioid peptide) evokes a pattern of biological responses closely resembling those observed after MS. Specifically, intracisternal (ic) injection of bE reduces basal ODC activity as well as tissue ODC responsiveness to key trophic factors including GH and insulin in peripheral organs, and dramatically reduces DNA synthesis in developing rat tissues in a naloxene-reversible manner. Also, these effects caused by bE occur only during the first three weeks of postnatal life. Finally, we have shown that the suppression of peripheral DNA synthesis and ODC activity induced by both MS and bE occur through CNS mechanisms independent of the pituitary gland.

These studies highlight the integrative need of appropriate sensory input CNS function and end-organ physiology to maintain normal maturational development. The combined demonstrations which show that a specific type of tactile stimulation restores normal growth physiology both in MS rat pups and in isolated human preterm neonates suggest that similar hormonal and biochemical mechanisms might be involved. We suggest that this animal model can be used to investigate all aspects of this condition ranging from the afferent sensory stimuli triggering the response and the neuronal and neurochemical mechanisms “transducing” these stimuli to the specific biomolecular mechanisms operating in the cells of the target tissues.