Pulmonary Surfactant: An Immunological Perspective

Zissis C. Chroneos^{1,*}, Zvezdana Sever-Chroneos¹ and Virginia L. Shepherd^{2,3}

¹The Center of Biomedical Research, University of Texas Health Science Center at Tyler, Tyler, ²Department of Pathology, Vanderbilt University Medical Center and ³Department of Veteran Affairs, Nashville

Key Words

Lung • Surfactant protein • Alveolar macrophages • Collectin • Receptor • Inflammation • Innate immunity

Abstract

Pulmonary surfactant has two crucial roles in respiratory function; first, as a biophysical entity it reduces surface tension at the air water interface, facilitating gas exchange and alveolar stability during breathing, and, second, as an innate component of the lung's immune system it helps maintain sterility and balance immune reactions in the distal airways. Pulmonary surfactant consists of 90% lipids and 10% protein. There are four surfactant proteins named SP-A, SP-B, SP-C, and SP-D; their distinct interactions with surfactant phospholipids are necessary for the ultra-structural organization, stability, metabolism, and lowering of surface tension. In addition, SP-A and SP-D bind pathogens, inflict damage to microbial membranes, and regulate microbial phagocytosis and activation or deactivation of inflammatory responses by alveolar macrophages. SP-A and SP-D, also known as pulmonary collectins, mediate microbial phagocytosis via SP-A and SP-D receptors and the

coordinated induction of other innate receptors. Several receptors (SP-R210, CD91/calreticulin, SIRP α , and toll-like receptors) mediate the immunological functions of SP-A and SP-D. However, accumulating evidence indicate that SP-B and SP-C and one or more lipid constituents of surfactant share similar immuno-regulatory properties as SP-A and SP-D. The present review discusses current knowledge on the interaction of surfactant with lung innate host defense.

Copyright © 2010 S. Karger AG, Basel

Introduction

Pulmonary surfactant, a complex of lipids and proteins lining the alveolar surface, is responsible for lowering surface tension at the air-liquid interface thereby preventing alveolar collapse at the end of expiration [1, 2]. However, lung surfactant is also an integral component of the lung's innate immune system helping to control inflammation and to prevent microbial infections of the distal lung [3, 4]. Inside the alveolus, large and small aggregate vesicles of pulmonary surfactant lipids are

spatially and dynamically coordinated with the surface active monolayer at the air-liquid interface [5]. Surfactant composition and pool size is controlled by several physiological processes, including secretion, re-uptake, and recycling by alveolar type II epithelial cells and degradation by both alveolar type II epithelial cells and macrophages [6, 7]. All surfactant proteins contribute to the internalization of surfactant subtypes by type II epithelial cells [8-11]. The principal lipid constituents of surfactant are phospholipids, 80-85%, and cholesterol, 8-10% by weight. Phosphatidyl choline species comprise about 75% of surfactant phospholipid. Saturated dipalmitoyl phosphatidyl choline (DPPC) is the most abundant and critical surface-active phospholipid species that constitutes 50% (w/w) or more of lung surfactant phospholipid. Remaining phospholipids include phosphatidyl glycerol (12%), phosphatidyl ethanolamine (5%), phosphatidyl inositol (4%), phosphatidyl serine (1.5%), sphingomyelin (1%), and lysophospholipid (<1%). Optimal surfactant function requires the presence of four proteins named SP-A, SP-B, SP-C, and SP-D. Of the four, SP-A is the most abundant comprising 50-70% of surfactant proteins (w/w). However, SP-B, which comprises 10% (w/w) of surfactant protein is the most critical; its absence or dysfunction due to mutations results in respiratory failure and death shortly after birth [12-15]. Secreted surfactant consists of different subtypes that can be distinguished by differential or density gradient centrifugation as surface active large aggregate (LA) and surface inactive or “spent” small aggregate surfactant (SA) forms [16]. The LA and SA forms can be isolated by differential centrifugation at 40,000 and 100,000 x g, respectively. Greater than 95% of SP-A, SP-B, and SP-C fractionate with LA surfactant [17], while less than 10% of SP-D associates with LA surfactant [11]. The LA surfactant contains several morphologically distinct forms that are referred to as lamellar body-like, tubular myelin, and multilamellar vesicles. SP-A and SP-B are necessary for the conversion of lamellar bodies into tubular myelin [13, 18-20], while SP-D modulates conversion of LA to SA surfactant [11, 21]. Recent findings in inducible SP-D transgenic mice showed that SP-D preferentially interacts with newly synthesized LA surfactant enriched in phosphatidyl inositol, an action that helps maintain surfactant pool size in neonatal and adult lung through conversion of LA surfactant into catabolically active SA forms [11]. In this study most of induced SP-D fractionated with SA surfactant [11]. Earlier studies identified phosphatidyl inositol as the major phospholipid binding site for SP-D in lamellar body enriched surfactant

[22]. SP-B and SP-C facilitate the formation and stability of interfacial films and surface active monolayer at the air-water interface [5]. Among surfactant components, SP-A and SP-D have been extensively characterized as host defense components of pulmonary surfactant [3, 23]. However, increasing evidence indicates that all surfactant proteins and one or more surfactant phospholipids regulate immunological homeostasis, inflammation, and innate lung host defense [4, 24].

Pulmonary surfactant in lung disease

Abnormal surfactant levels and composition in humans have been associated with respiratory dysfunction and inflammation in ALI/ARDS [25, 26], pulmonary fibrosis [27], emphysema [9, 28], cystic fibrosis [29], COPD [30], and RDS in newborns [1]. Genetic variation or deletion in surfactant protein genes is associated with COPD [31, 32], interstitial lung diseases [33, 34], cancer [35, 36], pulmonary infections [37, 38], congenital alveolar proteinosis [14, 39, 40], and enhanced development of broncho-pulmonary dysplasia [41]. In parallel, studies in SP-A, SP-B, SP-C, and SP-D-deficient mice support the crucial roles for surfactant proteins in surfactant stability and monolayer formation [13, 19, 21, 28, 42], metabolism [10, 21, 43, 44], pathogenesis of acute and chronic inflammation [28, 45-50], respiratory distress [13, 51, 52], lung injury [51, 53-55], and susceptibility to infection [46, 48, 56-58]. Deficiencies in surfactant protein components is associated with dys-regulated inflammatory responses [28, 43, 48, 49, 56] and presence of abnormal surfactant vesicles [26, 27]. Collectively, the results from genetically manipulated mice support the concept that genetic polymorphisms in one or more surfactant protein genes enhance susceptibility to pulmonary inflammatory disease in humans.

Genetic susceptibility may alter interactions between surfactant and alveolar cells promoting the development of lung disease. An inducible mouse model of SP-B expression was used to demonstrate that re-expression of SP-B in the lungs of SP-B-deficient mice suppressed inflammation caused by surfactant dysfunction [49]. SP-B is critical for the biophysical action of surfactant that lowers surface tension at the air-liquid interface; its absence, and by extension impaired surfactant function, resulted in activation of inflammatory pathways in alveolar macrophages and alveolar type II epithelial cells. Notable in this case is the activation of the transcription factor STAT3, which mediates the inflammatory as well anti-

inflammatory pathways of IL-6 family and IL-10 receptors in macrophages [59], respectively. In addition, recent studies demonstrated that STAT3 expression in alveolar type II epithelial cells is critical for pulmonary homeostasis; STAT3 regulates surfactant lipid synthesis and secretion [60, 61], oxidant stress [62, 63], and it protects epithelial cells from acute lung injury by LPS or adenoviral infections [60, 62].

A combination of genetic and environmental modifications of surfactant may further enhance the development of pulmonary disorders. In this regard, oxidative stress defined as the loss of balance between oxidant and antioxidant pathways underlies the pathogenesis of several inflammatory lung diseases including ARDS, COPD, and fibrosis [64]. Air pollution and ozone in particular, infections, and smoking are among environmental exposures that contribute to pulmonary oxidant stress [65]. Oxidized surfactant displays inferior anti-inflammatory and surface tension lowering properties [66, 67]. Reactive oxygen species alter the structure of surfactant lipids and proteins inactivating biophysical and immunological properties of surfactant components [68-76]. The macrophage scavenger receptors SR-A and MARCO clear oxidized surfactant lipids as a source of inflammation in the alveolus [77]. Under conditions of oxidative stress, SP-A, the major protein component of surfactant, plays a protective role against oxidative modification and in preserving the surface tension lowering properties of surfactant lipids [66, 78]. Genetic polymorphisms in SP-A may influence oxidant induced dysfunction of surfactant.

Decreased catabolism of surface-inactive surfactant aggregate forms is associated with respiratory dysfunction in patients with asthma, cystic fibrosis, and pneumonia [79, 80]. Impaired surfactant catabolism by alveolar macrophages results in accumulation of surfactant and eventual respiratory failure in patients with primary or secondary alveolar proteinosis (PAP) [81-84]. Loss of GM-CSF function is the main cause of primary PAP. GM-CSF is critical for the local differentiation of alveolar macrophages and their ability to degrade surfactant lipids and proteins [82, 85]. This discovery revealed the importance of alveolar macrophages in surfactant catabolism. In mice, alveolar macrophages contribute about 50% of surfactant catabolism [86]. Correspondingly, temporary depletion of alveolar macrophages results in as 8-10 fold increase in surfactant pool size [87] in rats. A variety of conditions including immunodeficiency syndromes, chronic infection, hematologic malignancies, and environmental exposures contribute to the

development of secondary PAP [84]. It is not clear how conditions that lead to secondary PAP impair alveolar macrophage function. However, secondary PAP in *Pneumocystis carinii* pneumonia involves stimulation of surfactant protein synthesis through the interaction between epithelial cell CD40 and CD154 on immune cells [88]. Moreover, accumulation of surfactant in this model is associated with re-distribution of SP-A and SP-D from large aggregate to small aggregate surfactant, oxidative de-oligomerization of SP-D, and alterations in relative expression levels of SP-B and SP-C [89-91]. Structural modifications and redistribution of surfactant proteins are known to occur in primary PAP surfactant as well [92, 93]. Congenital PAP, a rare group of genetic disorders caused by mutations in SP-B and SP-C genes [14, 39, 40], is characterized by mis-processing of SP-B and SP-C pro-proteins and secretion of metabolically inept surfactant containing abnormal forms of SP-B and SP-C [94, 95]. These studies emphasize that alveolar macrophage dysfunction on one hand, and endogenous or exogenous biochemical disruption of surfactant on the other, disturb the interaction of metabolic and immune processes signifying the development of lung disease.

SP-A and SP-D

Structural characterizations of SP-A and SP-D have led to the recognition that SP-A and SP-D interface critical physiological and immune functions of pulmonary surfactant. Both SP-A and SP-D are members of the collectin family of proteins, a classification that is based on homologous amino-terminal collagen-like domains and carboxy-terminal carbohydrate recognition domains (CRD). Mannose binding lectin (MBL), the founding member of the collectin family, is the first component of the lectin pathway of complement in the circulation [96]. Collectins are a subgroup of a large superfamily of proteins containing Ca^{++} -dependent CRD domains. However, the CRD domains of SP-A and SP-D are adapted to interact with distinct surfactant phospholipids [97, 98]. The collagen-like domain of collectins forms trimers and disulfide-dependent oligomerization of trimers then results in deca-octameric SP-A or MBL structures resembling a bouquet of flowers, or cross-hatched SP-D dodecamers. Thus formed, collectins serve as humoral pattern recognition molecules where multiple CRD domains bind, in a calcium-dependent manner, carbohydrate-based ligands on the surface of diverse microorganisms [96]. On the other hand, SP-A and SP-D serve discrete but

complementary roles in intra-alveolar surfactant dynamics and modulation of surfactant turnover by alveolar macrophages and type II epithelial cells, thus contributing to physiological and immunological integrity of the distal airspace. Swapping of amino-terminal segments including the collagen-like domains between SP-A and SP-D is not sufficient to interchange SP-A with SP-D function *in vivo* indicating that SP-A and SP-D are not physiologically redundant. SP-A binds and aggregates phosphatidyl choline, an interaction that facilitates formation of tubular myelin in large aggregate surfactant [99, 100]. Binding of SP-D to phosphatidyl inositol facilitates conversion of large aggregate surfactant to smaller lamellar forms that are metabolized by alveolar type II epithelial cells [18]. Proper surfactant ultrastructure appears critical for the post-natal SP-D-mediated surfactant maturation and the turnover of surfactant lipids by alveolar type II epithelial cells [10, 18, 19, 21, 43, 44, 99, 101-106].

Both SP-A and SP-D serve a pivotal function in lung humoral and innate immunity; they bind a wide range of pathogens, suppress microbial growth, and damage bacterial membranes on one hand, and modulate alveolar macrophage phagocytosis of monodisperse or agglutinated microorganisms [3, 23, 107-111] on the other. In addition to binding carbohydrates, SP-A binds lipid and protein moieties on microbial cell wall molecules. Thus, SP-A binds the lipid A moiety of rough lipopolysaccharides [112], disaturated phosphatidyl glycerol on the surface of *Mycoplasma pneumoniae* [109], the 65 kDa surface cytotoxin MPN372 on *Mycoplasma pneumoniae* [113, 114], and the adhesins Apa and Eap on *Mycobacterium tuberculosis* [115] and *Staphylococcus aureus* [116], respectively. SP-D interacts with heptose in the inner core oligosaccharide of rough LPS and mannose in O-antigen carbohydrate chains of smooth LPS [117, 118]. Extraction of LPS from the cell wall of gram-negative bacteria underlies the direct bactericidal activity of SP-A and SP-D [108]. LPS is a major immunoreactive contaminant in inhaled air that can disrupt surfactant membranes. Both SP-A and SP-D are involved in clearance and detoxification of LPS by alveolar macrophages [119, 120]. Further, SP-A- and SP-D-deficient mice are unable to resolve neutrophilic inflammation caused by gram-negative bacteria or LPS [43, 58]. Clathrin-dependent endocytosis of SP-A activates a PKC ζ -dependent pathway that blocks the ability of LPS to induce inflammation in alveolar macrophages [121]. Concordant to their ability to remove LPS from the surface of bacteria, SP-A and SP-D remove LPS from the membrane of surfactant vesicles [122, 123] as well,

and also bind and aggregate LPS in model surfactant monolayer membranes [124, 125]. These findings support the notion that SP-A and SP-D preserve basal surfactant function through their ability to both sterilize pathogens and remove inflammatory molecules that are inhaled into airway secretions.

In different circumstances, SP-A and SP-D enhance pathogen-dependent activation of pro-inflammatory responses of alveolar macrophages during ingestion of SP-A- or SP-D-opsonized microbes *in vitro* [126-131], a proactive clearance mechanism that also operates *in vivo* during initial clearance of experimental pulmonary infections [45, 46, 58, 90, 129, 132-137]. In contrast, SP-A and SP-D suppress secretion of pro-inflammatory cytokines and oxidant intermediates when macrophages are challenged with pathogen-derived cell wall components such as lipopolysaccharide [121, 138-144] or mycobacterial cell wall extract [145]; this protective SP-A function has been noted in both *in vitro* macrophage culture and *in vivo* animal models [58, 140, 146]. SP-A initially enhances clearance of *Mycoplasma pneumoniae* [133, 134] but subsequently turns off inflammation [146], indicating that SP-A works in a temporal fashion to resolve inflammation. In this context, SP-A and SP-D regulate phagocytosis of apoptotic cells [147, 148]. In addition, SP-A enhances secretion of anti-inflammatory IL-10 and can induce secretion of TGF β in the presence of mycobacterial cell wall antigen or apoptotic cells in peripheral blood monocytes and alveolar macrophages [145, 149], processes that may help resolve inflammation and restore steady-state concentrations of pulmonary surfactant in the long-term. In this regard, TGF β activation by epithelial β 6 integrin is necessary for the maintenance of normal surfactant pool size; lack of β 6 integrin results in inflammation and development of PAP [150]. These findings indicate that precise coordination of surfactant metabolism with immune responses is necessary to resolve inflammation and restore airway homeostasis.

SP-B, SP-C, and surfactant lipids

The hydrophobic SP-B and SP-C have been characterized extensively for their ability to effect lamellar body formation, secretion, and formation of the surfactant monolayer that is critical to the lowering of surface tension at the air/water interface [5]. However, increasing evidence indicate that SP-B and SP-C are also involved in immunomodulation that is critical for the stability and host defense of the airways [24]. Inducible transgenic

mice were used to show that reduction in SP-B expression and the associated abnormalities in reducing surface tension divulged an inflammatory response in alveolar macrophages and type II epithelial cells [49]. SP-B-deficiency also impaired the ability of the lung to counteract LPS-induced inflammation [50]. SP-B is the most critical effector of the surface tension lowering properties of surfactant, a function that also influences the shape and phagocytic activity of alveolar macrophages [151].

More recently, SP-C-deficient mice were found to be susceptible to bacterial and viral infections and to suffer from excessive inflammation [48, 56]. Bleomycin-induced fibrosis was more severe in SP-C-deficient mice [152]. The role of SP-C in suppressing inflammation is in part related to the ability of its amino-terminal segment to bind LPS [153-155]. On the other hand, SP-C-containing surfactant vesicles bind and interfere with toll-like receptor mediated inflammatory responses in macrophages [56]. Exogenous surfactant and surfactant lipids in the absence of proteins exhibit anti-inflammatory properties as well [156-158]. SP-B and SP-C appear to confer differential roles in the ability of surfactant to suppress inflammation; the presence of SP-B rather than SP-C in surfactant was more important in suppressing LPS-induced production of nitric oxide by alveolar macrophages [158]. Surfactant is a bovine derived therapeutic surfactant lipoprotein that is used to treat respiratory distress syndrome in premature infants; it contains SP-B and SP-C but not SP-A or SP-D [159]. In addition, surfactant vesicles containing phosphatidyl glycerol (POPG) block LPS-induced inflammation by macrophages [160]. POPG is an acidic phospholipid that is uniquely present in pulmonary surfactant. It was shown that POPG binds CD14, TLR4, and MD-2, thus antagonizing the ability of LPS to induce inflammation [160]. On the other hand, surfactant lipids and dipalmitoyl phosphatidyl choline in particular block epithelial cell expressed TLR4 activation by limiting translocation of TLR4 to membrane lipid raft micro domains [161]. Phospholipid vesicles containing dipalmitoyl phosphatidyl choline, the major phospholipid in pulmonary surfactant, induced expression of several macrophage innate immune receptors [162], a property that has also been noted for purified SP-A and SP-D [163-168]. In these cases, SP-D influences alveolar macrophage differentiation as indicated by expression of CD11c [169-171], while surfactant DPPC and SP-A induced expression and function of several innate immune receptors that so far include the complement receptor CR3 (CD11b), the scavenger receptors SR-A, CD36, and

LOX-1, the mannose receptor, toll-like receptors, Fc receptors and the complement receptor CR1 [157, 164, 169, 170]. Adoptive transfer experiments and studies in SP-D-inducible transgenic mice indicate that SP-D contributes to the local differentiation of freshly recruited monocytes in the alveolar space [169-171]. Surfactant lipids and proteins may subsequently become ligands for one or more of these receptors. It is also notable that phosphatidyl glycerol facilitates clearance of surfactant phospholipids by alveolar macrophages [172, 173]. These studies indicate that alveolar macrophages and alveolar type II epithelial cells coordinate surfactant metabolism and innate host defense properties of pulmonary surfactant.

SP-A and SP-D receptors

Several receptor systems have been identified that mediate homeostasis and immunomodulatory activities of SP-A and SP-D by alveolar macrophages. These include CD91/calreticulin, SIRP α , toll-like receptors, and SP-R210.

CD91/calreticulin and SIRP α

Earlier studies reported that calreticulin, an endoplasmic reticulum chaperone, appears on the cell surface of normal and malignant cells [174, 175]. In parallel, other studies identified the collagen-binding C1q receptor C1qR as calreticulin and additional studies demonstrated that the collagen-like domains of collectins (SP-A, SP-D, and MBL) also bind calreticulin [176] in macrophages. CD91, also known as lipoprotein related receptor 1 [177], was then identified as a receptor for secreted chaperones including calreticulin [178]. Gardai et al [179, 180] have subsequently demonstrated that calreticulin bridges SP-A-coated apoptotic cells and bacteria to CD91, resulting in the ingestion of SP-A-bound cargo, and a concomitant pro-inflammatory response. The same studies also showed that the CRD domains of collectins bind the signal inhibitory receptor SIRP α , thus mediating anti-inflammatory activities of collectins [179]. SIRP α was initially defined as a counter-receptor for the widely expressed CD47 [181], that transmits “eat me” vs. “don’t eat me” signals to initiate phagocytosis of apoptotic cells and exogenous intruders but not intact host cells by macrophages. The CD47-SIRP α interaction inhibits “self” engulfment through de-phosphorylation of myosin IIA [182]. However, SP-A and SP-D also bind SIRP α to suppress ingestion of apoptotic cells by alveolar

macrophages [148]; this interaction is thought to maintain a low rate of alveolar macrophage phagocytosis in the resting state without inciting inflammation. In this model, the interaction of lung collectins with apoptotic cells or bacteria increases the affinity of SP-A or SP-D with calreticulin, which favors the inflammatory/clearance CD91/calreticulin pathway over the basal inhibitory SIRP α pathway. On the other hand, LPS-mediated suppression of SIRP α [183] correlates with the enhanced ability of exudate macrophages to ingest apoptotic cells [148]. However, recent studies in transgenic mice and CD91-deficient cells showed that CD91 is not the only cell-surface receptor for calreticulin [184, 185]. Other studies identified the class A scavenger receptor SR-A as a calreticulin receptor on antigen presenting cells [186]. Interestingly, the pairing of CD47 with SIRP α inhibits phagocytosis via IgG Fc or complement receptors [187], but does not affect the phagocytosis of senescent or oxidized erythrocytes via scavenger receptors [188]. Therefore, SP-A and SP-D engagement of SIRP α may suppress phagocytosis via some but not all innate immune receptors. On the other hand, more recent studies demonstrated that CD91/calreticulin acts as a receptor for modified SP-D [76]. S-nitrosylation of thiols in the SP-D collagen domain enhanced the interaction of SP-D with CD91/calreticulin in the absence of apoptotic cell or bacterial targets. The increased binding of S-nitrosylated SP-D with calreticulin was associated with reduction in the oligomerization state of nitrosylated SP-D. In this case, postranslational modification of SP-D and its binding to calreticulin may overcome SIRP α -mediated local immunosuppression by SP-D. This alternative engagement of the CD91/calreticulin receptor has pathophysiological significance in that SP-D is subject to modification by reactive nitrogen and oxygen radicals during inflammation *in vivo* [75]. Modified SP-D exacerbates inflammation and injury through recruitment of inflammatory cells during *Pneumocystis carinii* pneumonia [89].

CD14 and toll-like receptors

Several studies have demonstrated that both SP-A and SP-D antagonize the inflammatory actions of pathogen-derived molecules through their ability to directly bind the LPS receptor CD14 and the toll-like receptors TLR2 and TLR4 [189-193], and the TLR4 adaptor MD-2 [194]. SP-A and SP-D were shown to bind these innate receptors via their CRD domains. SP-A and SP-D binding blocks the interaction of pathogen immuno-stimulatory molecules with TLR receptor components. On the other

hand, SP-A can also modulate expression and function of TLRs in human alveolar macrophages indirectly [165]. As mentioned above, all surfactant proteins and lipids modulate, directly or indirectly, the activity of toll-like receptors in the airway.

SP-R210

The 210 kDa SP-A receptor was identified in rat lung and human U937 cell membranes [195]. Polyclonal antibodies to SP-R210 blocked SP-A binding to alveolar macrophages and type II epithelial cells and also blocked the ability of SP-A to inhibit surfactant phospholipid secretion in alveolar type II epithelial cells. These studies provided initial evidence that SP-R210 is involved in surfactant metabolism. Subsequent studies defined the role of SP-R210 in immunological responses [126, 127, 145, 196]. SP-R210 facilitates phagocytosis and killing of SP-A-opsonized *Mycobacterium bovis* BCG, by a mechanism involving induction of nitric oxide and secretion of TNF α [126, 127]. Interestingly, SP-R210 was shown to inhibit T lymphocyte proliferation [196] in peripheral blood mononuclear cells, indicating that SP-R210 can also act to suppress inflammation. Antibodies against the SP-A binding domain confirmed the ability of SP-R210 to inhibit Th1 lymphocyte proliferation in the context of an immune response to mycobacterial antigen [145]. The suppression of lymphocyte proliferation by SP-R210 was mediated via induction of the anti-inflammatory cytokines IL-10 and TGF β . A molecule with properties similar to SP-R210 was recently identified in myometrial membranes where it mediates the anti-inflammatory response of SP-A during late gestation [197]. LPS and mycobacterial cell wall antigen stimulate inflammatory responses via CD14 and toll-like receptors [190, 193, 198]. In this regard, ligation of SP-R210 elicited secretion of IL-10 and suppressed, but did not abolish mycobacterial antigen-stimulated secretion of TNF α [145]. Thus, it appears that SP-A, via SP-R210, enhances macrophage activation inducing ingestion and killing of SP-A-opsonized pathogens but subsequently, it works to suppress inflammation.

Macrophage expressed SP-R210 was recently identified as a short trans-membrane isoform of the novel myosin 18A (Myo18A) [199] that was named as SP-R210_s. The Myo18A gene was also identified by Obinata and colleagues in fibroblasts [200]. Subsequent independent studies showed that the *Myo18A* gene encodes two major splice variants of Myo18A that are differentially expressed in somatic and immune cells [199, 201]. With the exception of alveolar type II epithelial cells

[195, 199], non-immune cells and tissues express a long intracellular isoform with an amino-terminal PDZ domain [199, 201] that was also named SP-R210_L. The long SP-R210_L isoform regulates retrograde membrane flow; this function involves formation of a tripartite complex that bridges the PDZ domain of SP-R210_L with LRAP35a and myosin IIA [202]. The size of SP-R210_L mRNA varies from 7.5-8.5 kbases in different tissues. SP-R210_S does not contain the amino-terminal PDZ domain; it is expressed as a type II membrane protein on the surface of macrophages where it acts as an SP-A receptor. The size of the SP-R210_S mRNA ranges between 6.5-7.0 kbases in different macrophage populations and cell lines. Macrophages also express a third 150-170 kDa splice variant called SP-R210_{S1} [145] expressed from 6.0-6.5 kbase mRNA species. SP-R210_S is the predominant isoform expressed on alveolar macrophages while SP-R210_{S1} is expressed in monocytes. The role of macrophage SP-R210 isoforms in surfactant catabolism is inferred from results in GM-CSF-deficient alveolar macrophages. Alveolar macrophages from GM-CSF-deficient mice lack high affinity binding sites for SP-A, and exhibit decreased uptake and degradation of LA surfactant [82].

Conclusions

Tightly controlled mechanisms exist to preserve surfactant homeostasis and eradicate a myriad of airborne pathogens to keep the lung sterile. This complex mixture of lipids and proteins lowers surface tension at the air-liquid interface preventing collapse of alveolar sacs during exhalation on one hand and intimately involved in innate host defense of the distal airways on the other. A host of studies in humans, animal models, and in cultured cells indicate that the integrity of surfactant and its various biophysical forms are critical components of the lung innate immune system. Studies on partial or individual surfactant components have revealed various mechanisms by which surfactant and its components interact with alveolar macrophages to prevent infection and precisely regulate inflammatory responses. The ability of several surfactant components to modulate innate immune receptors on alveolar macrophages, which are variously involved in phagocytosis of microbes (scavenger, complement, and mannose receptors), and uptake of damaged proteins and lipids (scavenger receptors) may help eliminate infectious agents and immunoreactive macromolecules from airway secretions.

It appears that considerable effort is invested to suppress the inflammatory actions of toll-like receptors in alveolar epithelial cells and macrophages as several surfactant proteins and a combination of surfactant lipids and proteins bind and antagonize the effects of LPS and infectious organisms. Within this realm, recent studies showed that TLR4 is an important immunological source of acute lung injury caused by acid inspiration, influenza, pulmonary contusion, and hemorrhagic shock [203-205]; these injury models did not involve administration of LPS indicating that insults that activate TLR inflammatory pathways are not restricted to specific pathogens or infection. The domain-dependent interactions of SP-A and SP-D with CD91/calreticulin and SIRP α control opposing macrophage reactions that maintain basal responses but enhance the clearance of apoptotic cells under inflammatory conditions that suppress SIRP α . The CD91/calreticulin pathway may contribute to oxidant-induced lung injury through recognition of nitrosylated SP-D. SP-R210 appears to coordinate macrophage activation with ingestion of SP-A-opsonized mycobacteria but to suppress antigen-mediated inflammation. Differential expression of SP-R210 isoforms in resident and recruited macrophages may mediate opposing inflammatory activities of SP-A during resolution of inflammation. SP-R210 isoforms, via secretion of IL-10, TGF β , and modulation of surfactant secretion may have reciprocal roles in surfactant metabolism by alveolar type II epithelial cells and macrophages.

Abbreviations

ALI (acute lung injury); ARDS (adult respiratory distress syndrome); COPD (chronic obstructive pulmonary disease); RDS (respiratory distress syndrome); PAP (pulmonary alveolar proteinosis); LA (large aggregate); SA (small aggregate); CRD (carbohydrate recognition domain); GM-CSF (granulocyte macrophage colony stimulating factor); SP (surfactant protein); TLR (toll-like receptors).

Acknowledgements

We acknowledge the financial support of the Juvenile Diabetes Research Foundation, Potts Memorial Foundation, University of Texas Health Science Center President's Council grants, and NIH/NHLBI Public Health grant HL068127.

References

- 1 Halliday HL: Surfactants: Past, present and future. *J Perinatol* 2008;28 Suppl 1:S47-56.
- 2 Lhert F, Yan W, Biswas SC, Hall SB: Effects of hydrophobic surfactant proteins on collapse of pulmonary surfactant monolayers. *Biophys J* 2007;93:4237-4243.
- 3 Wright JR: Immunoregulatory functions of surfactant proteins. *Nat Rev Immunol* 2005;5:58-68.
- 4 Christmann U, Buechner-Maxwell VA, Witonsky SG, Hite RD: Role of lung surfactant in respiratory disease: Current knowledge in large animal medicine. *J Vet Intern Med* 2009;23:227-242.
- 5 Perez-Gil J: Structure of pulmonary surfactant membranes and films: The role of proteins and lipid-protein interactions. *Biochim Biophys Acta* 2008;1778:1676-1695.
- 6 Rooney SA, Young SL, Mendelson CR: Molecular and cellular processing of lung surfactant. *FASEB J* 1994;8:957-967.
- 7 Rooney SA: Lung surfactant. *Environ Health Perspect* 1984;55:205-226.
- 8 Horowitz AD, Moussavian B, Whitsett JA: Roles of sp-a, sp-b, and sp-c in modulation of lipid uptake by pulmonary epithelial cells in vitro. *Am J Physiol* 1996;270:L69-79.
- 9 Botas C, Poulain F, Akiyama J, Brown C, Allen L, Goerke J, Clements J, Carlson E, Gillespie AM, Epstein C, Hawgood S: Altered surfactant homeostasis and alveolar type ii cell morphology in mice lacking surfactant protein d. *Proc Natl Acad Sci U S A* 1998;95:11869-11874.
- 10 Bates SR, Dodia C, Tao JQ, Fisher AB: Surfactant protein-a plays an important role in lung surfactant clearance: Evidence using the surfactant protein-a gene-targeted mouse. *Am J Physiol Lung Cell Mol Physiol* 2008;294:L325-333.
- 11 Ikegami M, Grant S, Korfhagen T, Scheule RK, Whitsett JA: Surfactant protein-d regulates the postnatal maturation of pulmonary surfactant lipid pool sizes. *J Appl Physiol* 2009;106:1545-1552.
- 12 Tokieda K, Whitsett JA, Clark JC, Weaver TE, Ikeda K, McConnell KB, Jobe AH, Ikegami M, Iwamoto HS: Pulmonary dysfunction in neonatal sp-b-deficient mice. *Am J Physiol* 1997;273:L875-882.
- 13 Clark JC, Wert SE, Bachurski CJ, Stahlman MT, Stripp BR, Weaver TE, Whitsett JA: Targeted disruption of the surfactant protein b gene disrupts surfactant homeostasis, causing respiratory failure in newborn mice. *Proc Natl Acad Sci U S A* 1995;92:7794-7798.
- 14 Noguee LM, Garnier G, Dietz HC, Singer L, Murphy AM, deMello DE, Colten HR: A mutation in the surfactant protein b gene responsible for fatal neonatal respiratory disease in multiple kindreds. *J Clin Invest* 1994;93:1860-1863.
- 15 Schuerman FA, Griesse M, Gille JP, Brasch F, Noorduyt LA, van Kaam AH: Surfactant protein b deficiency caused by a novel mutation involving multiple exons of the sp-b gene. *Eur J Med Res* 2008;13:281-286.
- 16 Wright JR, Clements JA: Metabolism and turnover of lung surfactant. *Am Rev Respir Dis* 1987;136:426-444.
- 17 Baritussio A, Alberti A, Quaglini D, Pettenazzo A, Dalzoppo D, Sartori L, Pasquali-Ronchetti I: Sp-a, sp-b, and sp-c in surfactant subtypes around birth: Reexamination of alveolar life cycle of surfactant. *Am J Physiol* 1994;266:L436-447.
- 18 Yao LJ, Fraher LJ, Veldhuizen RA, Samuelson S, Borron P, Malloy J, McCaig L, Lewis JF: Effects of alveolar surfactant aggregates on T-lymphocyte proliferation. *Biochim Biophys Acta* 2001;1535(3):266-74.
- 19 Korfhagen TR, Bruno MD, Ross GF, Huelsman KM, Ikegami M, Jobe AH, Wert SE, Stripp BR, Morris RE, Glasser SW, Bachurski CJ, Iwamoto HS, Whitsett JA: Altered surfactant function and structure in sp-a gene targeted mice. *Proc Natl Acad Sci U S A* 1996;93:9594-9599.
- 20 Veldhuizen RA, Yao LJ, Hearn SA, Possmayer F, Lewis JF: Surfactant-associated protein a is important for maintaining surfactant large-aggregate forms during surface-area cycling. *Biochem J* 1996;313:835-840.
- 21 Ikegami M, Na CL, Korfhagen TR, Whitsett JA: Surfactant protein d influences surfactant ultrastructure and uptake by alveolar type ii cells. *Am J Physiol Lung Cell Mol Physiol* 2005;288:L552-561.
- 22 Persson AV, Gibbons BJ, Shoemaker JD, Moxley MA, Longmore WJ: The major glycolipid recognized by sp-d in surfactant is phosphatidylinositol. *Biochemistry* 1992;31:12183-12189.
- 23 Kuroki Y, Takahashi M, Nishitani C: Pulmonary collectins in innate immunity of the lung. *Cell Microbiol* 2007;9(8):1871-9.
- 24 Mulugeta S, Beers MF: Surfactant protein c: Its unique properties and emerging immunomodulatory role in the lung. *Microbes Infect* 2006;8:2317-2323.
- 25 Lewis JF, Veldhuizen R: The role of exogenous surfactant in the treatment of acute lung injury. *Annu Rev Physiol* 2003;65:613-642.
- 26 Todd DA, Marsh MJ, George A, Henderson NG, Barr H, Sebastian S, Clark GT, Koster G, Clark HW, Postle AD: Surfactant phospholipids, surfactant proteins, and inflammatory markers during acute lung injury in children. *Pediatr Crit Care Med* 2009, in press.
- 27 Gunther A, Schmidt R, Nix F, Yabut-Perez M, Guth C, Rosseau S, Siebert C, Grimminger F, Morr H, Velcovsky HG, Seeger W: Surfactant abnormalities in idiopathic pulmonary fibrosis, hypersensitivity pneumonitis and sarcoidosis. *Eur Respir J* 1999;14:565-573.
- 28 Wert SE, Yoshida M, LeVine AM, Ikegami M, Jones T, Ross GF, Fisher JH, Korfhagen TR, Whitsett JA: Increased metalloproteinase activity, oxidant production, and emphysema in surfactant protein d gene-inactivated mice. *Proc Natl Acad Sci U S A* 2000;97:5972-5977.
- 29 Griesse M, Essl R, Schmidt R, Rietschel E, Ratjen F, Ballmann M, Paul K: Pulmonary surfactant, lung function, and endobronchial inflammation in cystic fibrosis. *Am J Respir Crit Care Med* 2004;170:1000-1005.
- 30 Ohlmeier S, Vuolanto M, Toljamo T, Vuopala K, Salmenkivi K, Myllarniemi M, Kinnula VL: Proteomics of human lung tissue identifies surfactant protein a as a marker of chronic obstructive pulmonary disease. *J Proteome Res* 2008;7:5125-5132.
- 31 Foreman MG, DeMeo DL, Hersh CP, Carey VJ, Fan VS, Reilly JJ, Shapiro SD, Silverman EK: Polymorphic variation in surfactant protein b is associated with copd exacerbations. *Eur Respir J* 2008;32:938-944.
- 32 Guo X, Lin HM, Lin Z, Montano M, Sansores R, Wang G, DiAngelo S, Pardo A, Selman M, Floros J: Surfactant protein gene a, b, and d marker alleles in chronic obstructive pulmonary disease of a mexican population. *Eur Respir J* 2001;18:482-490.
- 33 Guillot L, Epaul R, Thouvenin G, Jonard L, Mohsni A, Couderc R, Counil F, de Blic J, Taam RA, Le Bourgeois M, Reix P, Flamein F, Clement A, Feldmann D: New surfactant protein c gene mutations associated with diffuse lung disease. *J Med Genet* 2009;46:490-494.
- 34 Brasch F, Griesse M, Tredano M, Johnen G, Ochs M, Rieger C, Mulugeta S, Muller KM, Bahau M, Beers MF: Interstitial lung disease in a baby with a de novo mutation in the sftpc gene. *Eur Respir J* 2004;24:30-39.

- 35 Jiang F, Caraway NP, Nebiyou Bekele B, Zhang HZ, Khanna A, Wang H, Li R, Fernandez RL, Zaidi TM, Johnston DA, Katz RL: Surfactant protein a gene deletion and prognostics for patients with stage i non-small cell lung cancer. *Clin Cancer Res* 2005;11:5417-5424.
- 36 Wang Y, Kuan PJ, Xing C, Cronkhite JT, Torres F, Rosenblatt RL, DiMaio JM, Kinch LN, Grishin NV, Garcia CK: Genetic defects in surfactant protein a2 are associated with pulmonary fibrosis and lung cancer. *Am J Hum Genet* 2009;84:52-59.
- 37 Puthothu B, Forster J, Heinze J, Heinzmann A, Krueger M: Surfactant protein b polymorphisms are associated with severe respiratory syncytial virus infection, but not with asthma. *BMC Pulm Med* 2007;7:6.
- 38 Hallman M, Haataja R: Surfactant protein polymorphisms and neonatal lung disease. *Semin Perinatol* 2006;30:350-361.
- 39 Lin Z, deMello DE, Wallot M, Floros J: An sp-b gene mutation responsible for sp-b deficiency in fatal congenital alveolar proteinosis: Evidence for a mutation hotspot in exon 4. *Mol Genet Metab* 1998;64:25-35.
- 40 Tredano M, Griese M, Brasch F, Schumacher S, de Blic J, Marque S, Houdayer C, Elion J, Couderc R, Bahuau M: Mutation of sftpc in infantile pulmonary alveolar proteinosis with or without fibrosing lung disease. *Am J Med Genet A* 2004;126A:18-26.
- 41 Pavlovic J, Papagaroufalis C, Xanthou M, Liu W, Fan R, Thomas NJ, Apostolidou I, Papathoma E, Megaloyianni E, DiAngelo S, Floros J: Genetic variants of surfactant proteins a, b, c, and d in bronchopulmonary dysplasia. *Dis Markers* 2006;22:277-291.
- 42 Glasser SW, Burhans MS, Korfhagen TR, Na CL, Sly PD, Ross GF, Ikegami M, Whitsett JA: Altered stability of pulmonary surfactant in sp-c-deficient mice. *Proc Natl Acad Sci U S A* 2001;98:6366-6371.
- 43 Ikegami M, Scoville EA, Grant S, Korfhagen T, Brondyk W, Scheule RK, Whitsett JA: Surfactant protein-d and surfactant inhibit endotoxin-induced pulmonary inflammation. *Chest* 2007;132:1447-1454.
- 44 Korfhagen TR, Sheftelyevich V, Burhans MS, Bruno MD, Ross GF, Wert SE, Stahlman MT, Jobe AH, Ikegami M, Whitsett JA, Fisher JH: Surfactant protein-d regulates surfactant phospholipid homeostasis in vivo. *J Biol Chem* 1998;273:28438-28443.
- 45 LeVine AM, Hartshorn K, Elliott J, Whitsett J, Korfhagen T: Absence of sp-a modulates innate and adaptive defense responses to pulmonary influenza infection. *Am J Physiol Lung Cell Mol Physiol* 2002;282:L563-572.
- 46 LeVine AM, Bruno MD, Huelsman KM, Ross GF, Whitsett JA, Korfhagen TR: Surfactant protein a-deficient mice are susceptible to group b streptococcal infection. *J Immunol* 1997;158:4336-4340.
- 47 Glasser SW, Detmer EA, Ikegami M, Na CL, Stahlman MT, Whitsett JA: Pneumonitis and emphysema in sp-c gene targeted mice. *J Biol Chem* 2003;278:14291-14298.
- 48 Glasser SW, Senft AP, Whitsett JA, Maxfield MD, Ross GF, Richardson TR, Prows DR, Xu Y, Korfhagen TR: Macrophage dysfunction and susceptibility to pulmonary pseudomonas aeruginosa infection in surfactant protein c-deficient mice. *J Immunol* 2008;181:621-628.
- 49 Ikegami M, Whitsett JA, Martis PC, Weaver TE: Reversibility of lung inflammation caused by sp-b deficiency. *Am J Physiol Lung Cell Mol Physiol* 2005;289:L962-970.
- 50 Epaul R, Ikegami M, Whitsett JA, Jobe AH, Weaver TE, Akinbi HT: Surfactant protein b inhibits endotoxin-induced lung inflammation. *Am J Respir Cell Mol Biol* 2003;28:373-378.
- 51 Nessler LL, Melton KR, Ikegami M, Na CL, Wert SE, Rice WR, Whitsett JA, Weaver TE: Partial sp-b deficiency perturbs lung function and causes air space abnormalities. *Am J Physiol Lung Cell Mol Physiol* 2005;288:L1154-1161.
- 52 Melton KR, Nessler LL, Ikegami M, Tichelaar JW, Clark JC, Whitsett JA, Weaver TE: Sp-b deficiency causes respiratory failure in adult mice. *Am J Physiol Lung Cell Mol Physiol* 2003;285:L543-549.
- 53 Haddad IY, Milla C, Yang S, Panoskaltsis-Mortari A, Hawgood S, Lacey DL, Blazar BR: Surfactant protein a is a required mediator of keratinocyte growth factor after experimental marrow transplantation. *Am J Physiol Lung Cell Mol Physiol* 2003;285:L602-610.
- 54 Yang S, Milla C, Panoskaltsis-Mortari A, Hawgood S, Blazar BR, Haddad IY: Surfactant protein a decreases lung injury and mortality after murine marrow transplantation. *Am J Respir Cell Mol Biol* 2002;27:297-305.
- 55 Haque R, Umstead TM, Ponnuru P, Guo X, Hawgood S, Phelps DS, Floros J: Role of surfactant protein-a (sp-a) in lung injury in response to acute ozone exposure of sp-a deficient mice. *Toxicol Appl Pharmacol* 2007;220:72-82.
- 56 Glasser SW, Witt TL, Senft AP, Baatz JE, Folger D, Maxfield MD, Akinbi HT, Newton DA, Prows DR, Korfhagen TR: Surfactant protein c-deficient mice are susceptible to respiratory syncytial virus infection. *Am J Physiol Lung Cell Mol Physiol* 2009;297:L64-72.
- 57 LeVine AM, Gwozdz J, Stark J, Bruno M, Whitsett J, Korfhagen T: Surfactant protein-a enhances respiratory syncytial virus clearance in vivo. *J Clin Invest* 1999;103:1015-1021.
- 58 LeVine AM, Kurak KE, Bruno MD, Stark JM, Whitsett JA, Korfhagen TR: Surfactant protein-a-deficient mice are susceptible to pseudomonas aeruginosa infection. *Am J Respir Cell Mol Biol* 1998;19:700-708.
- 59 Maritano D, Sugrue ML, Tininini S, Dewilde S, Strobl B, Fu X, Murray-Tait V, Chiarle R, Poli V: The stat3 isoforms alpha and beta have unique and specific functions. *Nat Immunol* 2004;5:401-409.
- 60 Matsuzaki Y, Besnard V, Clark JC, Xu Y, Wert SE, Ikegami M, Whitsett JA: Stat3 regulates abca3 expression and influences lamellar body formation in alveolar type ii cells. *Am J Respir Cell Mol Biol* 2008;38:551-558.
- 61 Xu Y, Ikegami M, Wang Y, Matsuzaki Y, Whitsett JA: Gene expression and biological processes influenced by deletion of stat3 in pulmonary type ii epithelial cells. *BMC Genomics* 2007;8:455.
- 62 Matsuzaki Y, Xu Y, Ikegami M, Besnard V, Park KS, Hull WM, Wert SE, Whitsett JA: Stat3 is required for cytoprotection of the respiratory epithelium during adenoviral infection. *J Immunol* 2006;177:527-537.
- 63 Hokuto I, Ikegami M, Yoshida M, Takeda K, Akira S, Perl AK, Hull WM, Wert SE, Whitsett JA: Stat-3 is required for pulmonary homeostasis during hyperoxia. *J Clin Invest* 2004;113:28-37.
- 64 Cho HY, Kleeberger SR: Genetic mechanisms of susceptibility to oxidative lung injury in mice. *Free Radic Biol Med* 2007;42:433-445.
- 65 Park HS, Kim SR, Lee YC: Impact of oxidative stress on lung diseases. *Respirology* 2009;14:27-38.
- 66 Bailey TC, Maruscak AA, Petersen A, White S, Lewis JF, Veldhuizen RA: Physiological effects of oxidized exogenous surfactant in vivo: Effects of high tidal volume and surfactant protein a. *Am J Physiol Lung Cell Mol Physiol* 2006;291:L703-709.
- 67 Bailey TC, Da Silva KA, Lewis JF, Rodriguez-Capote K, Possmayer F, Veldhuizen RA: Physiological and inflammatory response to instillation of an oxidized surfactant in a rat model of surfactant deficiency. *J Appl Physiol* 2004;96:1674-1680.
- 68 Mikerov AN, Umstead TM, Gan X, Huang W, Guo X, Wang G, Phelps DS, Floros J: Impact of ozone exposure on the phagocytic activity of human surfactant protein a (sp-a) and sp-a variants. *Am J Physiol Lung Cell Mol Physiol* 2008;294:L121-130.

- 69 Manzanares D, Rodriguez-Capote K, Liu S, Haines T, Ramos Y, Zhao L, Doherty-Kirby A, Lajoie G, Possmayer F: Modification of tryptophan and methionine residues is implicated in the oxidative inactivation of surfactant protein b. *Biochemistry* 2007;46:5604-5615.
- 70 Rodriguez-Capote K, Manzanares D, Haines T, Possmayer F: Reactive oxygen species inactivation of surfactant involves structural and functional alterations to surfactant proteins sp-b and sp-c. *Biophys J* 2006;90:2808-2821.
- 71 Kuzmenko AI, Wu H, Wan S, McCormack FX: Surfactant protein a is a principal and oxidation-sensitive microbial permeabilizing factor in the alveolar lining fluid. *J Biol Chem* 2005;280:25913-25919.
- 72 Kuzmenko AI, Wu H, Bridges JP, McCormack FX: Surfactant protein oxidation: Surfactant lipid peroxidation damages surfactant protein a (sp-a) and inhibits interactions with phospholipid vesicles. *J Lipid Res* 2004;45:1061-1068.
- 73 Wang G, Bates-Kenney SR, Tao JQ, Phelps DS, Floros J: Differences in biochemical properties and in biological function between human sp-a1 and sp-a2 variants, and the impact of ozone-induced oxidation. *Biochemistry* 2004;43:4227-4239.
- 74 Davis IC, Zhu S, Sampson JB, Crow JP, Matalon S: Inhibition of human surfactant protein a function by oxidation intermediates of nitrite. *Free Radic Biol Med* 2002;33:1703-1713.
- 75 Matalon S, Shrestha K, Kirk M, Waldheuser S, McDonald B, Smith K, Gao Z, Belaouaj A, Crouch EC: Modification of surfactant protein d by reactive oxygen-nitrogen intermediates is accompanied by loss of aggregating activity, in vitro and in vivo. *FASEB J* 2009; 23:1415-1430.
- 76 Guo CJ, Atochina-Vasserman EN, Abramova E, Foley JP, Zaman A, Crouch E, Beers MF, Savani RC, Gow AJ: S-nitrosylation of surfactant protein-d controls inflammatory function. *PLoS Biol* 2008;6:e266.
- 77 Dahl M, Bauer AK, Arredouani M, Soininen R, Tryggvason K, Kleeberger SR, Kobzik L: Protection against inhaled oxidants through scavenging of oxidized lipids by macrophage receptors marco and sr-ai/ii. *J Clin Invest* 2007;117:757-764.
- 78 Rodriguez Capote K, McCormack FX, Possmayer F: Pulmonary surfactant protein-a (sp-a) restores the surface properties of surfactant after oxidation by a mechanism that requires the cys6 interchain disulfide bond and the phospholipid binding domain. *J Biol Chem* 2003;278:20461-20474.
- 79 Mander A, Langton-Hewer S, Bernhard W, Warner JO, Postle AD: Altered phospholipid composition and aggregate structure of lung surfactant is associated with impaired lung function in young children with respiratory infections. *Am J Respir Cell Mol Biol* 2002;27:714-721.
- 80 Beers MF, Hamvas A, Moxley MA, Gonzales LW, Guttentag SH, Solarin KO, Longmore WJ, Noguee LM, Ballard PL: Pulmonary surfactant metabolism in infants lacking surfactant protein b. *Am J Respir Cell Mol Biol* 2000;22:380-391.
- 81 Seymour JF, Presneill JJ: Pulmonary alveolar proteinosis: Progress in the first 44 years. *Am J Respir Crit Care Med* 2002;166:215-235.
- 82 Yoshida M, Ikegami M, Reed JA, Chroneos ZC, Whitsett JA: Gm-csf regulates protein and lipid catabolism by alveolar macrophages. *Am J Physiol Lung Cell Mol Physiol* 2001;280:L379-386.
- 83 Ikegami M, Ueda T, Hull W, Whitsett JA, Mulligan RC, Dranoff G, Jobe AH: Surfactant metabolism in transgenic mice after granulocyte macrophage-colony stimulating factor ablation. *Am J Physiol* 1996;270:L650-658.
- 84 Huizar I, Kavuru MS: Alveolar proteinosis syndrome: Pathogenesis, diagnosis, and management. *Curr Opin Pulm Med* 2009;15:491-498.
- 85 Shibata Y, Berclaz PY, Chroneos ZC, Yoshida M, Whitsett JA, Trapnell BC: Gm-csf regulates alveolar macrophage differentiation and innate immunity in the lung through pu.1. *Immunity* 2001;15:557-567.
- 86 Gurel O, Ikegami M, Chroneos ZC, Jobe AH: Macrophage and type ii cell catabolism of sp-a and saturated phosphatidylcholine in mouse lungs. *Am J Physiol Lung Cell Mol Physiol* 2001;280:L1266-1272.
- 87 Forbes A, Pickell M, Foroughian M, Yao LJ, Lewis J, Veldhuizen R: Alveolar macrophage depletion is associated with increased surfactant pool sizes in adult rats. *J Appl Physiol* 2007;103:637-645.
- 88 Shibasaki M, Hashimoto K, Okamoto M, Hayashi Y, Imaizumi K, Hashimoto N, Ozaki N, Yokoi T, Takagi K, Hasegawa Y, Shimokata K, Kawabe T: Up-regulation of surfactant protein production in a mouse model of secondary pulmonary alveolar proteinosis. *Am J Respir Cell Mol Biol* 2009;40:536-542.
- 89 Atochina-Vasserman EN, Gow AJ, Abramova H, Guo CJ, Tomer Y, Preston AM, Beck JM, Beers MF: Immune reconstitution during pneumocystis lung infection: Disruption of surfactant component expression and function by s-nitrosylation. *J Immunol* 2009;182:2277-2287.
- 90 Atochina EN, Gow AJ, Beck JM, Haczku A, Inch A, Kadire H, Tomer Y, Davis C, Preston AM, Poulain F, Hawgood S, Beers MF: Delayed clearance of pneumocystis carinii infection, increased inflammation, and altered nitric oxide metabolism in lungs of surfactant protein-d knockout mice. *J Infect Dis* 2004;189:1528-1539.
- 91 Atochina EN, Beck JM, Scanlon ST, Preston AM, Beers MF: Pneumocystis carinii pneumonia alters expression and distribution of lung collectins sp-a and sp-d. *J Lab Clin Med* 2001;137:429-439.
- 92 Kobayashi M, Takeuchi T, Ohtsuki Y: Differences in the immunolocalization of surfactant protein (sp)-a, sp-d, and kl-6 in pulmonary alveolar proteinosis. *Pathol Int* 2008;58:203-207.
- 93 Voss T, Schafer KP, Nielsen PF, Schafer A, Maier C, Hannappel E, Maassen J, Landis B, Klemm K, Przybylski M: Primary structure differences of human surfactant-associated proteins isolated from normal and proteinosis lung. *Biochim Biophys Acta* 1992;1138:261-267.
- 94 Galetskiy D, Woischnik M, Ripper J, Griesse M, Przybylski M: Aberrant processing forms of lung surfactant proteins sp-b and sp-c revealed by high-resolution mass spectrometry. *Eur J Mass Spectrom* (Chichester, Eng) 2008;14:379-390.
- 95 Vorbroker DK, Proffitt SA, Noguee LM, Whitsett JA: Aberrant processing of surfactant protein c in hereditary sp-b deficiency. *Am J Physiol* 1995;268:L647-656.
- 96 Gupta G, Suroliya A: Collectins: Sentinels of Innate Immunity. *Bioessays* 2007; 29: 452-463.
- 97 Crouch E, McDonald B, Smith K, Cafarella T, Seaton B, Head J: Contributions of phenylalanine 335 to ligand recognition by human surfactant protein d: Ring interactions with sp-d ligands. *J Biol Chem* 2006;281:18008-18014.
- 98 Head JF, Mealy TR, McCormack FX, Seaton BA: Crystal structure of trimeric carbohydrate recognition and neck domains of surfactant protein a. *J Biol Chem* 2003;278:43254-43260.
- 99 Palaniyar N, Ikegami M, Korfhagen T, Whitsett J, McCormack FX: Domains of surfactant protein a that affect protein oligomerization, lipid structure and surface tension. *Comp Biochem Physiol A Mol Integr Physiol* 2001;129:109-127.
- 100 Yu SH, McCormack FX, Voelker DR, Possmayer F: Interactions of pulmonary surfactant protein sp-a with monolayers of dipalmitoylphosphatidylcholine and cholesterol: Roles of sp-a domains. *J Lipid Res* 1999;40:920-929.

- 101 Chabot S, Koumanov K, Lambeau G, Gelb MH, Balloy V, Chignard M, Whitsett JA, Touqui L: Inhibitory effects of surfactant protein a on surfactant phospholipid hydrolysis by secreted phospholipases a2. *J Immunol* 2003;171:995-1000.
- 102 Wu YZ, Manevich Y, Baldwin JL, Dodia C, Yu K, Feinstein SI, Fisher AB: Interaction of surfactant protein a with peroxiredoxin 6 regulates phospholipase a2 activity. *J Biol Chem* 2006;281:7515-7525.
- 103 Bates SR, Kazi AS, Tao JQ, Yu KJ, Gonder DS, Feinstein SI, Fisher AB: Role of p63 (ckap4) in binding of surfactant protein-a to type ii pneumocytes. *Am J Physiol Lung Cell Mol Physiol* 2008;295:L658-669.
- 104 Wright JR, Youmans DC: Degradation of surfactant lipids and surfactant protein a by alveolar macrophages in vitro. *Am J Physiol* 1995;268:L772-780.
- 105 Zhang L, Ikegami M, Korfhagen TR, McCormack FX, Yoshida M, Senior RM, Shipley JM, Shapiro SD, Whitsett JA: Neither sp-a nor nh2-terminal domains of sp-a can substitute for sp-d in regulation of alveolar homeostasis. *Am J Physiol Lung Cell Mol Physiol* 2006;291:L181-190.
- 106 Palaniyar N, Zhang L, Kuzmenko A, Ikegami M, Wan S, Wu H, Korfhagen TR, Whitsett JA, McCormack FX: The role of pulmonary collectin n-terminal domains in surfactant structure, function, and homeostasis in vivo. *J Biol Chem* 2002;277:26971-26979.
- 107 Wu H, Kuzmenko A, Wan S, Schaffer L, Weiss A, Fisher JH, Kim KS, McCormack FX: Surfactant proteins a and d inhibit the growth of gram-negative bacteria by increasing membrane permeability. *J Clin Invest* 2003;111:1589-1602.
- 108 McCormack FX: New concepts in collectin-mediated host defense at the air-liquid interface of the lung. *Respirology* 2006;11 Suppl:S7-10.
- 109 Piboonpocanun S, Chiba H, Mitsuzawa H, Martin W, Murphy RC, Harbeck RJ, Voelker DR: Surfactant protein a binds mycoplasma pneumoniae with high affinity and attenuates its growth by recognition of disaturated phosphatidylglycerols. *J Biol Chem* 2005;280:9-17.
- 110 Pikaar JC, Voorhout WF, van Golde LM, Verhoef J, Van Strijp JA, van Iwaarden JF: Opsonic activities of surfactant proteins a and d in phagocytosis of gram-negative bacteria by alveolar macrophages. *J Infect Dis* 1995;172:481-489.
- 111 McCormack FX, Gibbons R, Ward SR, Kuzmenko A, Wu H, Deepe GS, Jr.: Macrophage-independent fungicidal action of the pulmonary collectins. *J Biol Chem* 2003;278:36250-36256.
- 112 Van Iwaarden JF, Pikaar JC, Storm J, Brouwer E, Verhoef J, Oosting RS, van Golde LM, van Strijp JA: Binding of surfactant protein a to the lipid moiety of bacterial lipopolysaccharides. *Biochem J* 1994;303:407-411.
- 113 Kannan TR, Provenzano D, Wright JR, Baseman JB: Identification and characterization of human surfactant protein a binding protein of mycoplasma pneumoniae. *Infect Immun* 2005;73:2828-2834.
- 114 Kannan TR, Baseman JB: Adp-ribosylating and vacuolating cytotoxin of mycoplasma pneumoniae represents unique virulence determinant among bacterial pathogens. *Proc Natl Acad Sci U S A* 2006;103:6724-6729.
- 115 Ragas A, Roussel L, Puzo G, Riviere M: The mycobacterium tuberculosis cell-surface glycoprotein apa as a potential adhesin to colonize target cells via the innate immune system pulmonary c-type lectin surfactant protein a. *J Biol Chem* 2007;282:5133-5142.
- 116 Sever-Chroneos Z, Geisbrecht B, Chroneos ZC: Surfactant protein a interacts with the staphylococcus aureus adhesin eap/map. *Am J Resp Crit Care Med* 2008;177:A585.
- 117 Sahly H, Ofek I, Podschun R, Brade H, He Y, Ullmann U, Crouch E: Surfactant protein d binds selectively to klebsiella pneumoniae lipopolysaccharides containing mannose-rich o-antigens. *J Immunol* 2002;169:3267-3274.
- 118 Wang H, Head J, Kosma P, Brade H, Muller-Loennies S, Sheikh S, McDonald B, Smith K, Cafarella T, Seaton B, Crouch E: Recognition of heptoses and the inner core of bacterial lipopolysaccharides by surfactant protein d. *Biochemistry* 2008;47:710-720.
- 119 Stamme C, Wright JR: Surfactant protein a enhances the binding and deacylation of e. Coli lps by alveolar macrophages. *Am J Physiol* 1999;276:L540-547.
- 120 van Rozendaal BA, van de Lest CH, van Eijk M, van Golde LM, Voorhout WF, van Helden HP, Haagsman HP: Aerosolized endotoxin is immediately bound by pulmonary surfactant protein d in vivo. *Biochim Biophys Acta* 1999;1454:261-269.
- 121 Moulakakis C, Stamme C: Role of clathrin-mediated endocytosis of surfactant protein a by alveolar macrophages in intracellular signaling. *Am J Physiol Lung Cell Mol Physiol* 2009;296:L430-441.
- 122 Kuzmenko AI, Wu H, McCormack FX: Pulmonary collectins selectively permeabilize model bacterial membranes containing rough lipopolysaccharide. *Biochemistry* 2006;45:2679-2685.
- 123 Canadas O, Garcia-Verdugo I, Keough KM, Casals C: Sp-a permeabilizes lipopolysaccharide membranes by forming protein aggregates that extract lipids from the membrane. *Biophys J* 2008;95:3287-3294.
- 124 Wang L, Brauner JW, Mao G, Crouch E, Seaton B, Head J, Smith K, Flach CR, Mendelsohn R: Interaction of recombinant surfactant protein d with lipopolysaccharide: Conformation and orientation of bound protein by irras and simulations. *Biochemistry* 2008;47:8103-8113.
- 125 Garcia-Verdugo I, Canadas O, Taneva SG, Keough KM, Casals C: Surfactant protein a forms extensive lattice-like structures on 1,2-dipalmitoylphosphatidylcholine/rough-lipopolysaccharide-mixed monolayers. *Biophys J* 2007;93:3529-3540.
- 126 Weikert LF, Lopez JP, Abdolrasulnia R, Chroneos ZC, Shepherd VL: Surfactant protein a enhances mycobacterial killing by rat macrophages through a nitric oxide-dependent pathway. *Am J Physiol Lung Cell Mol Physiol* 2000;279:L216-223.
- 127 Weikert LF, Edwards K, Chroneos ZC, Hager C, Hoffman L, Shepherd VL: Sp-a enhances uptake of bacillus calmette-guerin by macrophages through a specific sp-a receptor. *Am J Physiol* 1997;272:L989-995.
- 128 Lopez JP, Vigerust DJ, Shepherd VL: Mitogen-activated protein kinases and nfkappab are involved in sp-a-enhanced responses of macrophages to mycobacteria. *Respir Res* 2009;10:60.
- 129 Chiba H, Piboonpocanun S, Mitsuzawa H, Kuronuma K, Murphy RC, Voelker DR: Pulmonary surfactant proteins and lipids as modulators of inflammation and innate immunity. *Respirology* 2006;11 Suppl:S2-6.
- 130 Barr FE, Pedigo H, Johnson TR, Shepherd VL: Surfactant protein-a enhances uptake of respiratory syncytial virus by monocytes and u937 macrophages. *Am J Respir Cell Mol Biol* 2000;23:586-592.
- 131 Kostina E, Ofek I, Crouch E, Friedman R, Sirota L, Klinger G, Sahly H, Keisari Y: Noncapsulated klebsiella pneumoniae bearing mannose-containing o antigens is rapidly eradicated from mouse lung and triggers cytokine production by macrophages following opsonization with surfactant protein d. *Infect Immun* 2005;73:8282-8290.
- 132 LeVine AM, Kurak KE, Wright JR, Watford WT, Bruno MD, Ross GF, Whitsett JA, Korfhagen TR: Surfactant protein-a binds group b streptococcus enhancing phagocytosis and clearance from lungs of surfactant protein-a-deficient mice. *Am J Respir Cell Mol Biol* 1999;20:279-286.

- 133 Hickman-Davis JM, Gibbs-Erwin J, Lindsey JR, Matalon S: Role of surfactant protein-a in nitric oxide production and mycoplasma killing in congenic c57bl/6 mice. *Am J Respir Cell Mol Biol* 2004;30:319-325.
- 134 Hickman-Davis JM, Lindsey JR, Zhu S, Matalon S: Surfactant protein a mediates mycoplasma killing in congenic alveolar macrophages. *Am J Physiol* 1998;274:L270-277.
- 135 Giannoni E, Sawa T, Allen L, Wiener-Kronish J, Hawgood S: Surfactant proteins a and d enhance pulmonary clearance of *Pseudomonas aeruginosa*. *Am J Respir Cell Mol Biol* 2006;34:704-710.
- 136 Jounblat R, Clark H, Eggleton P, Hawgood S, Andrew PW, Kadioglu A: The role of surfactant protein d in the colonisation of the respiratory tract and onset of bacteraemia during pneumococcal pneumonia. *Respir Res* 2005;6:126.
- 137 LeVine AM, Elliott J, Whitsett JA, Srikiatkachorn A, Crouch E, DeSilva N, Korfhagen T: Surfactant protein-d enhances phagocytosis and pulmonary clearance of respiratory syncytial virus. *Am J Respir Cell Mol Biol* 2004;31:193-199.
- 138 Sano H, Sohma H, Muta T, Nomura S, Voelker DR, Kuroki Y: Pulmonary surfactant protein a modulates the cellular response to smooth and rough lipopolysaccharides by interaction with cd14. *J Immunol* 1999;163:387-395.
- 139 McIntosh JC, Mervin-Blake S, Conner E, Wright JR: Surfactant protein a protects growing cells and reduces tnfr-alpha activity from lps-stimulated macrophages. *Am J Physiol* 1996;271:L310-319.
- 140 Borron P, McIntosh JC, Korfhagen TR, Whitsett JA, Taylor J, Wright JR: Surfactant-associated protein a inhibits lps-induced cytokine and nitric oxide production in vivo. *Am J Physiol Lung Cell Mol Physiol* 2000;278:L840-847.
- 141 Arias-Diaz J, Garcia-Verdugo I, Casals C, Sanchez-Rico N, Vara E, Balibrea JL: Effect of surfactant protein a (sp-a) on the production of cytokines by human pulmonary macrophages. *Shock* 2000;14:300-306.
- 142 Moulakakis C, Adam S, Seitzer U, Schromm AB, Leitges M, Stamme C: Surfactant protein a activation of atypical protein kinase c zeta in ikappab-alpha-dependent anti-inflammatory immune regulation. *J Immunol* 2007;179:4480-4491.
- 143 Wu Y, Adam S, Hamann L, Heine H, Ulmer AJ, Buwitt-Beckmann U, Stamme C: Accumulation of inhibitory kappab-alpha as a mechanism contributing to the anti-inflammatory effects of surfactant protein-a. *Am J Respir Cell Mol Biol* 2004;31:587-594.
- 144 Stamme C, Muller M, Hamann L, Gutschmann T, Seydel U: Surfactant protein a inhibits lipopolysaccharide-induced immune cell activation by preventing the interaction of lipopolysaccharide with lipopolysaccharide-binding protein. *Am J Respir Cell Mol Biol* 2002;27:353-360.
- 145 Samten B, Townsend JC, Sever-Chroneos Z, Pasquinelli V, Barnes PF, Chronesos ZC: An antibody against the surfactant protein a (sp-a)-binding domain of the sp-a receptor inhibits t cell-mediated immune responses to mycobacterium tuberculosis. *J Leukoc Biol* 2008;84:115-123.
- 146 Ledford JG, Goto H, Potts EN, Degan S, Chu HW, Voelker DR, Sunday ME, Cianciolo GJ, Foster WM, Kraft M, Wright JR: Sp-a preserves airway homeostasis during mycoplasma pneumoniae infection in mice. *J Immunol* 2009;182:7818-7827.
- 147 Schagat TL, Wofford JA, Wright JR: Surfactant protein a enhances alveolar macrophage phagocytosis of apoptotic neutrophils. *J Immunol* 2001;166:2727-2733.
- 148 Janssen WJ, McPhillips KA, Dickinson MG, Linderman DJ, Morimoto K, Xiao YQ, Oldham KM, Vandivier RW, Henson PM, Gardai SJ: Surfactant proteins a and d suppress alveolar macrophage phagocytosis via interaction with sirp alpha. *Am J Respir Crit Care Med* 2008;178:158-167.
- 149 Reidy MF, Wright JR: Surfactant protein a enhances apoptotic cell uptake and tgfbeta1 release by inflammatory alveolar macrophages. *Am J Physiol Lung Cell Mol Physiol* 2003;285:L854-861.
- 150 Koth LL, Alex B, Hawgood S, Nead MA, Sheppard D, Erle DJ, Morris DG: Integrin beta6 mediates phospholipid and collectin homeostasis by activation of latent tgfbeta1. *Am J Respir Cell Mol Biol* 2007;37:651-659.
- 151 Akei H, Whitsett JA, Buroker M, Ninomiya T, Tatsumi H, Weaver TE, Ikegami M: Surface tension influences cell shape and phagocytosis in alveolar macrophages. *Am J Physiol Lung Cell Mol Physiol* 2006;291:L572-579.
- 152 Lawson WE, Polosukhin VV, Stathopoulos GT, Zoia O, Han W, Lane KB, Li B, Donnelly EF, Holburn GE, Lewis KG, Collins RD, Hull WM, Glasser SW, Whitsett JA, Blackwell TS: Increased and prolonged pulmonary fibrosis in surfactant protein c-deficient mice following intratracheal bleomycin. *Am J Pathol* 2005;167:1267-1277.
- 153 Garcia-Verdugo I, Garcia de Paco E, Espinassous Q, Gonzalez-Horta A, Synguelakis M, Kanellopoulos J, Rivas L, Chaby R, Perez-Gil J: Synthetic peptides representing the n-terminal segment of surfactant protein c modulate lps-stimulated tnfr-alpha production by macrophages. *Innate Immun* 2009;15:53-62.
- 154 Augusto LA, Li J, Synguelakis M, Johansson J, Chaby R: Structural basis for interactions between lung surfactant protein c and bacterial lipopolysaccharide. *J Biol Chem* 2002;277:23484-23492.
- 155 Augusto L, Le Blay K, Auger G, Blanot D, Chaby R: Interaction of bacterial lipopolysaccharide with mouse surfactant protein c inserted into lipid vesicles. *Am J Physiol Lung Cell Mol Physiol* 2001;281:L776-785.
- 156 Morris RH, Tonks AJ, Jones KP, Ahluwalia MK, Thomas AW, Tonks A, Jackson SK: Dppc regulates cox-2 expression in monocytes via phosphorylation of creb. *Biochem Biophys Res Commun* 2008;370:174-178.
- 157 Tonks A, Parton J, Tonks AJ, Morris RH, Finall A, Jones KP, Jackson SK: Surfactant phospholipid dppc downregulates monocyte respiratory burst via modulation of pkc. *Am J Physiol Lung Cell Mol Physiol* 2005;288:L1070-1080.
- 158 Miles PR, Bowman L, Rao KM, Baatz JE, Huffman L: Pulmonary surfactant inhibits lps-induced nitric oxide production by alveolar macrophages. *Am J Physiol* 1999;276:L186-196.
- 159 Blanco O, Perez-Gil J: Biochemical and pharmacological differences between preparations of exogenous natural surfactant used to treat respiratory distress syndrome: Role of the different components in an efficient pulmonary surfactant. *Eur J Pharmacol* 2007;568:1-15.
- 160 Kuronuma K, Mitsuzawa H, Takeda K, Nishitani C, Chan ED, Kuroki Y, Nakamura M, Voelker DR: Anionic pulmonary surfactant phospholipids inhibit inflammatory responses from alveolar macrophages and u937 cells by binding the lipopolysaccharide interacting proteins cd14 and md2. *J Biol Chem* 2009;284:25488-25500.
- 161 Abate W, Alghaithy AA, Parton J, Jones KP, Jackson SK: Surfactant lipids regulate lps-induced interleukin-8 production in a549 lung epithelial cells by inhibiting translocation of tlr4 into lipid raft domains. *J Lipid Res* 2009;in press.

- 162 Gille C, Spring B, Bernhard W, Gebhard C, Basile D, Lauber K, Poets CF, Orlikowsky TW: Differential effect of surfactant and its saturated phosphatidylcholines on human blood macrophages. *J Lipid Res* 2007;48:307-317.
- 163 Kuronuma K, Sano H, Kato K, Kudo K, Hyakushima N, Yokota S, Takahashi H, Fujii N, Suzuki H, Kodama T, Abe S, Kuroki Y: Pulmonary surfactant protein a augments the phagocytosis of streptococcus pneumoniae by alveolar macrophages through a casein kinase 2-dependent increase of cell surface localization of scavenger receptor a. *J Biol Chem* 2004;279:21421-21430.
- 164 Gil M, McCormack FX, Levine AM: Surfactant protein-a modulates cell surface expression of cr3 on alveolar macrophages and enhances cr3-mediated phagocytosis. *J Biol Chem* 2009;284:7495-7504.
- 165 Henning LN, Azad AK, Parsa KV, Crowther JE, Tridandapani S, Schlesinger LS: Pulmonary surfactant protein a regulates tlr expression and activity in human macrophages. *J Immunol* 2008;180:7847-7858.
- 166 Kudo K, Sano H, Takahashi H, Kuronuma K, Yokota S, Fujii N, Shimada K, Yano I, Kumazawa Y, Voelker DR, Abe S, Kuroki Y: Pulmonary collectins enhance phagocytosis of mycobacterium avium through increased activity of mannose receptor. *J Immunol* 2004;172:7592-7602.
- 167 Beharka AA, Gaynor CD, Kang BK, Voelker DR, McCormack FX, Schlesinger LS: Pulmonary surfactant protein a up-regulates activity of the mannose receptor, a pattern recognition receptor expressed on human macrophages. *J Immunol* 2002;169:3565-3573.
- 168 Tenner AJ, Robinson SL, Borchelt J, Wright JR: Human pulmonary surfactant protein (sp-a), a protein structurally homologous to clq, can enhance fcr- and crl-mediated phagocytosis. *J Biol Chem* 1989;264:13923-13928.
- 169 Senft AP, Korfhagen TR, Whitsett JA, LeVine AM: Surfactant protein d regulates the cell surface expression of alveolar macrophage beta(2)-integrins. *Am J Physiol Lung Cell Mol Physiol* 2007;292:L469-475.
- 170 Senft A, Korfhagen T, Whitsett JA, Levine AM: Surfactant protein-d regulates the cell surface expression of alveolar macrophage {beta}2 integrins. *Am J Physiol Lung Cell Mol Physiol* 2007;292:L469-475.
- 171 Guth AM, Janssen WJ, Bosio CM, Crouch EC, Henson PM, Dow SW: Lung environment determines unique phenotype of alveolar macrophages. *Am J Physiol Lung Cell Mol Physiol* 2009;296:L936-946.
- 172 Poelma DL, Lachmann B, Haitzma JJ, Zimmermann LJ, van Iwaarden JF: Influence of phosphatidylglycerol on the uptake of liposomes by alveolar cells and on lung function. *J Appl Physiol* 2005;98:1784-1791.
- 173 Quintero OA, Wright JR: Metabolism of phosphatidylglycerol by alveolar macrophages in vitro. *Am J Physiol Lung Cell Mol Physiol* 2000;279:L399-407.
- 174 Arosa FA, de Jesus O, Porto G, Carmo AM, de Sousa M: Calreticulin is expressed on the cell surface of activated human peripheral blood t lymphocytes in association with major histocompatibility complex class i molecules. *J Biol Chem* 1999;274:16917-16922.
- 175 White TK, Zhu Q, Tanzer ML: Cell surface calreticulin is a putative mannoside lectin which triggers mouse melanoma cell spreading. *J Biol Chem* 1995;270:15926-15929.
- 176 Stuart GR, Lynch NJ, Day AJ, Schwaeble WJ, Sim RB: The clq and collectin binding site within clq receptor (cell surface calreticulin). *Immunopharmacology* 1997;38:73-80.
- 177 Basu S, Binder RJ, Ramalingam T, Srivastava PK: Cd91 is a common receptor for heat shock proteins gp96, hsp90, hsp70, and calreticulin. *Immunity* 2001;14:303-313.
- 178 Lillis AP, Van Duyn LB, Murphy-Ullrich JE, Strickland DK: Ldl receptor-related protein 1: Unique tissue-specific functions revealed by selective gene knockout studies. *Physiol Rev* 2008;88:887-918.
- 179 Gardai SJ, Xiao YQ, Dickinson M, Nick JA, Voelker DR, Greene KE, Henson PM: By binding sirpalpha or calreticulin/cd91, lung collectins act as dual function surveillance molecules to suppress or enhance inflammation. *Cell* 2003;115:13-23.
- 180 Vandivier RW, Ogden CA, Fadok VA, Hoffmann PR, Brown KK, Botto M, Walport MJ, Fisher JH, Henson PM, Greene KE: Role of surfactant proteins a, d, and clq in the clearance of apoptotic cells in vivo and in vitro: Calreticulin and cd91 as a common collectin receptor complex. *J Immunol* 2002;169:3978-3986.
- 181 van Beek EM, Cochrane F, Barclay AN, van den Berg TK: Signal regulatory proteins in the immune system. *J Immunol* 2005;175:7781-7787.
- 182 Tsai RK, Discher DE: Inhibition of "Self" Engulfment through deactivation of myosin-ii at the phagocytic synapse between human cells. *J Cell Biol* 2008;180:989-1003.
- 183 Kong XN, Yan HX, Chen L, Dong LW, Yang W, Liu Q, Yu LX, Huang DD, Liu SQ, Liu H, Wu MC, Wang HY: Lps-induced down-regulation of signal regulatory protein {alpha} contributes to innate immune activation in macrophages. *J Exp Med* 2007;204:2719-2731.
- 184 Lillis AP, Greenlee MC, Mikhailenko I, Pizzo SV, Tenner AJ, Strickland DK, Bohlson SS: Murine low-density lipoprotein receptor-related protein 1 (lrp) is required for phagocytosis of targets bearing lrp ligands but is not required for clq-triggered enhancement of phagocytosis. *J Immunol* 2008;181:364-373.
- 185 Walters JJ, Berwin B: Differential cd91 dependence for calreticulin and pseudomonas exotoxin-a endocytosis. *Traffic (Copenhagen, Denmark)* 2005;6:1173-1182.
- 186 Berwin B, Hart JP, Rice S, Gass C, Pizzo SV, Post SR, Nicchitta CV: Scavenger receptor-a mediates gp96/grp94 and calreticulin internalization by antigen-presenting cells. *The EMBO journal* 2003;22:6127-6136.
- 187 Oldenburg PA, Gresham HD, Lindberg FP: Cd47-signal regulatory protein alpha (sirpalpha) regulates fcgamma and complement receptor-mediated phagocytosis. *J Exp Med* 2001;193:855-862.
- 188 Olsson M, Oldenburg PA: Cd47 on experimentally senescent murine rbc inhibits phagocytosis following fcgamma receptor-mediated but not scavenger receptor-mediated recognition by macrophages. *Blood* 2008;112:4259-4267.
- 189 Ohya M, Nishitani C, Sano H, Yamada C, Mitsuzawa H, Shimizu T, Saito T, Smith K, Crouch E, Kuroki Y: Human pulmonary surfactant protein d binds the extracellular domains of toll-like receptors 2 and 4 through the carbohydrate recognition domain by a mechanism different from its binding to phosphatidylinositol and lipopolysaccharide. *Biochemistry* 2006;45:8657-8664.
- 190 Yamada C, Sano H, Shimizu T, Mitsuzawa H, Nishitani C, Himi T, Kuroki Y: Surfactant protein a directly interacts with tlr4 and md-2 and regulates inflammatory cellular response. Importance of supratrimeric oligomerization. *J Biol Chem* 2006;281:21771-21780.

- 191 Sato M, Sano H, Iwaki D, Kudo K, Konishi M, Takahashi H, Takahashi T, Imaizumi H, Asai Y, Kuroki Y: Direct binding of toll-like receptor 2 to zymosan, and zymosan-induced nf-kappa b activation and tnf-alpha secretion are down-regulated by lung collectin surfactant protein a. *J Immunol* 2003;171:417-425.
- 192 Murakami S, Iwaki D, Mitsuzawa H, Sano H, Takahashi H, Voelker DR, Akino T, Kuroki Y: Surfactant protein a inhibits peptidoglycan-induced tumor necrosis factor-alpha secretion in u937 cells and alveolar macrophages by direct interaction with toll-like receptor 2. *J Biol Chem* 2002;277:6830-6837.
- 193 Sano H, Chiba H, Iwaki D, Sohma H, Voelker DR, Kuroki Y: Surfactant proteins a and d bind cd14 by different mechanisms. *J Biol Chem* 2000;275:22442-22451.
- 194 Yamazoe M, Nishitani C, Takahashi M, Katoh T, Arikawa S, Shimizu T, Mitsuzawa H, Sawada K, Voelker DR, Takahashi H, Kuroki Y: Pulmonary surfactant protein d inhibits lipopolysaccharide (lps)-induced inflammatory cell responses by altering lps binding to its receptors. *J Biol Chem* 2008;283:35878-35888.
- 195 Chroneos ZC, Abdolrasulnia R, Whitsett JA, Rice WR, Shepherd VL: Purification of a cell-surface receptor for surfactant protein a. *J Biol Chem* 1996;271:16375-16383.
- 196 Borron P, McCormack FX, Elhalwagi BM, Chroneos ZC, Lewis JF, Zhu S, Wright JR, Shepherd VL, Possmayer F, Inchley K, Fraher LJ: Surfactant protein a inhibits t cell proliferation via its collagen-like tail and a 210-kda receptor. *Am J Physiol* 1998;275:L679-686.
- 197 Garcia-Verdugo I, Leiber D, Robin P, Billon-Denis E, Chaby R, Tanfin Z: Direct interaction of surfactant protein a with myometrial binding sites: Signaling and modulation by bacterial lipopolysaccharide. *Biol Reprod* 2007;76:681-691.
- 198 Quesniaux VJ, Nicolle DM, Torres D, Kremer L, Guerardel Y, Nigou J, Puzo G, Erard F, Ryffel B: Toll-like receptor 2 (tlr2)-dependent-positive and tlr2-independent-negative regulation of proinflammatory cytokines by mycobacterial lipomannans. *J Immunol* 2004;172:4425-4434.
- 199 Yang CH, Szeliga J, Jordan J, Faske S, Sever-Chroneos Z, Dorsett B, Christian RE, Settlege RE, Shabanowitz J, Hunt DF, Whitsett JA, Chroneos ZC: Identification of the surfactant protein a receptor 210 as the unconventional myosin 18a. *J Biol Chem* 2005;280:34447-34457.
- 200 Furusawa T, Ikawa S, Yanai N, Obinata M: Isolation of a novel pdz-containing myosin from hematopoietic supportive bone marrow stromal cell lines. *Biochem Biophys Res Commun* 2000;270:67-75.
- 201 Mori K, Furusawa T, Okubo T, Inoue T, Ikawa S, Yanai N, Mori KJ, Obinata M: Genome structure and differential expression of two isoforms of a novel pdz-containing myosin (myspdz) (myo18a). *J Biochem (Tokyo)* 2003;133:405-413.
- 202 Tan I, Yong J, Dong JM, Lim L, Leung T: A tripartite complex containing mrck modulates lamellar actomyosin retrograde flow. *Cell* 2008;135:123-136.
- 203 Lv T, Shen X, Shi Y, Song Y: Tlr4 is essential in acute lung injury induced by unresuscitated hemorrhagic shock. *J Trauma* 2009;66:124-131.
- 204 Hoth JJ, Wells JD, Brownlee NA, Hiltbold EM, Meredith JW, McCall CE, Yoza BK: Toll-like receptor 4-dependent responses to lung injury in a murine model of pulmonary contusion. *Shock* 2009;31:376-381.
- 205 Imai Y, Kuba K, Neely GG, Yaghubian-Malhami R, Perkmann T, van Loo G, Ermolaeva M, Veldhuizen R, Leung YH, Wang H, Liu H, Sun Y, Pasparakis M, Kopf M, Mech C, Bavari S, Peiris JS, Slutsky AS, Akira S, Hultqvist M, Holmdahl R, Nicholls J, Jiang C, Binder CJ, Penninger JM: Identification of oxidative stress and toll-like receptor 4 signaling as a key pathway of acute lung injury. *Cell* 2008;133:235-249.