Preface

The phenomenon of aging is characterized by various degenerative changes which differentially affect the highly specialized structures within the eye, such as the purely cellular lens, the brain-derived retina and the connective tissue of the uvea and sclera. The eye can therefore serve as an excellent model system to study age-related degenerative diseases.

Glaucoma is a common age-related ocular disease. Studies in glaucomatous eyes have shown that there is a correlation between optic nerve fiber loss and changes in the outflow tissues, with the latter being presumably responsible for increased intraocular pressure (IOP). Another finding in glaucoma is that increased expression of the stress protein α B-crystallin occurs in TM of glaucomatous eyes. α B-crystallin is one of the proteins normally protecting lens fiber proteins from unfolding due to various stress factors. In this volume, an overview article deals with the molecular biology of this molecule, and original articles give further insights into the distribution and possible functional significance of nonlenticular α B-crystallin. A better understanding of the role of protective molecules in the eye might open new perspectives in the treatment of degenerative diseases.

The commonest causes of blindness are the tapetoretinal degenerations and age-related macular degeneration.

Retinal transplantation studies have been undertaken that may ultimately lead to therapy for these diseases. New kinds of laminated transplants with promising survival times have been implanted and studied using different immunohistochemical methods. Further, an attempt has been made to investigate factors that influence the development of the retinal tissues, since less differentiated embryonic retina has the potential to be well integrated into degenerated retina. In the essentially cellular lens, it is difficult to define when development ends and ageing begins. An overview article examines cells of the deep lens fibers. These cells lose organelles and finally undergo denucleation, even though they remain viable for the life of the lens. The membranous channel system of the lens fibers plays a role in retaining transparency of the lens. In vitro, oxidative damage is one factor responsible for inducing changes in these lenticular ion channels. Protection against oxidative damage might therefore protect against cataract formation.

The studies were kindly supported by the European Commission as part of the Biomed project BMH4-CT96-1593 (Brussels).

We thank especially Mr. Kallasvaara for his help and his patience.

E. Lütjen-Drecoll, Erlangen