Rare Benign Neoplasms of the Larynx: Rhabdomyoma and Lipoma

by Zbären et al. in ORL 1995;57:351-355

I.	Friedmann
----	-----------

Northwick Park Hospital, Harrow, Middlesex, UK

Prof. I. Friedmann, Northwick Park Hospital, Watford Road, Harrow, Middlesex HA1 3UJ (UK)

Dear Sir,

I have been reading with great interest the comprehensive paper by Zbären et al. [1] on 'Rare benign neoplasms of the larynx: Rhabdomyoma and lipoma' in ORL 1995; 57:351-355 and I would like to make two points.

First a general point about so-called 'reviews of the literature': A recent paper on lipomas of the larynx [2] has suggested that approximately 80 cases had been reported adding 3 cases of their own. The present authors basing their review equally on Zakrzewski's total of 65 reported by 1965 suggest that about 20 more cases had been reported bringing the total to 85 not including the 3 new cases nor the 2 cases we have seen [3]. One of these patients was an elderly woman who was found dead by her son on the kitchen floor 'with some meat protruding from her lips'. The illustration (fig. 26.1) shows a large pedunculated tumour diagnosed microscopically as a benign lipoma which the

poor woman had disguised for years complaining of a cough from time to time. The other case only illustrated was from an elderly man and was removed surgically. A similar case, only the 3rd one in my long experience, was a subglottic lipoma removed surgically from the larynx of an 83-year-old man complaining of dysphagia of long duration (to be published). There was no evidence of malignancy in any of these cases but Wenig [2] suggests that some of the liposarcomas may have been overlooked in the surgical material and might only be recognised retrospectively on a recurrent lipoma. Neither of our cases would fall into this category.

This case would bring the total to 86 or 91. I have suggested that reference to and discussion of published cases of any neoplasm were essential in any paper but neither the claim of a 'first case' [4] nor the 'totals' quoted might prove to be correct.

References

Zbären P, Lang H, Becker M: Rare benign neoplasms of the larynx: Rhabdomyoma and lipoma. ORL 1995;57:351-355. Wenig BM: Lipomas of the larynx: A review of the literature with the addition of three new cases. J Laryngol Otol 1995;109:353-357. Friedmann I, Ferlito A: Adipose tissue and synovial tumours; in Friedmann I, Ferlito A (eds): Granulomas and Neoplasms of the Larynx. Edinburgh, Churchill Livingstone, 1988, chapter 26, p 303. Treasure T: What is the place of the clinical case report in medical publishing? J R Soc Med 1995;88:279.

KÄIUER

E-Mail karger@karger.ch Fax+ 41 61 306 12 34 <http://www.karger.ch>

© 1996 S. KargerAG, Basel 0301-1569/96/O583-0182\$10.00/O