Impaired Neurotransmitter Release in Alzheimer's and Parkinson's Diseases

Jie Shen

Center for Neurologic Diseases, Brigham and Women's Hospital, Program in Neuroscience, Harvard Medical School, Boston, Mass., USA

Key Words

Synapse • Age-related neurodegeneration • Presenilin • Parkin • Nicastrin

Abstract

Mutations in several causative genes have been linked to monogenic forms of Alzheimer's disease (AD) or Parkinson's disease (PD). To look for possible common pathogenic mechanisms underlying age-related neurodegeneration in AD and PD, we employed genetic approaches to investigate systematically the roles of these gene products (e.g. presenilins (PS) for AD; Parkin, DJ-1, PINK1 and LRRK2 for PD) in the mouse brain, especially in neural circuits that are particularly vulnerable in AD or PD. Our series of genetic studies revealed that PS play cell type-specific roles in the developing brain with the most prominent function in the maintenance of neural progenitor cells. In the adult cerebral cortex, where the pathogenesis of AD occurs, loss of PS results in progressive memory impairment and age-related neurodegeneration. Specifically, PS are involved in the regulation of long-term potentiation and NMDA receptor functions. Interestingly, our further genetic dissection in the hippocampal Schaeffer collateral pathway highlighted the importance of presynaptic PS in the activity-dependent regulation of glutamate release and long-term potentiation induction via

modulation of calcium release from intracellular stores. Intriguingly, our independent genetic analysis of Parkin, DJ-1, PINK1 and LRRK2 showed a common defect in activity-dependent dopamine release caused by PD-linked mutations in these genes. Together, our genetic studies suggest that presynaptic dysfunction might be a converging early pathogenic event before neurodegeneration in AD and PD.

Copyright © 2010 S. Karger AG, Basel

Alzheimer's disease (AD) is the most common form of dementia and neurodegenerative disorder. AD is characterized clinically by progressive memory loss and deterioration of cognitive functions, and neuropathologically by extracellular amyloid plaques, intracellular neurofibrillary tangles, and synaptic and neuronal loss. Presenilins (PS1 and PS2) are the major causative genes of early-onset (<65 years of age) familial AD and harbor ~90% of FAD mutations identified to date. PS are essential components of γ -secretase, a multi-subunit protease complex that catalyzes the intramembranous cleavage of a number of type I transmembrane proteins, including Notch and the amyloid precursor protein. Notch is a key physiological substrate of γ -secretase, as evidenced by similar developmental phenotypes exhibited by PS and Notch mutant mice [reviewed in 1], and the dependence of Notch

signaling on the γ -secretase-mediated release of its intracellular domain [2, 3].

PS play important roles during embryonic development. *PS1*^{-/-} mice exhibit perinatal lethality and somitogenesis defects [4, 5], while mice deficient for both PS1 and PS2 die before embryonic day 9.5 and display early patterning defects [6]. Our investigations of PS function in neural development revealed essential roles for PS in maintenance of neural progenitor cells, differentiation and migration of postmitotic neurons and generation of radial glia [1, 4, 7–9]. PS1 also regulates neuronal migration in a non-cell-autonomous manner by controlling proliferation of meningeal fibroblasts, which in turn affects the survival of Cajal Reizus neurons, pioneer neurons that are important for proper neuronal migration [8]. PS regulate these processes primarily through the Notch signaling pathway [1–3, 7, 10]. However, it is unclear whether Notch is the key functional mediator of PS in the adult brain.

To investigate the role of PS in synaptic function in the adult cerebral cortex, which is the most relevant experimental system for the investigation of the pathogenesis of AD, we generated a viable *PS1* conditional knockout, in which expression of PS1 is selectively eliminated in excitatory pyramidal neurons of the forebrain beginning at postnatal day ~18 [11]. PS is normally expressed highly in pyramidal neurons of the cerebral cortex. This hypomorphic *PS* mutant mouse exhibits a specific but mild deficit in spatial memory [11]. Synaptic transmission and plasticity in the hippocampal CA3-CA1 synapse, however, are normal [11]. Analysis of conditional *PS*-null mice lacking both PS1 and PS2 in the postnatal forebrain revealed impairments in hippocampal memory and long-term potentiation (LTP) prior to any neuropathological changes, demonstrating a requirement for PS in normal synaptic plasticity and memory [12]. More specifically, we found a selective reduction in NMDA receptor-mediated responses and synaptic levels of NMDA receptors and α CaMKII in mutant mice. Furthermore, in the absence of PS, levels of CBP and transcription of CREB/CBP target genes are reduced [12, 13], even though subsequently we found that CREB-mediated transcription is regulated indirectly by PS [14]. Strikingly, *PS* cDKO mice develop in an age-dependent manner synaptic, dendritic and neuronal degeneration with accompanying astrogliosis and hyperphosphorylation of tau, demonstrating an essential role for PS in neuronal survival [12, 15]. Furthermore, PS promote memory and neuronal survival in a γ -secretase-dependent manner, as conditional inactivation of nicastrin, another component of the γ -secretase

complex, in the adult cerebral cortex similarly resulted in progressive memory impairment and neurodegeneration [16]. Based on these in vivo findings and a large number of reports on the effects of FAD-linked mutations in culture and in vitro systems as well as in *C. elegans*, we proposed that PS mutations may cause dementia and neurodegeneration in AD via a partial loss-of-function mechanism [17]. The fact that synaptic impairments precede progressive neurodegeneration suggests that synaptic dysfunction caused by loss of PS function promotes subsequent neuronal degeneration.

To determine the precise synaptic site of PS function, we performed a systematic genetic analysis through the restriction of PS inactivation to hippocampal CA1 or CA3 neurons [18]. This strategy permitted analysis of the effects of PS inactivation in either presynaptic or postsynaptic neurons of the Schaeffer collateral pathway. We found that LTP induced by theta burst stimulation is decreased after presynaptic but not postsynaptic deletion of PS. Moreover, presynaptic but not postsynaptic inactivation of PS impairs short-term plasticity and synaptic facilitation. The probability of evoked glutamate release, measured with the open-channel NMDA receptor antagonist MK-801, is reduced by presynaptic inactivation of PS. Strikingly, depletion of calcium internal stores by thapsigargin or inhibition of calcium release from these stores by ryanodine receptor inhibitors mimics and occludes the effects of presynaptic PS inactivation. Collectively, our genetic and electrophysiological studies demonstrate that loss of PS function impairs LTP induction and glutamatergic neurotransmitter release by a presynaptic mechanism. These findings, which distinguish unequivocally between presynaptic and postsynaptic functions of PS, raise the possibility that presynaptic mechanisms may play a primary role in AD pathophysiology. In support of this hypothesis, PS are localized to presynaptic terminals [18], and amyloid precursor protein C-terminal fragments, precursors of A β , accumulate in presynaptic terminals of *PS1* conditional knockout mice [19].

Parkinson's disease (PD) is the most common movement disorder characterized by resting tremor, rigidity and bradykinesia. These clinical features are thought to result from reduced dopaminergic input to the striatum, which is caused by the loss of dopaminergic neurons in the substantia nigra. The occurrence of PD is largely sporadic, but clinical syndromes resembling sporadic PD have been linked to mutations in at least 5 distinct genes (*α -synuclein*, *parkin*, *DJ-1*, *PINK1* and *LRRK2*). The recessive inheritance mode of the mutations and the existence of large deletions in the *parkin*, *DJ-1* and *PINK1*

genes indicate a loss-of-function pathogenic mechanism. Parkin can function as an E3 ubiquitin ligase [20–23] or as a transcription regulator [24]. DJ-1 itself can be oxidized, thus protecting cells from oxidative damage [25, 26]. PINK1 is a protein kinase localized in the mitochondrion and other subcellular compartments [27–30]. Our previous generation and multidisciplinary analysis of *parkin*^{-/-}, *DJ-1*^{-/-} and *PINK1*^{-/-} mice have demonstrated that each of these gene products is required for normal dopaminergic function and evoked dopamine release in nigrostriatal terminals [31–34]. Furthermore, inactivation of each or all three of these recessive PD genes does not cause dopaminergic neurodegeneration [31, 33–35]. In addition, our studies have further shown that loss of PINK1 or Parkin impairs mitochondrial function [36, 37]. A role for PINK1 and Parkin in mitochondria is also supported by genetic studies in *Drosophila* [38–42]. Thus, two converging cellular pathogenic mechanisms have emerged from genetic studies of recessive parkinsonism

[43]. Specifically, presynaptic dopaminergic dysfunction in evoked release may be a central pathogenic precursor before leading to frank dopaminergic neuron loss in PD.

In summary, our genetic approaches to the studies of AD and PD [44] have uncovered a novel pathogenic mechanism, suggesting that defects in presynaptic neurotransmitter release may represent a convergent mechanism leading to neurodegeneration in affected circuits in AD and PD. Therapeutic strategies directed toward restoring normal neurotransmitter release may be effective in combating circuit dysfunction and neurodegeneration in these disorders.

Acknowledgements

The author would like to thank Adair Swain for assistance. This work was supported by grants from the NINDS.

References

- Wines-Samuelson M, Shen J: Presenilins in the developing, adult, and aging cerebral cortex. *Neuroscientist* 2005;11:441–451.
- De Strooper B, Annaert W, Cupers P, Saftig P, Craessaerts K, Mumm JS, Schroeter EH, Schrijvers V, Wolfe MS, Ray WJ, Goate A, Kopan R: A presenilin-1-dependent gamma-secretase-like protease mediates release of Notch intracellular domain. *Nature* 1999;398:518–522.
- Song W, Nadeau P, Yuan M, Yang X, Shen J, Yankner BA: Proteolytic release and nuclear translocation of Notch-1 are induced by presenilin-1 and impaired by pathogenic presenilin-1 mutations. *Proc Natl Acad Sci USA* 1999;96:6959–6963.
- Shen J, Bronson RT, Chen DF, Xia W, Selkoe DJ, Tonegawa S: Skeletal and CNS defects in Presenilin-1-deficient mice. *Cell* 1997;89:629–639.
- Wong PC, Zheng H, Chen H, Becher MW, Sirinathsinghji DJ, Trumbauer ME, Chen HY, Price DL, Van der Ploeg LH, Sisodia SS: Presenilin 1 is required for Notch1 and Dll1 expression in the paraxial mesoderm. *Nature* 1997;387:288–292.
- Donoviel DB, Hadjantonakis AK, Ikeda M, Zheng H, Hyslop PS, Bernstein A: Mice lacking both presenilin genes exhibit early embryonic patterning defects. *Genes Dev* 1999;13:2801–2810.
- Handler M, Yang X, Shen J: Presenilin-1 regulates neuronal differentiation during neurogenesis. *Development* 2000;127:2593–2606.
- Wines-Samuelson M, Handler M, Shen J: Role of presenilin-1 in cortical lamination and survival of Cajal-Retzius neurons. *Dev Biol* 2005;277:332–346.
- Kim WY, Shen J: Presenilins are required for maintenance of neural stem cells in the developing brain. *Mol Neurodegener* 2008;3:2.
- Yang X, Klein R, Tian X, Cheng HT, Kopan R, Shen J: Notch activation induces apoptosis in neural progenitor cells through a p53-dependent pathway. *Dev Biol* 2004;269:81–94.
- Yu H, Saura CA, Choi SY, Sun LD, Yang X, Handler M, Kawarabayashi T, Younkin L, Fedeles B, Wilson MA, Younkin S, Kandel ER, Kirkwood A, Shen J: APP processing and synaptic plasticity in presenilin-1 conditional knockout mice. *Neuron* 2001;31:713–726.
- Saura CA, Choi SY, Beglopoulos V, Malkani S, Zhang D, Shankaranarayana Rao BS, Chattarji S, Kelleher RJ 3rd, Kandel ER, Duff K, Kirkwood A, Shen J: Loss of presenilin function causes impairments of memory and synaptic plasticity followed by age-dependent neurodegeneration. *Neuron* 2004;42:23–36.
- Beglopoulos V, Shen J: Regulation of CRE-dependent transcription by presenilins: prospects for therapy of Alzheimer's disease. *Trends Pharmacol Sci* 2006;27:33–40.
- Watanabe H, Smith MJ, Heilig E, Beglopoulos V, Kelleher RJ 3rd, Shen J: Indirect regulation of presenilins in CREB-mediated transcription. *J Biol Chem* 2009;284:13705–13713.
- Beglopoulos V, Sun X, Saura CA, Lemere CA, Kim RD, Shen J: Reduced beta-amyloid production and increased inflammatory responses in presenilin conditional knock-out mice. *J Biol Chem* 2004;279:46907–46914.
- Tabuchi K, Chen G, Sudhof TC, Shen J: Conditional forebrain inactivation of nicastrin causes progressive memory impairment and age-related neurodegeneration. *J Neurosci* 2009;29:7290–7301.
- Shen J, Kelleher RJ 3rd: The presenilin hypothesis of Alzheimer's disease: evidence for a loss-of-function pathogenic mechanism. *Proc Natl Acad Sci USA* 2007;104:403–409.
- Zhang C, Wu B, Beglopoulos V, Wines-Samuelson M, Zhang D, Dragatsis I, Sudhof TC, Shen J: Presenilins are essential for regulating neurotransmitter release. *Nature* 2009;460:632–636.
- Saura CA, Chen G, Malkani S, Choi SY, Takahashi RH, Zhang D, Gouras GK, Kirkwood A, Morris RG, Shen J: Conditional inactivation of presenilin 1 prevents amyloid accumulation and temporarily rescues contextual and spatial working memory impairments in amyloid precursor protein transgenic mice. *J Neurosci* 2005;25:6755–6764.
- Imai Y, Soda M, Takahashi R: Parkin suppresses unfolded protein stress-induced cell death through its E3 ubiquitin-protein ligase activity. *J Biol Chem* 2000;275:35661–35664.

- 21 Shimura H, Hattori N, Kubo S, Mizuno Y, Asakawa S, Minoshima S, Shimizu N, Iwai K, Chiba T, Tanaka K, Suzuki T: Familial Parkinson disease gene product, parkin, is a ubiquitin-protein ligase. *Nat Genet* 2000;25: 302–305.
- 22 Shimura H, Schlossmacher MG, Hattori N, Frosch MP, Trockenbacher A, Schneider R, Mizuno Y, Kosik KS, Selkoe DJ: Ubiquitination of a new form of alpha-synuclein by parkin from human brain: implications for Parkinson's disease. *Science* 2001;293:263–269.
- 23 Zhang Y, Gao J, Chung KK, Huang H, Dawson VL, Dawson TM: Parkin functions as an E2-dependent ubiquitin-protein ligase and promotes the degradation of the synaptic vesicle-associated protein, CDCrel-1. *Proc Natl Acad Sci USA* 2000;97:13354–13359.
- 24 Alves da Costa C, Sunyach C, Giaime E, West A, Corti O, Brice A, Abou-Sleiman PM, Wood NW, Takahashi H, Goldberg MS, Shen J, Checler F: Transcriptional repression of p53 by parkin and impairment by autosomal recessive juvenile Parkinson's disease-associated mutations. *Nat Cell Biol* 2009;11: 1370–1375.
- 25 Kahle PJ, Waak J, Gasser T: DJ-1 and prevention of oxidative stress in Parkinson's disease and other age-related disorders. *Free Radic Biol Med* 2009;47:1354–13561.
- 26 Mitumoto A, Nakagawa Y, Takeuchi A, Okawa K, Iwamatsu A, Takanezawa Y: Oxidized forms of peroxiredoxins and DJ-1 on two-dimensional gels increased in response to sublethal levels of paraquat. *Free Radic Res* 2001;35:301–310.
- 27 Gandhi S, Muqit MM, Stanyer L, Healy DG, Abou-Sleiman PM, Hargreaves I, Heales S, Ganguly M, Parsons L, Lees AJ, Latchman DS, Holton JL, Wood NW, Revesz T: PINK1 protein in normal human brain and Parkinson's disease. *Brain* 2006;129:1720–1731.
- 28 Pridgeon JW, Olzmann JA, Chin LS, Li L: PINK1 protects against oxidative stress by phosphorylating mitochondrial chaperone TRAP1. *PLoS Biol* 2007;5:e172.
- 29 Lin W, Kang UJ: Characterization of PINK1 processing, stability, and subcellular localization. *J Neurochem* 2008;106:464–474.
- 30 Weihofen A, Ostaszewski B, Minami Y, Selkoe DJ: Pink1 Parkinson mutations, the Cdc37/Hsp90 chaperones and Parkin all influence the maturation or subcellular distribution of Pink1. *Hum Mol Genet* 2008;17: 602–616.
- 31 Goldberg MS, Fleming SM, Palacino JJ, Cepeda C, Lam HA, Bhatnagar A, Meloni EG, Wu N, Ackerson LC, Klapstein GJ, Gajendiran M, Roth BL, Chesselet MF, Maidment NT, Levine MS, Shen J: Parkin-deficient mice exhibit nigrostriatal deficits but not loss of dopaminergic neurons. *J Biol Chem* 2003;278:43628–43635.
- 32 Goldberg MS, Pisani A, Haburcak M, Vorthers TA, Kitada T, Costa C, Tong Y, Martella G, Tscherter A, Martins A, Bernardi G, Roth BL, Pothos EN, Calabresi P, Shen J: Nigrostriatal dopaminergic deficits and hypokinesia caused by inactivation of the familial Parkinsonism-linked gene DJ-1. *Neuron* 2005;45:489–496.
- 33 Kitada T, Pisani A, Karouani M, Haburcak M, Martella G, Tscherter A, Platania P, Wu B, Pothos EN, Shen J: Impaired dopamine release and synaptic plasticity in the striatum of parkin^{-/-} mice. *J Neurochem* 2009;110: 613–621.
- 34 Kitada T, Pisani A, Porter DR, Yamaguchi H, Tscherter A, Martella G, Bonsi P, Zhang C, Pothos EN, Shen J: Impaired dopamine release and synaptic plasticity in the striatum of PINK1-deficient mice. *Proc Natl Acad Sci USA* 2007;104:11441–11446.
- 35 Yamaguchi H, Shen J: Absence of dopaminergic neuronal degeneration and oxidative damage in aged DJ-1-deficient mice. *Mol Neurodegener* 2007;2:10.
- 36 Gautier CA, Kitada T, Shen J: Loss of PINK1 causes mitochondrial functional defects and increased sensitivity to oxidative stress. *Proc Natl Acad Sci USA* 2008;105:11364–11369.
- 37 Palacino JJ, Sagi D, Goldberg MS, Krauss S, Motz C, Wacker M, Klose J, Shen J: Mitochondrial dysfunction and oxidative damage in parkin-deficient mice. *J Biol Chem* 2004;279:18614–18622.
- 38 Clark IE, Dodson MW, Jiang C, Cao JH, Huh JR, Seol JH, Yoo SJ, Hay BA, Guo M: *Drosophila* pink1 is required for mitochondrial function and interacts genetically with parkin. *Nature* 2006;441:1162–1166.
- 39 Greene JC, Whitworth AJ, Kuo I, Andrews LA, Feany MB, Pallanck LJ: Mitochondrial pathology and apoptotic muscle degeneration in *Drosophila* parkin mutants. *Proc Natl Acad Sci USA* 2003;100:4078–4083.
- 40 Park J, Lee SB, Lee S, Kim Y, Song S, Kim S, Bae E, Kim J, Shong M, Kim JM, Chung J: Mitochondrial dysfunction in *Drosophila* PINK1 mutants is complemented by parkin. *Nature* 2006;441:1157–1161.
- 41 Pesah Y, Pham T, Burgess H, Middlebrooks B, Verstreken P, Zhou Y, Harding M, Bellen H, Mardon G: *Drosophila* parkin mutants have decreased mass and cell size and increased sensitivity to oxygen radical stress. *Development* 2004;131:2183–2194.
- 42 Yang Y, Gehrke S, Imai Y, Huang Z, Ouyang Y, Wang JW, Yang L, Beal MF, Vogel H, Lu B: Mitochondrial pathology and muscle and dopaminergic neuron degeneration caused by inactivation of *Drosophila* Pink1 is rescued by Parkin. *Proc Natl Acad Sci USA* 2006;103:10793–10798.
- 43 Shen J, Cookson MR: Mitochondria and dopamine: new insights into recessive parkinsonism. *Neuron* 2004;43:301–304.
- 44 Beglopoulos V, Shen J: Gene-targeting technologies for the study of neurological disorders. *Neuromolecular Med* 2004;6:13–30.