*John A. Grant
John F. Sarwark*

Chicago, Ill., USA

Image of the Month

This is a 13-year-old girl who presented with very mild and infrequent back pain. She played soccer and swam competitively. Her parents had noted some difficulty walking long distances and some posture changes. She had a normal neurological examination and denied bladder or bowel problems at the time of diagnosis. Exertion occasionally made the back pain, which was not radicular in nature, worse. She was offered an operation but declined. She wore a brace with somewhat cavalier regularity for 2 years, during which time she continued to swim. Finally, she noted more back pain on walking long distances and presented for surgery. At that time, she had a normal neurological examination and a normal manual muscle test performed by a physical therapist.

Because of the severity of the grade and angle of the spondylolisthesis (spondyloptosis) it was opted to offer a through-and-through fibular strut graft in addition to a bilateral postero-lateral fusion [1]. At surgery, a Gill decompression of the posterior and relatively free element at L5 was performed. The dural sac and relevant nerve roots were then identified and retracted laterally. The dural sac was held in place with K wires and a 10-mm reamer was passed over a K wire which, under fluoroscopic control, had been placed centrally through the bodies of L5 and S1 from posterior to anterior. The fibular dowel was then sized and impacted into the resultant channel and a bilateral posterolateral fusion was performed.

Reference

- 1 Bohlman H, Cook S: One-stage decompression and interbody fusion for lumbosacral spondyloptosis through a posterior approach. *J Bone Joint Surg* 1982;64:415-418.