A Hypomorphic Allele in the *FGF8* Gene Contributes to Holoprosencephaly and Is Allelic to Gonadotropin-Releasing Hormone Deficiency in Humans

R.F. Arauz^a B.D. Solomon^a D.E. Pineda-Alvarez^a A.L. Gropman^{a, b}
J.A. Parsons^c E. Roessler^a M. Muenke^a

^aMedical Genetics Branch, National Human Genome Research Institute, National Institutes of Health, Bethesda, Md.,

^bDepartment of Neurology, Children's National Medical Center, Washington, D.C., and ^cDepartment of Neurology, The Children's Hospital, Aurora, Colo., USA

Key Words

Anterior pituitary • *FGF8* • Holoprosencephaly • HPE • IHH

Abstract

Holoprosencephaly (HPE), the most common malformation of the human forebrain, may arise due to interacting genetic and environmental factors. To date, at least 12 contributory genes have been identified. Fibroblast growth factor 8 (*Fgf8*) belongs to the FGF family of genes expressed in several developmental signaling centers, including the anterior neural ridge, which is implicated in midline anomalies in mice. In humans, *FGF8* mutations have been previously reported in facial clefting and in hypogonadotropic hypogonadism, but have not been reported in patients with HPE. We screened 360 probands with HPE for sequence variations in *FGF8* using High Resolution DNA Melting (HRM) and sequenced all identified variations. Here we describe a total of 8 sequence variations in HPE patients, including a putative loss-of-function mutation in 3 members of a family with variable forms of classic HPE, and relate these findings to the phenotypes seen in other conditions.

Copyright © 2010 S. Karger AG, Basel

Holoprosencephaly is the most common congenital malformation of the human forebrain, with a prevalence of 1 in 250 conceptions and 1 in 10,000–20,000 live births [reviewed in Muenke and Beachy, 2000; Dubourg et al., 2007; Monuki, 2007; Roessler and Muenke, 2010]. HPE results from incomplete division of the forebrain early in gestation and is exquisitely sensitive to both genetic and environmental insults during this critical period. HPE is traditionally categorized by the degree of anatomical malformations, from most severe to least severe, into alobar, semilobar, lobar, and middle interhemispheric variant types [Muenke and Beachy, 2000; Cohen, 2006; Dubourg et al., 2007; Solomon et al., 2010]. Facial features in patients with HPE range from the most severe form with cyclopia (a single eye), proboscis (a nose-like structure above a single eye), and cleft lip and palate, to less severe microforms including single maxillary central incisor (SMCI), microcephaly, and hypotelorism (closely spaced eyes) but with structurally normal brains by conventional neuroimaging [Muenke and Beachy, 2000; Cohen, 2006]. Severe neurological impairment is found in virtually all patients with brain malformations [Muenke and Beachy, 2000; Dubourg et al., 2007; Lacbawan et al., 2009].

Although all potential causes of HPE are not well characterized, there is mounting evidence showing that inter-

acting environmental and genetic factors are involved, and that severe manifestations may be due to a complex interaction of etiologies [Ming and Muenke, 2002; Bendauid et al., 2006; Dubourg et al., 2007; Roessler and Muenke, 2010].

Human *FGF8* is a member of the large fibroblast growth factor (FGF) family and maps to chromosomal region 10q24 [Lorenzi et al., 1995; White et al., 1995; Payson et al., 1996; Yoshiura et al., 1997]. *Fgf8* contains 6 exons and alternative splicing can result in 4 possible coding regions for the amino-terminus [Gemel et al., 1996; Olsen et al., 2006]. Vertebrate *Fgf8* is involved in a variety of activities including mitotic induction and cell survival, and has been implicated in a broad range of biological processes, including cell growth, tissue repair, and tumor growth invasion [Yoshiura et al., 1997]. The temporal and spatial expression pattern of *Fgf8* suggests its key involvement in embryonic development. In support of this, animal models show involvement of *Fgf8* in limb development, gastrulation, left-right axis determination, and organ morphogenesis [Ohuchi et al., 1994; Johnson and Tabin, 1997; Meyers and Martin, 1999].

Recent studies have shed light onto the conserved vertebrate process by which secreted factors produced in patterning centers regulate early organ formation such as the developing forebrain [Rubenstein et al., 1998; Storm et al., 2006; Monuki, 2007; Fernandes and Hébert, 2008]. Specifically, in telencephalic specification multiple patterning centers contribute to the development of the midline of the forebrain, an axis essential for the division of the cerebral hemispheres that fails in the HPE sequence. Communication between these centers coordinates telencephalic development [Monuki, 2007; Fernandes and Hébert, 2008]. Through these patterning centers, *Fgf8* signaling affects the telencephalon at several levels. *Fgf8* disruption in both mouse and zebrafish results in a small telencephalon [Meyers et al., 1998]. The rostral patterning center expresses multiple *Fgf* genes, including *Fgf8* [Maruoka et al., 1998]. *Fgf8* expression in the rostral patterning center aids proliferation and inhibits differentiation and dorsal telencephalic fates [Xuan et al., 1995]. The dorsal patterning center, a center known to express *Zic2* (mutations in *ZIC2* cause HPE in humans), also expresses *Bmp* and *Wnt* genes, and directs development of cortical hem and dorsocaudal structures [Brown et al., 1998; Grove et al., 1998; Galceran et al., 2000]. Finally, the ventral patterning center expresses *Shh* (again, mutations in which cause HPE in humans) [Crossley and Martin, 1995; Shimamura et al., 1995; Roessler et al., 1996, 2009]. *Shh* interacts with *Fgf8*, and regulation of *Shh* expression and development of ventral telen-

cephalon is known to be dependent on FGF signaling in zebrafish and other vertebrates [Shinya et al., 2001; Garel et al., 2003; Lupo et al., 2006; Storm et al., 2006; reviewed in Hoch et al., 2009]. These observations provide evidence for cross-linked communication between patterning centers and for the obligatory role of the FGF8 in communication between ventral and dorsal centers. Therefore, the testable prediction would be to measure the actual presence or absence of *FGF8* variants among our HPE cases.

While mutations in *FGF8* have not previously been studied in human HPE, heterozygous mutations in *FGF8* can be found in other diseases in humans [reviewed in Krejci et al., 2009]. Mutations in *FGF8* were found in 6 unrelated patients with idiopathic hypogonadotropic hypogonadism (IHH) with and without anosmia [Falardeau et al., 2008; for a review, see Bianco and Kaiser, 2009]. Functional studies show that all of these mutations resulted in diminished function of various degrees. Interestingly, *Fgf8* hypomorphic mice show olfactory bulb anomalies, with reduced neuronal gonadotropin cells in the hypothalamus. Of note, one of the above patients with adult-onset IHH has an identical mutation to the one we found in HPE patients [Falardeau et al., 2008; see below].

Additionally, an *FGF8* mutation (whose likely pathogenicity was inferred by structural analysis) has also been reported in one patient with nonsyndromic cleft lip and palate [Riley et al., 2007]. Importantly, all patients described so far with *FGF8* mutations have phenotypes consistent with midline defects [Krejci et al., 2009].

As *Fgf8* appears to act in midline induction by serving as an intermediate factor in a cross-regulatory interaction between ventral and dorsal domains, we felt that *FGF8* was an excellent candidate gene for human HPE [Storm et al., 2006; Monuki, 2007; Fernandes and Hébert, 2008; Paek and Hébert, 2009]. Here we will present the use of high-resolution DNA melting (HRM), with bi-directional sequencing of variants as a robust and sensitive screening methodology to evaluate new HPE candidate genes with respect to their potential roles as primary or secondary factors in forebrain development pathologies.

Materials and Methods

The genomic organization of the *FGF8* gene (NM_033164.1, corresponding to the longest transcript: variant F) was obtained from public database programs (<http://www.ncbi.nlm.nih.gov>) and the Bioinformatics site annotation provided by <http://www.genome.ucsc.edu>. Guidelines for the naming of the sequence variants conform to the recommendations of the human nomenclature committee (www.hgvs.org/mutnomen). Oligonucleotide primers were designed using the Primer3 v. 0.4.0 and Oligo™ 6.8

Table 1. Primers for PCR amplification of the human *FGF8* gene

Exon	Forward	Reverse	Product size, bp	T _m , °C ^a
1	5'-AGCGGCGCACAGCGATTTCGGTG-3'	5'-GAGGCAGAGGACCAGCAAGTGC-3'	343	66.4
2	5'-CCGCTTTTGTCTCCCACAG-3'	5'-GACCTCAGGAGGGGTGCTAC-3'	245	56.4
3	5'-CCAGGGGATGGATGTTTCG-3'	5'-CCACCTGTTGGGAGACAC-3'	303	64
4	5'-CCTGGGCGGAGTAGCATTA-3'	5'-TCCCACAAGCTACCTTCAGC-3'	264	65
5	5'-GCCTTCTGCCTACCTTGTG-3'	5'-ATCTGCCAATAGCCATCCTG-3'	254	57
6.1	5'-GGGTGCCCTACAGGATGAG-3'	5'-ACCAGCGTGAGGTCCACTT-3'	206	63.5
6.2	5'-GGGCGGGTAGTTGAGGAAC-3'	5'-ACGGAGATTGTGCTGGAGA-3'	207	62.4
6.3	5'-GGTGAAGGCCATGTACCAG-3'	5'-GCGAGTTGTGAGGGATTAGAGA-3'	270	64.2

^aAnnealing temperature for PCR amplification.

for the 6 coding exons and exon/intron boundaries of all known isoforms, as described (table 1).

Screened Populations

Our study consists of 360 unrelated cases from our collection of probands with HPE, which encompasses the entire clinical spectrum of severity in our HPE patients. In addition, our study also includes 184 unrelated individuals acquired as de-identified samples from the Coriell Institute for Medical Research (Camden, N.J.) to serve as controls. High-resolution melting profile was performed on the 360 unrelated HPE patients and the 184 unrelated controls simultaneously. Investigation of all samples was in accordance with a National Human Genome Research Institute (NHGRI) Institutional Review Board-approved research protocol.

PCR Amplification

Roche LightCycler® 480: All exons, except exon 1 (see below), were amplified in a 7.5-μl reaction volume using 5–10 ng DNA template, 3.5 μl of Roche High-resolution melting Master (containing FastStart Taq DNA polymerase, reaction buffer, dNTP mix, and HRM dye (Roche)), 0.6 μl of magnesium chloride (Roche) and 0.3 mM of each primer. All reactions were done employing a Roche LightCycler® 480 (Roche, Ind.). PCR amplification parameters were: incubation at 95°C for 10 min followed by 45 cycles of denaturation at 95°C for 10 s, annealing (depending on primer, see table 1) for 15 s, and extension at 72°C at amplicon size divided for 25 s.

PCR Assay for Exon 1: Exon 1 was amplified using a 25-μl reaction volume using 5–10 ng DNA template, 0.2 mM of dNTP, 0.75 mM of each primer, 2.5 μl of 10× PCR buffer (Invitrogen), 2.5 μl of 10× Enhancer buffer (Invitrogen), 0.75 μl of magnesium sulfate (Invitrogen) and 0.2 μl Taq polymerase. Every reaction was performed using a PTC-225 thermocycler (MJ Research, Mass.). PCR cycling parameters were: 95°C for 4 min followed by 40 cycles at 95°C, annealing at 66.4°C, extension at 72°C for 30 s, and a final extension step of 72°C for 7 min.

High-Resolution DNA Melting Acquisition

High-resolution melting curve analysis was used to identify putative sequence variants in *FGF8* in 360 HPE samples and 184 normal controls using the LightCycler® 480 II system (Roche ap-

plied science, Indianapolis, Ind.). Samples in all exons, except exon 1, were melted by increasing the temperature from 65 to 95°C, while exon 1 was melted from 65 to 99°C at a programmed rate of 0.2°C/s with fluorescence acquisition of 25 times per 1°C of temperature increase.

High-Resolution DNA Melting Analysis

384 well DNA-containing sample plates were used to scan for mutations; within these plates, 16 water-containing blank samples were used as negative controls. HRM curve analysis was performed according to the manufacturer's recommendations using LightCycler® 480 Software release v.1.5.0 (Idaho Technology Inc., Idaho and Roche Applied Sciences, Ind.). Amplicons of individuals showing evidence for variants were normalized, temperature-shifted fluorescence-over-temperature plots were identified, and samples were sequenced bi-directionally to confirm the presence of sequence changes.

DNA Sequencing

Samples with melting profiles deviating from the profile of the wild type were sequenced to define the variant. Sequencing was performed bi-directionally at the core DNA Sequencing Facility, National Institute of Neurological Disorders and Stroke, NIH.

Clinical Data

In accordance with our NHGRI IRB-approved procedures, families with positive clinical or molecular findings were contacted and consented to participate in our comprehensive clinical study on HPE.

Results

Molecular Findings in HPE Patients and Controls

We find both novel and known variants in *FGF8* in both HPE patients and controls (table 2). Most of these variations occur in non-coding intronic or flanking sequences that do not have any known functional effects. In 1 HPE-affected family, we find a hypomorphic muta-

Table 2. Variants detected in *FGF8* by HRM and bi-directional sequencing. DNA mutation numbering is based on the cDNA sequence. A of ATG translation initiation codon is +1 and the initiation codon is codon 1 based on NM_033163.1 (for a review, see Krejci et al. [2009])

Exon	Nucleotide and amino acid variants	Location	Number of HPE patients	Number of controls
1	c.1-35C>T [5'UTR-35C>T, rs2735432]	5'UTR	8	3
1	c.1-6G>T [5'UTR-6G>T]	5'UTR	1	0
3	c.70-127G>C [IVS2-127G>C]	intronic	1	0
4	c.232C>T [p.R78C]	missense	0	1
5	c.444+19G>A [IVS5+19G>A, rs3218234]	intronic	12	4
5	c.444+30C>A [IVS5+30C>A]	intronic	1	0
5	c.444+36C>A [IVS5+36C>A]	intronic	1	0
6	c.678G>T [p.P226P]	synonymous substitution	1	0
6	c.686C>T [p.T229M]	missense	2	0

tion in a highly evolutionary conserved region of *FGF8*: c.686C>T, resulting in p.T229M (fig. 1). This mutation results in the substitution of a hydrophobic amino acid (methionine) for a hydrophilic amino acid (threonine) in a region of the molecule present in all 4 known biologically active isoforms. Of note, this specific mutation has been reported in a patient with idiopathic hypogonadotropic hypogonadism, was shown to have a deleterious functional effect using in vitro models, and is predicted to alter activity of *FGF8* in humans [Falardeau et al., 2008]. While the precise mechanism underlying its reduced activity is unknown, due to its location in the extreme carboxy-terminus of the protein, it has been recently demonstrated that factors other than direct binding to receptors likely contribute to sharp morphogenetic gradients in biological tissues [Yu et al., 2009]. By bidirectional sequencing, no coding region mutations were found in the four most common HPE-associated genes in this family (*SHH*, *ZIC2*, *SIX3*, and *TGIF*).

Clinical Findings

Three members of the family (fig. 2) tested positive for a mutation in *FGF8*: c.686C>T, resulting in p.T229M. This maternally-inherited mutation was initially found in dizygotic twin females. The first twin (II.1) had semilobar HPE, and features consistent with classic non-syndromic HPE, including microcephaly, cleft palate, seizures, diabetes insipidus (DI), and severe neurological impairment. Other than the presence of DI, she had no evidence for other endocrinological disturbances. The other twin (II.2) had an SMCI and hypotelorism. Brain MRI performed at approximately 1 year of age was described as showing subtle midline anomalies, with the presence of bilateral olfactory bulb dysplasia, but a repeat

MRI performed at 8 years of age at the NIH showed no evidence of midline abnormalities. Significantly, this child was quite intellectually gifted, supported by standardized scholastic evaluations showing that she performed several years above her chronological age in reading and mathematics. The only clinical manifestation in the twins' mother (I.1) was mild hypotelorism; she was also highly intelligent by history. Her brain MRI showed no midline or other anomalies.

Neither living family member with the mutation showed any signs of endocrinological disturbance despite a comprehensive evaluation. Specifically, there were no signs of hypogonadotropic hypogonadism by history, serum screening, or by radiologic anatomical study.

Discussion

We tested 360 individuals with HPE for mutations in *FGF8*. Surprisingly, we detected only 1 family with a putative loss-of-function mutation in this gene. The low incidence of mutations of the *FGF8*-coding region argues against mutations in this gene as a common genetic factor to explain the occurrence of midline forebrain anomalies. However, we cannot exclude the possibility that unscreened genetic elements could play a role in human disease, for example, those that control the levels or tissue distribution of *FGF8* transcripts [Komisarczuk et al., 2009], or its various isoforms, which were not the focus of this study. Nor can we exclude the ability of other FGFs to compensate for any putative diminished *FGF8* function.

Several additional conclusions can be drawn from this study. First of all, this family simultaneously highlights

Fig. 1. HRM detection of variant c.686C>T, resulting in p.T229M in exon 6 of *FGF8*. **a** Temperature melt profile of the variant (in magenta) is distinct from wild types. **b** Differential plot illustrates the variant (in magenta) is distinct from wild types. **c** Representative chromatogram shows the DNA sequence variation detected in all 3 members of the family.

Table 3. Putative disease-related *FGF8* variants

Patients	Nucleotide change	Amino acid substitution	Diagnosis	Associated phenotypes	Reference
1a	c.686C>T	p.T229M	HPE	semilobar HPE, cleft palate, and seizures	this report
1b	c.686C>T	p.T229M	microform HPE	microform HPE	this report
1c	c.686C>T	p.T229M	microform HPE	very subtle microform HPE	this report
2	c.686C>T	p.T229M	AHH	type II diabetes	Falardeau et al. [2008]
3	c.40C>A	p.H14N	nIHH	high arched palate, osteoporosis, fractures	Falardeau et al. [2008]
4	c.77C>T	p.P26L	KS	none	Falardeau et al. [2008]
5	c.118T>C	p.F40L	nIHH	none	Falardeau et al. [2008]
6	c.298A>G	p.K100E	nIHH	microphallus	Falardeau et al. [2008]
7	c.379C>G	p.R127G	KS	cleft lip and palate, osteoporosis, hearing loss, hypotelorism, flat nasal bridge, camptodactyly, hyperlaxity	Falardeau et al. [2008]
8	c.217G>C	D73H	NS CLP	BL CLP	Riley et al. [2007]

Abbreviations: AHH, adult-onset hypogonadotropic hypogonadism; BL, bilateral; KS, Kallmann syndrome; nIHH, normosmic idiopathic hypogonadotropic hypogonadism; NS CLP, nonsyndromic cleft lip and palate.

Fig. 2. Pedigree of family segregating *FGF8* mutation. As the legend indicates, individuals with the right side shaded blue segregate the c.686C>T mutation in *FGF8*. Individuals I.1 and II.2 display microform features, indicated by yellow shading, while patient II.1 had semilobar HPE, indicated by black shading. Individuals II.3 and II.4 were not tested, but by history and photo review had no findings consistent with microform HPE or other related issues.

the importance and limitation of functional studies, both from the perspective of understanding the genetic pathogenesis of HPE, and in terms of counseling families in the clinical setting. The p.T229M mutation found in this family and in an apparently unrelated family with IHH occurs in an evolutionarily conserved region of *FGF8* suggesting its biological relevance. Evidence from animal models and cell-based assays shows that this mutation has a modest deleterious functional effect. Specifically, this heterozygous mutation is predicted to alter FGF8 ac-

tivity in humans [Falardeau et al., 2008]. However, it is reasonable to infer that additional genetic or environmental factors, presently unknown, would be required to produce severe phenotypes. Interestingly, several families with IHH have been described that segregate more than one type of mutation for this condition, and, therefore, an oligogenic model has been proposed [Pitteloud et al., 2007; Cole et al., 2008] that resembles the ‘multiple hit’ model of HPE in many respects [Ming and Muenke, 2002].

The family reported here shows that the same mutation can result in highly variable clinical consequences; this mutation must therefore not solely determine phenotypic outcome. This highly variable intrafamilial expressivity is typical in other families with mutations in HPE-related genes [Roessler et al., 1996; Solomon et al., 2009]. This inter- and intrafamilial variable expressivity is typical for HPE and other complex traits including IHH [Pitteloud et al., 2007; Cole et al., 2008]. This argues that *FGF8* may act as one of numerous factors in the model of the pathogenesis of HPE resulting from interacting genetic and environmental influences [Ming and Muenke, 2002; Roessler and Muenke, 2010].

Second, previous studies have shown that patients with mutations in *FGF8* are associated with midline defects such as cleft lip and palate and central pituitary/hypothalamic dysfunction. Using our cohort of patients with HPE, we show that a mutation in *FGF8* may contribute to a midline defect at the most severe end of the phenotypic spectrum (table 3). However, we cannot rule out

the possibility that other genetic and/or environmental factors are equally relevant to the pathologies seen in this family.

Our results suggest that mutations in an evolutionarily conserved region of *FGF8* can affect human midline and forebrain development. This report, in conjunction with previous reports, provides evidence that *FGF8* is involved in midline patterning in humans and that failure of its proper expression can infrequently result in human diseases. Despite evidence for the key role of *FGF8* in the cascade involved in midline forebrain patterning, this study revealed a low prevalence of mutations in *FGF8* in our cohort of patients with HPE. Biologically, *FGF8* appears to play important roles in forebrain development, but may only act to contribute to frank disease in the presence of other, currently undetermined factors. Fur-

thermore, the fact that the family described here did not show any signs of IHH indicates that other factors are almost certainly yet to be identified. Future directions should be centered on screening more patients with a variety of midline defects for mutations in *FGF8* in order to determine the entire spectrum of phenotypic consequences. In conjunction, it will be important to continue to explore the presence of other contributory biological factors in these individuals.

Acknowledgements

We are extremely grateful to the family presented in this report and to all the other research participants. This research was supported by the Intramural Research Program of the National Human Genome Research Institute, National Institutes of Health.

References

- Bendavid C, Haddad BR, Griffin A, Huizing M, Dubourg C, et al: Multicolour fish and quantitative PCR can detect submicroscopic deletions in holoprosencephaly patients with a normal karyotype. *J Med Genet* 43:496–500 (2006).
- Bianco SDC, Kaiser UB: The genetic and molecular basis of idiopathic hypogonadotropic hypogonadism. *Nat Rev Endocrinol* 5:569–576 (2009).
- Brown SA, Warburton D, Brown LY, Yu C, Roeder ER, et al: Holoprosencephaly due to mutations in *ZIC2*, a homologue of drosophila odd-paired. *Nat Genet* 20:180–183 (1998).
- Cohen M: Holoprosencephaly: Clinical, anatomic, and molecular dimensions. *Birth Defects Res A Clin Mol Teratol* 76:658–673 (2006).
- Cole LW, Sidis Y, Zhang C, Quinton R, Plummer L, et al: Mutations in Prokineticin 2 and Prokineticin receptor 2 genes in human gonadotrophin-releasing hormone deficiency: molecular genetics and clinical spectrum. *J Clin Endocrinol Metab* 93:3551–3559 (2008).
- Crossley PH, Martin GR: The mouse *Fgf8* gene encodes a family of polypeptides and is expressed in regions that direct outgrowth and patterning in the developing embryo. *Development* 121:439–451 (1995).
- Dubourg C, Bendavid C, Pasquier L, Henry C, Odent S, David V: Holoprosencephaly. *Orphanet J Rare Dis* 2:8 (2007).
- Falardeau J, Chung WCJ, Beenken A, Raivio T, Plummer L, et al: Decreased *FGF8* signaling causes deficiency of gonadotropin-releasing hormone in humans and mice. *J Clin Invest* 118:2822–2831 (2008).
- Fernandes M, Hébert JM: The ups and downs of holoprosencephaly: dorsal versus ventral patterning forces. *Clin Genet* 73:413–423 (2008).
- Galceran J, Miyashita-Lin EM, Devaney E, Rubenstein JLR, Grosschedl R: Hippocampus development and generation of dentate gyrus granule cells is regulated by LEF1. *Development* 127:469–482 (2000).
- Garel S, Huffman KJ, Rubenstein JLR: Molecular regionalization of the neocortex is disrupted in *Fgf8* hypomorphic mutants. *Development* 130:1903–1914 (2003).
- Gemell J, Gorry M, Ehrlich GD, MacArthur CA: Structure and sequence of human *FGF8*. *Genomics* 35:253–257 (1996).
- Grove EA, Tole S, Limon J, Yip LW, Ragsdale CW: The hem of the embryonic cerebral cortex is defined by the expression of multiple *Wnt* genes and is compromised in *Gli3*-deficient mice. *Development* 125:2315–2325 (1998).
- Hoch RV, Rubenstein JLR, Pleasure S: Genes and signaling events that establish regional patterning of the mammalian forebrain. *Sem Cell Dev Biol* 20:378–386 (2009).
- Johnson RL, Tabin CJ: Molecular models for vertebrate limb development. *Cell* 90:979–990 (1997).
- Komisarczuk AZ, Kawakami K, Becker TS: Cis-regulation and chromosomal rearrangement of the *fgf8* locus after the teleost/terapod split. *Dev Biol* 336:301–312 (2009).
- Krejci P, Prochazkova J, Bryja V, Kozubik A, Wilcox WR: Molecular pathology of the fibroblast growth factor family. *Hum Mut* 30:1245–1255 (2009).
- Lacabawan F, Solomon BD, Roessler E, El-Jaick K, Domené S, et al: Clinical spectrum of *SIX3*-associated mutations in holoprosencephaly: correlation between genotype, phenotype and function. *J Med Genet* 46:389–398 (2009).
- Lorenzi MV, Long JE, Miki T, Aaronson SA: Expression cloning, developmental and chromosomal localization of fibroblast growth factor-8. *Oncogene* 10:2051–2055 (1995).
- Lupo G, Harris WA, Lewis KE: Mechanisms of ventral patterning in the vertebrate nervous system. *Nature Rev Neurosci* 7:103–114 (2006).
- Maruoka Y, Ohbayashi N, Hoshikawa M, Itoh N, Hogan BLM, Furuta Y: Comparison of the expression of three highly related genes, *Fgf8*, *Fgf17* and *Fgf18*, in the mouse embryo. *Mech Dev* 74:175–177 (1998).
- Meyers EN, Martin GR: Differences in left-right axis pathways in mouse and chick: Functions of *FGF8* and *SHH*. *Science* 285:403–406 (1999).
- Meyers EN, Lewandoski M, Martin GR: An *Fgf8* mutant allelic series generated by Cre- and Fbp-mediated recombination. *Nat Genet* 18:136–141 (1998).
- Ming JE, Muenke M: Multiple hits during early embryonic development: Digenic diseases and holoprosencephaly. *Am J Hum Genet* 17:1017–1032 (2002).
- Monuki ES: The morphogen signaling network in forebrain development and holoprosencephaly. *J Neuropathol Exp Neurol* 66:566–575 (2007).
- Muenke M, Beachy PA: Genetics of ventral forebrain development and holoprosencephaly. *Curr Opin Genet Dev* 10:262–269 (2000).

- Ohuchi H, Yoshioka H, Tanaka A, Kawakami Y, Nohno T, Noji S: Involvement of androgen-induced growth factor (*FGF-8*) gene in mouse embryogenesis and morphogenesis. *Biochem Biophys Res Commun* 204:882–888 (1994).
- Olsen SK, Li JY, Bromleigh C, Eliseenkova AV, Ibrahimi OA, et al: Structural basis by which alternative splicing modulates the organizer activity of *FGF8* in the brain. *Genes Dev* 20:185–198 (2006).
- Paek H, Hébert JM: *FGF* signaling is strictly required to maintain early telencephalic precursor cell survival. *Development* 136:2457–2465 (2009).
- Payson RA, Wu J, Liu Y, Chiu MI: The human *FGF-8* localizes on chromosome 10q24 and is subjected to induction by androgen in breast cancer cells. *Oncogene* 1:47–53 (1996).
- Pitteloud N, Quinton R, Pearce S, Raivio T, Acierno J, et al: Digenic mutations account for variable phenotypes in idiopathic hypogonadotropic hypogonadism. *J Clin Invest* 117:457–463 (2007).
- Riley BM, Mansilla MA, Ma J, Daack-Hirsch S, Maher BS, et al: Impaired *FGF* signaling contributes to cleft lip and palate. *Proc Natl Acad Sci USA* 104:4512–4517 (2007).
- Roessler E, Muenke M: The molecular genetics of holoprosencephaly. *Am J Med Genet C Semin Med Genet* 154C:52–61 (2010).
- Roessler E, Belloni E, Gudenz K, Jay P, Berta P, et al: Mutations in the human sonic hedgehog gene cause holoprosencephaly. *Nature Genetics* 14:357–360 (1996).
- Roessler E, El-Jaick KB, Dubourg C, Vélez JI, Solomon BD, et al: The mutational spectrum of holoprosencephaly-associated changes within the *SHH* gene in humans predicts loss-of-function through either key structural alterations of the ligand or its altered synthesis. *Hum Mutat* 30:E921–E935 (2009).
- Rubenstein JL, Shimamura K, Martinez S, Puelles L: Regionalization of the prosencephalic neural plate. *Ann Rev Neurosci* 21:445–477 (1998).
- Shimamura K, Hartigan DJ, Martinez S, Puelles L, Rubenstein JLR: Longitudinal organization of the anterior neural plate and neural tube. *Development* 121:3923–3933 (1995).
- Shinya M, Koshida S, Sawada A, Kuroiwa A, Takeda H: *Fgf* signalling through MAPK cascade is required for development of the subpallial telencephalon in zebrafish embryos. *Development* 128:4153–4164 (2001).
- Solomon BD, Lacbawan F, Jain M, Domené S, Roessler E, et al: A novel *SIX3* mutation segregates with holoprosencephaly in a large family. *Am J Med Genet A* 149A:919–925 (2009).
- Solomon BD, Mercier S, Vélez JI, Pineda-Alvarez DE, Wyllie A, et al: Analysis of genotype-phenotype correlations in human holoprosencephaly. *Am J Med Genet C Semin Med Genet* 154C:133–141 (2010).
- Storm EE, Garel S, Borello U, Hébert JM, Martinez S, et al: Dose-dependent functions of *Fgf8* in regulating telencephalic patterning centers. *Development* 133:1831–1844 (2006).
- White RA, Dowler LL, Angeloni SV, Pasztor LM, MacArthur CA: Assignment of *FGF8* to human chromosome 10q25–q26: mutations in *FGF8* may be responsible for some types of acrocephalosyndactyly linked to this region. *Genomics* 30:109–111 (1995).
- Xuan S, Baptista CA, Balas G, Tao W, Soares VC, Lai E: Winged helix transcription factor *BF-1* is essential for the development of the cerebral hemispheres. *Neuron* 14:1141–1152 (1995).
- Yoshiura K, Leysens NJ, Chang J, Ward D, Murray JC, Muenke M: Genomic structure, sequence, and mapping of human *FGF8* with no evidence for its role in craniosynostosis/limb defect syndromes. *Am J Med Genet* 72:354–362 (1997).
- Yu SR, Burkhardt M, Pies NM, Petrásek Z, Schlopp S, Schwille P, Brand M: *Fgf8* morphogen gradient forms by a source-sink mechanism with freely diffusing molecules. *Nature* 461:533–536 (2009).